

# Carbonyl Oxide Chemistry in Water Cluster: An Extended Computational Study

Chayanika Kashyap<sup>1</sup>, Indrani Baruah<sup>1</sup>, Shahnaz Rohman<sup>1</sup>, Sabnam Ullah<sup>1</sup>, Amlan Kalita<sup>1</sup>, Gargi Borgohain<sup>1</sup>, and Ankur Guha<sup>1</sup>

<sup>1</sup>Cotton University

May 5, 2020

## Abstract

An extended computational approach has been utilized to explore the reactions of acids with carbonyl oxide, also known as Criegee intermediate (CI). The reactions were explored inside water cluster containing 50 water molecules. All possibilities of product formation were considered. Among the considered acids, the rate of 1,4-insertion follows the order -  $\text{HCOO} < \text{HCl} < \text{HNO}_3$ . The most stable products of the reactions between the considered acids and CI have been identified.

**1. Introduction:** In atmosphere, ozone molecule occurs as open shell singlet di-radical. Ozone easily attack the double bond containing compounds and oxidized them by attacking at double bonded positions, which generates a highly reactive Criegee intermediate (CI), also known as carbonyl oxide.<sup>1-4</sup> It is found that tropospheric alkenes are the most significant source of Criegee intermediates.<sup>5-7</sup> Tropospheric alkenes are strongly localized and short chain alkenes are more usual to undergo ozonolysis. The general mechanism of alkene ozonolysis is shown below –

These produced CIs are found to have excess internal energy to undergo unimolecular processes, generally. But with acidic particles, these CIs undergo bimolecular reaction so efficiently.<sup>8-10</sup> Studies found that some acids like HCl,  $\text{H}_2\text{SO}_4$ ,  $\text{HNO}_3$ ,  $\text{HCOOH}$  etc. are embedded on the surface of stratospheric region condensing with  $\text{H}_2\text{O}$ , they are also known as polar stratospheric particles, play a major role in scavenging CIs. During these scavenging reaction with acids, atmospheric  $\text{H}_2\text{O}$  budget are remarkably influenced.<sup>11</sup>

Early experimental data showed fast reactions of CI with formic acids (rate constant,  $k_{\text{HCOOH}} = 1 \times 10^{-11} \text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ ), hydrochloric acids (rate constant,  $k_{\text{HCl}} = 4.0 \times 10^{-11} \text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ ) and nitric acids ( $k_{\text{HNO}_3} = 5.4 \times 10^{-10} \text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ ), forming good yields.<sup>12-16</sup> Their reaction rates are suggested with the help of their reaction rate co-efficient values.<sup>16-18</sup> The obtained high rate co-efficient values suggested that acids play a key role in scavenging CI in the atmosphere leading to the formation of low-volatile but highly oxidized molecules.<sup>19,20</sup> The obtained experimental data were also in agreement with theoretical values.<sup>11</sup> Vereecken performed quantum chemical calculation to find out the probable barrier less pathways of these CI-acid reactions in gaseous phase.<sup>11</sup> As it has been already discovered that water budget plays a significant role in atmosphere, so here in our study, all the possible pathways of the CI-acid reactions are studied in water cluster containing 50 water molecules using high level computational chemistry approach.


**2. Computational details:** The water cluster containing 50 molecules of water was energy minimized using TIP4P model for water by using leap module of AMBER18 package. Steepest descent method was used for energy minimization and conjugate gradient method to discard any unfavourable interactions. The

energy minimized water cluster was used for all other *abinitio* calculations. The gradient and hessian calculations were performed using M06-2X/6-311+G\* level of theory.<sup>21</sup> We have used two layer ONIOM<sup>22</sup> calculations by using CCSD/6-311++G\*\* for the reactants and M06-2X/6-311+G\* for the surrounding water molecules. Intermediates were characterized by all real values of the hessian matrix while transition states were characterized by one imaginary value of the hessian matrix. Open shell species were treated with unrestricted formalism. Unless otherwise noted, spin contamination values were negligibly small, < 0.5%. All these calculations were performed using Gaussian 16 suite of program.<sup>23</sup>

### 3. Result and discussion


#### 3.1 Water Cluster

Fig 1 shows the TIP4P minimum energy structure of water cluster containing 50 water molecules. This cluster has been used to simulate all the reaction pathways discussed below.



**Fig 1 .** TIP4P energy minimized structure of water cluster containing 50 water molecules.

**3.2 Formation of Criegee Intermediate (CI):** Carbonyl oxides or CIs are those atmospheric intermediate which significantly act as non-photolytic source of many tropospheric reaction processes. Johnson *et.al* proved that the carbonyl oxides are produced via alkene ozonolysis having significant amount of internal energy to undergo either unimolecular or bimolecular reactions depending upon their stability.<sup>20</sup> As tropospheric abundances of alkenes and ozone are relatively higher, it is worthwhile to study the formation of CI through the reaction between  $C_2H_4$  with  $O_3$ . Fig 2 shows the reaction energetic.




**Fig 2** . Energetic for the formation of CI by the reaction of  $C_2H_4$  with  $O_3$ . The optimized geometries of the intermediates and transition states were also shown. Bond lengths are in Å.

CI-formation takes place via ozonolysis of  $C_2H_4$ . First step is 1,3-dipolar cycloaddition of ozone to the double bond of ethane leading to primary ozonide or 1<sup>st</sup> molozonide, **1** . The reaction is found to be exergonic by 75.3 kcal/mol. In 2<sup>nd</sup> step, the carbonyls (as 1,3-compounds) again undergo 1,3-cycloaddition, leading to the formation of secondary ozonide **2** (more stable than primary ozonide). The 3<sup>rd</sup> step is the breaking of C-O bonds to generate the Criegee intermediate (carbonyl oxides) and  $H_2CO$  which involves a barrier of 25.1 kcal/mol. Overall, the formation of CI is exergonic suggesting the high rate of the reaction between  $C_2H_4$  and  $O_3$ .

We then turned our attention to investigate all the possible reaction channels of some acids with CI. The choice of the acids namely, hydrochloric acid, formic acid and nitric acid for the reaction with CI is stimulated by their high abundance in atmosphere.


**3.3 Reactions between hydrochloric acids and CI** : For the co-reactant HCl, two reaction channels are considered one involving the attack of H atom of HCl to the O atom of CI while the other one involves concomitant attack of H of HCl to the terminal O of CI and Cl of HCl to the C atom of CI resulting in the formation of a cyclic transition state **TS-2** (Fig 3). The energy profile diagram is shown in Fig 3.



**Fig 3 .** Energetic of the reaction between Cl and HCl in water cluster.

The first reaction route is normal proton abstraction by Cl from HCl which proceed via **TS1** with a barrier of 13.56 kcal/mol and generates a chloride radical, which can further initiate chain reactions. The second reaction channel involves 1,3-insertion of HCl to Cl which forms a 5-membered **TS2** with a barrier of 17.75 kcal/mol. This **TS2** gives a relevant **INT** ( $\text{ClCH}_2\text{OOH}$ ) (chlorinated methylhydroperoxide) or thermally fragmented into chlorinated methoxy and hydroxyl radicals (**PROD-1**). **INT** may lead to the most stable products chlorinated methanal and water (**PROD-3**) via a cyclic **TS3**. The energy barrier for this process is 19.1 kcal/mol. The formation of **PROD-3** is highly exergonic and is the most stable product.


**3.4 Reactions between formic acids and Cl:** Four reaction pathways have been shown for the reaction between Cl and formic in this study as shown in Fig 4.



**Fig 4 .** Energetic of the reaction between Cl and HCOOH in water cluster.

Among these pathways, three reactions are initiated through a cyclic **TS1** with a barrier of 46.44 kcal/mol which is formed due to 1,4-insertion of Cl to HCOOH. The **TS1** then leads to one common intermediate hydroperoxymethylformate **INT** (HPMF). The formation of this **INT** is endergonic by 44.01 kcal/mol. This **INT** is bifurcated to **TS2** and **TS3** at the energy level of 90.06 kcal/mol and 44.96 kcal/mol respectively. The **TS2** yields highly stabilized product (**PROD-2**) while **TS3** gives a mixture of radical product (**PROD-3**). Another mixture of product (**PROD-4**) is found from **TS3** via another **TS5**. The energy barrier associated with this path is very high. Another reaction route involves proton abstraction from formic acid by Cl to generate an acid radical via **TS4** with a low barrier of 10.17 kcal/mol.

**3.5 Reactions between nitric acids and Cl:** For Nitric acid, as the co-reactant for Cl, three product formation pathways are shown in Fig 5.



**Fig 5 .** Energetic of the reaction between CI and HNO<sub>3</sub> in water cluster.

The different sets of products can be formed for the reaction between CI and HNO<sub>3</sub>. The formation of **PROD-1** is exergonic while the formation of **PROD-2** is endergonic. The attainment of these two products is sourced from same intermediate **INT** which is at the -47.0 kcal/mol in the energy profile diagram. This **INT** is attained via a transition state (**TS1**) with a barrier of 11.42 kcal/mol. The **INT** then forms two products (**PROD-1** and **PROD-2**) via two transition states (**TS2** and **TS3**). **TS3** lies very high in energy and thus the formation of **PROD-2** is less likely. Third product **PROD-3** is obtained by proton abstraction of CI from nitric acid via a transition state **TS4**. The energy barrier for the formation of **PROD-3** is only 6.96 kcal/mol and the formation of **PROD-3** is slightly endergonic (1.99 kcal/mol). From the energy barrier values, it is evident that proton abstraction from nitric acid will be easiest having highest rate constant values which is in agreement with previous theoretical calculations in gaseous phase.<sup>11</sup>

#### 4. Conclusions

All the reaction channels of Criegee intermediate (CI) with different acids held within a water cluster having 50 water molecules have been studied using high level ONIOM method (CCSD:M06-2X). Several products have been identified with low barrier for their formation. As the atmosphere contains a large quantity of water, the reactions studied herein have a good implication towards understanding the degradation mechanism of the acids considered herein. The barrier for 1,4-insertion pathway depends on the acid. The trend of barrier for the 1,4-insertion pathway follows the order: HCOO < HCl < HNO<sub>3</sub>. This is also in tune with previous experimental values of rate constants.<sup>12-16</sup> This study provides all the possible pathways of degradation of these acids on reacting with CI.

#### Acknowledgement

A. K. G. thanks the Science and Engineering Research Board (SERB), Government of India for providing financial assistance in the form of a project (project no. ECR/2016/001466). G. B. thanks DST for financial support (DST/INSPIRE/04/2017/003051).

## References

- (1) N. A. I. Watson, J. A. Black, T. M. Stonelake, P. J. Knowles, J. M. Beames, An Extended Computational Study of Criegee Intermediate - Alcohol Reactions, *J. Phys. Chem. A.* 123 (2019) 218-229.
- (2) C. C. Womack, M.-A. Martin-Drumel, G. G. Brown, R. W. Field, M. C. McCarthy, Observation of the Simplest Criegee Intermediate  $\text{CH}_2\text{OO}$  in the Gas-Phase Ozonolysis of Ethylene. *Sci. Adv.* 1 (2015) e1400105.
- (3) R. Asatryan, J. W. Bozzelli, Formation of a Criegee Intermediate in the Low-Temperature Oxidation of Dimethyl Sulfoxide. *Phys. Chem. Chem. Phys.* 10 (2008) 1769–1780.
- (4) Y. Changa, H. Chang, J. J Lina, Kinetics of the simplest Criegee intermediate reaction with ozone studied by mid-infrared quantum cascade laser spectrometer, *Phys. Chem. Chem. Phys.* 20 (2018) 97-102.
- (5) R. Criegee, Mechanism of Ozonolysis. *Angew. Chem. Int. Ed.* 14 (1975) 745-752. (6) C. A. Taatjes, D. E. Shallcross, C. J. Percival, Research frontiers in the chemistry of Criegee intermediates and tropospheric ozonolysis, *Phys. Chem. Chem. Phys.*, 16 (2014) 1704. (7) D. L. Osborn, C. A. Taatjes, The physical chemistry of Criegee intermediates in the gas phase, *Int. Rev. Phys. Chem.* , 34 (2015) 309. (8) P.-L. Luo, C. -A. Chung, Y. -P. Lee, Rate coefficient of the reaction  $\text{CH}_2\text{OO} + \text{NO}_2$  probed with a quantum cascade laser near 11  $\mu\text{m}$ , *Phys. Chem. Chem. Phys.* 21 (2019) 17578-17583.
- (9) M. Svanberg, J. B. C. Pettersson, K. Bolton, *J. Phys. Chem. A.* 104 (2000) 5787-5798.
- (10)(a) M. A. Tolbert, *M. A. Science.* 272 (1996) 1597. (b) T. Peter, *Annu. ReV. Phys. Chem.* 48 (1997) 785. (c) J. Schreiner, C. Voigt, A. Kohlmann, F. Arnold, K. Mauersberger, N. Larsen, *Science.* 283 (1999) 968.
- (11) L. Vereecken, The Reaction of Criegee Intermediates with Acids and Enols, *Phys. Chem. Chem. Phys.* , 19 (2017) 28630.
- (12) O. Welz, A. J. Eskola, L. Sheps, B. Rotavera, J. D. Savee, A. M. Scheer, D. L. Osborn, D. Lowe, A. M. Booth, P. Xiao, M. Anwar, H. Khan, C. J. Percival, D. E. Shallcross, C. A. Taatjes, Rate Coefficients of C1 and C2 Criegee Intermediate Reactions with Formic and Acetic Acid Near the Collision Limit: Direct Kinetics Measurements and Atmospheric Implications; *Angew. Chem. Int. Ed.*, 126 (2014) 4635–4638.
- (13) M. Sipilä, T. Jokinen, T. Berndt, S. Richters, R. Makkonen, N. M. Donahue, R. L. Mauldin, T. Kurtén, P. Paasonen, N. Sarnela, M. Ehn, H. Junninen, M. P. Rissanen, J. Thornton, F. Stratmann, H. Herrmann, D. R. Worsnop, M. Kulmala, V. M. Kerminen, T. Petäjä, Reactivity of stabilized Criegee intermediates (sCIs) from isoprene and monoterpeneozonolysis toward  $\text{SO}_2$  and organic acids, *Atmos. Chem. Phys.* , 14 (2014) 12143–12153.
- (14) R. Chhantyal-Pun, M. R. McGillen, J. M. Beames, M. A. H. Khan, C. J. Percival, D. E. Shallcross, A. J. Orr-Ewing, Temperature Dependence of the Rates of Reaction of Trifluoroacetic Acid with Criegee Intermediates. *Angew. Chem. Int. Ed.* , 56 (2017) 9044–9047.
- (15) E. S. Foreman, K. M. Kapnas, C. Murray, Reactions between Criegee Intermediates and the Inorganic Acids HCl and  $\text{HNO}_3$  : Kinetics and Atmospheric Implications, *Angew. Chem. Int. Ed.*, 55 (2016) 10419-10422
- (16) L. Vereecken, H. Harder, A. Novelli, The reactions of Criegee intermediates with alkenes, ozone, and carbonyl oxides. *Phys. Chem. Chem. Phys.*, 16 (2014) 4039–4049.
- (17) L. Vereecken, H. Harder, A. Novelli, The reaction of Criegee intermediates with  $\text{NO}$ ,  $\text{RO}_2$ , and  $\text{SO}_2$ , and their fate in the atmosphere, *Phys. Chem. Chem. Phys.* , 14 (2012) 14682–14695.

- (18) M. Wang, L. Yao, J. Zheng, X. Wang, J. Chen, X. Yang, D. R. Worsnop, N. M. Donahue and L. Wang, Reactions of Atmospheric Particulate Stabilized Criegee Intermediates Lead to High-Molecular-Weight Aerosol Components. *Environ. Sci. Technol.* , 50 (2016) 5702–5710.
- (19) L. Yao, Y. Ma, L. Wang, J. Zheng, A. Khalizov, M. Chen, Y. Zhou, L. Qi, F. Cui, *Atmos. Environ.* , 94 (2014) 448–457.
- (20) D. Johnson, G. Marston, The Gas-Phase Ozonolysis of Unsaturated Volatile Organic Compounds in the Troposphere. *Chem. Soc. Rev.* 37 (2008) 699–716.
21. Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* 120 (2008) 215.
22. S. Dapprich, I. Komaromi, K. S. Byun, K. Morokuma, M. J. Frisch, *J. Mol. Struct. (Theochem)* 461-462 (1999) 1-21. (b) L. W. Chung, W. M. C. Sameera, R. Ramozzi, A. J. Page, M. Hatanaka, G. P. Petrova, T. V. Harris, X. Li, Z. Ke, F. Liu, H. Li, L. Ding, K. Morokuma, *Chem. Rev.* 115 (2015) 5678-5796.
23. Gaussian 16, Revision A.03, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2016.