

Less May Be Best: High Flow Rates Are Not Always the Answer

Jie Li¹, J Scott¹, and Jan Hau Lee²

¹Rush University

²KK Women's and Children's Hospital

August 14, 2020

References

1. Luo J, Duke T, Chisti MJ, Kepreotes E, Kalinowski V, Li J. Efficacy of high-flow nasal cannula vs standard oxygen therapy or nasal continuous positive airway pressure in children with respiratory distress: A meta-analysis. *J Pediatr.* 2019;215:199-208.
2. Li J, Jing GQ, Scott JB. Year in Review 2019: High-flow nasal cannula (HFNC) oxygen therapy for adult patients. *Respir Care.* 2020; 65(4): 545-557.
3. Lavizzari A, Veneroni C, Colnaghi M, et al. Respiratory mechanics during NCPAP and HHHFNC at equal distending pressures. *Arch Dis Child Fetal Neonatal Ed.* 2014;99(4):F315-F320.
4. Simon M, Braune S, Frings D, Wiontzek AK, Klose H, Kluge S. High-flow nasal cannula oxygen versus non-invasive ventilation in patients with acute hypoxaemic respiratory failure undergoing flexible bronchoscopy—a prospective randomised trial. *Crit Care* 2014;18(6):712. doi: 10.1186/s13054-014-0712-9.
5. Saksitthichok B, Petnak T, So-Ngern A, Boonsarngsuk V. A prospective randomized comparative study of high-flow nasal cannula oxygen and non-invasive ventilation in hypoxic patients undergoing diagnostic flexible bronchoscopy. *J Thorac Dis* 2019;11(5):1929-1939.
6. Douglas N, Ng I, Nazeem F, et al. A randomised controlled trial comparing high-flow nasal oxygen with standard management for conscious sedation during bronchoscopy. *Anaesthesia* 2018;73(2):169-176.
7. Lin Y, Zhang X, Li L, et al. High-flow nasal cannula oxygen therapy and hypoxia during gastroscopy with propofol sedation: A randomized multicenter clinical trial. *Gastrointest Endosc* 2019;90(4):591-601.
8. Teng WN, Ting CK, Wang YT, et al. High-flow nasal cannula and mandibular advancement bite block decrease hypoxic events during sedative esophagogastroduodenoscopy: A randomized clinical trial. *Biomed Res Int* 2019;4206795.
9. Riccio CA, Sarmiento S, Minhajuddin A, Nasir D, Fox AA. High-flow versus standard nasal cannula in morbidly obese patients during colonoscopy: A prospective, randomized clinical trial. *J Clin Anesth* 2019;54:19-24.
10. Nielsen KR, Ellington LE, Gray AJ, Stanberry LI, Smith LS, DiBlasi RM. Effect of high-flow nasal cannula on expiratory pressure and ventilation in infant, pediatric, and adult models. *Respir Care* 2018;63:147-57.
11. Coté CJ, Wilson S; American academy of pediatrics; American academy of pediatric dentistry. Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures. *Pediatrics.* 2019;143(6):e20191000.
12. Klotz D, Seifert V, Baumgartner J, Teufel U, Fuchs H. High-flow nasal cannula vs standard respiratory care in pediatric procedural sedation: A randomized controlled pilot trial. *Pediatr Pulmonol.* 2020;1-7.
13. Scott JB, Kaur R. Monitoring breathing frequency, pattern, and effort. *Respir Care.* 2020;65(6):793-806.
14. Neuman Y, Koslow M, Matveychuk A, Bar-Sef A, Guber A, Shitrit D. Increased hypoxemia in patients

with COPD and pulmonary hypertension undergoing bronchoscopy with biopsy. *Int J Chron Obstruct Pulmon Dis.* 2015;10:2627-2632.