

Long-term responses of benthic invertebrates to rotenone treatment

Gaute Kjærstad¹, Jo Arnekleiv¹, Gaute Velle², and Anders Gravbrøt Finstad¹

¹Norges teknisk-naturvitenskapelige universitet Vitenskapsmuseet Institutt for Naturhistorie

²NORCE Norwegian Research Centre AS

July 20, 2021

Abstract

Biological invasions are regarded as one of the largest threats to native biodiversity. The eradication of non-native parasites by culling of hosts are a controversial conservation strategy, particularly when using indiscriminate methods involving whole ecosystem collateral damage. While short-term effects are abundantly documented, long-term surveys are needed to detect potential wider ecosystem effects. Here, we report a six-year study on effects of the piscicide rotenone on invertebrate communities from a Norwegian water course using a Before-After-Control-Impact design. Kick-net samples of benthic invertebrates were collected from three lentic sampling stations and two lotic stations two to four times per year in both a control and a treated watershed. In general, only relatively minor short-term effects immediately after the treatment on species turnover, measured as temporal beta-diversity, of benthic invertebrates were observed both in lentic and lotic locations. However, the lotic fauna was temporarily severely negatively affected following a period of rotenone exposure from an upstream lake. Species turnover co-varied markedly between control and treatment locations, indicating that natural environmental variation override effects of rotenone treatment. Likewise, the abundance of invertebrate taxa varied considerably both over time and between control and treatment locations. Our study indicates minor short-term (i.e. < one month) or long-term (i.e. four years) effects of rotenone treatment on benthic invertebrates, but severe effects on the lotic fauna eight months after treatment. However, long-term effects are likely to be taxa-specific and vary depending on habitat connectivity and thus potential for re-colonization and will differ among locations and among taxa.

Long-term responses of benthic invertebrates to rotenone treatment

Running title: Rotenone effects on benthic invertebrates

Gaute Kjærstad¹ | Jo Vegar Arnekleiv¹ | Gaute Velle^{2,3} | Anders Gravbrøt Finstad¹

¹Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, Trondheim, Norway

²NORCE Norwegian Research Centre, LFI, Bergen, Norway

³Department of Biological Sciences, University of Bergen, Bergen, Norway

Correspondence

Gaute Kjærstad, Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, Trondheim, Norway.

Email: gaute.kjarstad@ntnu.no

Acknowledgements

We thank the Norwegian Environment Agency and the County Governor of Nordland for financial support. We are grateful to Mark Daverdin, Lars Rønning and Aslak D. Sjursen for assisting the field work. Antti P. Eloranta gave valuable comments on an earlier draft of the manuscript.

Abstract

Biological invasions are regarded as one of the largest threats to native biodiversity. The eradication of non-native parasites by culling of hosts are a controversial conservation strategy, particularly when using indiscriminate methods involving whole ecosystem collateral damage. While short-term effects are abundantly documented, long-term surveys are needed to detect potential wider ecosystem effects.

Here, we report a six-year study on effects of the piscicide rotenone on invertebrate communities from a Norwegian water course using a Before-After-Control-Impact design. Kick-net samples of benthic invertebrates were collected from three lentic sampling stations and two lotic stations two to four times per year in both a control and a treated watershed.

In general, only relatively minor short-term effects immediately after the treatment on species turnover, measured as temporal beta-diversity, of benthic invertebrates were observed both in lentic and lotic locations. However, the lotic fauna was temporarily severely negatively affected following a period of rotenone exposure from an upstream lake. Species turnover co-varied markedly between control and treatment locations, indicating that natural environmental variation override effects of rotenone treatment. Likewise, the abundance of invertebrate taxa varied considerably both over time and between control and treatment locations.

Our study indicates minor short-term (i.e. < one month) or long-term (i.e. four years) effects of rotenone treatment on benthic invertebrates, but severe effects on the lotic fauna eight months after treatment. However, long-term effects are likely to be taxa-specific and vary depending on habitat connectivity and thus potential for re-colonization and will differ among locations and among taxa.

Keywords

alien species, benthic invertebrates, beta diversity, piscicide, recolonization

1 | INTRODUCTION

Biological invasions are regarded as one of the largest threats to native biodiversity (Mack et al., 2000), including in freshwater (Sala et al., 2000, Caffrey et al., 2014). Alien species may display invasive properties inducing serious ecological consequences through both direct and indirect ecological interactions (Kumschick et al., 2015), as well as major economic impacts (Dittel & Epifanio, 2009). Such alien invasive species may spread rapidly and cause problems that escalate from local to regional following establishment. Early targeted eradication actions are often therefore the only effective management method. This approach may be particularly effective when species are required to move through environmental corridors, such as in freshwater ecosystems (Hulme, 2009). However, options for cost-effective and reliable measures are often limited since it may become virtually impossible to eradicate invasive aquatic species in anything but small, enclosed waterbodies (Holdich et al., 1999).

In aquatic ecosystems, one of the most widely used measures for eradication of invasive species is rotenone. This is a piscicide which is highly toxic to fish and to certain invertebrates but has low toxicity to birds and mammals (Ling, 2003). Rotenone is a botanical compound originating from plant species in the Leguminosae family and have been used for centuries as a mean of capturing fish in areas where these plants are naturally occurring (Brooks & Price, 1961, Meadows, 1973). It has also been used as a fish management tool for almost a century (Finlayson et al., 2010), either with the purpose of quantifying fish abundances, manipulating fish populations to maintain sport fisheries or treatment of rearing ponds (McClay, 2000, 2005). It is one of the few, or the only, cost-effective and widely accepted measure for eradicating populations of invasive fish. In addition, rotenone has been widely used for eradication of fish parasites through culling of host populations.

Use of rotenone is controversial. This is largely due to its indiscriminate impacts on the wider aquatic fauna. Short-term impacts on non-target organisms, like benthic invertebrates is well documented (e.g. Koksvik

& Aagaard, 1984, Melaas et al., 2001, Dalu et al., 2015). The observed effects often include reduction in the total abundance of benthic invertebrates compared to pre-treatment levels. However, there are large variations ranging from little or no reduction (Cook & Moore, 1969, Dudgeon, 1990, Bellingan et al., 2019) to up to 95% reduction (Hamilton et al., 2009). Part of the variation may be explained by taxa-specific tolerance to rotenone. Ephemeroptera, Plecoptera and Trichoptera, which are abundant in oxygen rich lotic habitats, are generally reported to be rotenone sensitive (Mangum & Madrigal, 1999, Arnekleiv et al., 2001, Eriksen et al., 2009), whereas Coleoptera, Odonata and Gastropoda, which are often abundant in lentic habitats, are considered rotenone tolerant (Chandler & Marking, 1982, Holcombe et al., 1987, Arnekleiv et al., 1997, Kjærstad & Arnekleiv, 2011). There are also species-specific differences in tolerance among closely related taxa. For example, species of gill breathing Ephemeroptera are particularly sensitive to rotenone (Arnekleiv et al. 2001). Recovery of the total benthic invertebrate density can occur within a year (Kjærstad & Arnekleiv, 2004, Pham et al., 2018). However, recovery of single taxa may potentially take several years (Arnekleiv et al., 1997, Mangum & Madrigal, 1999).

Long-term consequences of failing to act towards continuous spread of invasive fish or parasites may be severe to the ecosystem and ecosystem services. Short-term disturbances to the ecosystem are expected, and to certain degree accepted, as appropriate collateral damage. However, long-term consequences of rotenone treatment may be less accepted. It is currently a lack of studies assessing long-term ecosystem effects caused by rotenone treatments (Vinson et al., 2010). Therefore, this study focused on the long-term consequences of an intensive rotenone treatment on lotic and lentic benthic invertebrate communities. Specifically, we tested for impact of rotenone treatment on the long-term abundance and temporal turnover of key taxa by comparing time series from a rotenone treated watercourse and a nearby control watercourse.

2 | METHODS

2.1 Study sites

The Fusta (rotenone treated) and Drevja (control) water courses are situated in Northern Norway (Figure 1). The Fusta watercourse has a catchment area of 538 km². The largest and lowermost lake of the catchment, Lake Fustvatnet (38 m a.s.l.), has a surface area of 11 km² and a maximum depth of 68 m. Concentrations of total nitrogen and total phosphorus from water samples taken in August 2014 of 70 µg/L and <2 µg/L, respectively, indicate oligotrophic conditions. River Fusta, which is the outlet of Lake Fustvatnet, is 8.5 km long and drains to the sea. The river has a mean annual discharge of 34 m³ s⁻¹.

The Drevja watercourse (untreated), which is bordering the Fusta watercourse, has a catchment area of 178 km². Lake Drevvatn (47 m a.s.l.), the largest and lowermost lake in the catchment, has a surface area of 5 km² and a maximum depth of 40 m. Concentrations of total nitrogen and phosphorus from water samples taken in August 2014 of 55 µg/L and <2 µg/L, respectively, indicate oligotrophic conditions. River Drevjaelva, which is the outlet of Lake Drevvatn is 17.7 km long and drains to the sea. The river has a mean annual discharge of 12 m³ s⁻¹.

Both watercourses are surrounded by sparsely vegetated mountain areas, spruce dominated forests and scattered farmlands.

2.2 | Study design

We performed the study using a Before-After-Control-Impact design (Josefsson et al., 2020). Rotenone was used in the Fusta watercourse to eradicate the invasive parasite *Gyrodactylus salaris* through exterminating its host populations of salmonid fish. Three rotenone treatments were conducted in the watercourse, with two treatments of the River Fusta (August 2011 and 2012) and one subsequent treatment of the upstream Lake Fustvatnet (October 2012), causing the downstream River Fusta to be treated a third time. The neighboring Drevja watercourse was used as a control site (Figure 1). Sampling was conducted at two lotic and three lentic sampling stations in each watercourse during the autumn from 2011 to 2016, as well as one spring sampling in June 2013, eight months after the lake treatment. Sampling started in August 2011 before the first rotenone treatment and ended in Autumn 2016. An overview of times of sampling and rotenone

treatments is given in Table 1.

Figure 1

Riverine stations were situated in riffle areas dominated by cobbles (particle size of 64-256 mm) and pebbles (particle size 4-64 mm) and with scattered patches of river moss and alga. Two of the stations in each lake were situated in exposed areas where pebbles (particle size of 4-64 mm) dominated the substrate. One of the stations in each lake was situated in sheltered areas with a substrate dominated by finer inorganic and organic substrate (particle size < 2mm) and with little or no aquatic vegetation.

Table 1.

2.3 | Sampling and sample processing

Benthic invertebrates were collected using a kick net with a 25x25 cm frame and a mesh size of 0.25 mm. For each sample, an area of approximately 0.5 m² was covered with a sampling duration of one minute. At each combination of sampling occasion and sampling station, five replicate samples were taken. The samples were preserved in 96% ethanol before subsequent subsampling of each sample (1/10) in the laboratory. All invertebrates in the subsample were sorted and identified to the lowest possible taxonomic unit using a stereo microscope and following Engblom (1996) for Ephemeroptera, Lillehammer (1988) for Plecoptera, Rinne & Wiberg-Larsen (2017) for Trichoptera, Holmen (1987) and Nilsson & Holmen (1995) for Coleoptera and Gløer (2002) for Gastropoda. The number of specimens from the sub sample were then multiplied by 10. All specimens of the remaining sample that differed morphologically from the specimens in the sub sample, were picked out, identified to lowest possible taxon and counted.

2.4 | Rotenone treatment and fish reestablishment

The rotenone treatments of River Fusta were performed on August 18th 2011 and August 17th 2012 using a rotenone formulation CFT-Legumine with a 2.5% active gradient. In both treatments the release of rotenone took place over a 7-hour period from two stations in the river, in the upper and in the middle part (Stensli & Wist, 2014). The upper release was situated approximately 1 km upstream of our uppermost sampling station and the release in the middle part was just downstream of our lowermost sampling station. In addition, riverbanks of slow flowing areas, adjacent ponds, oxbow lakes and streams were rotenone treated with local point releases. The total use of CFT-Legumine in River Fusta was 754 and 573 L in 2011 and 2012, respectively. The water temperature was approximately 15° C during both treatments and the water flow was 12 m³s⁻¹ in 2011 and 11 m³s⁻¹ in 2012. In the lower part of River Fusta, treatment concentration of CFT-Legumine was higher than 0.5 ppm over a period of several hours, which was the desired minimum concentration, periodically reaching up to 7 ppm (Adolfson et al., 2014).

The rotenone treatment of Lake Fustvatnet took place between October 16th and October 20th 2012. The total use of CFT-Legumine in Lake Fustvatnet was 139,200 L, of which 96,600 L were used for the surface waters and 42,600 L were used for deeper areas. A CFT-Legumine formula of 3.3% rotenone was used, except for the surface layer where the treatment included 25,800 L with 2.5% rotenone (Moen & Bardal, 2014). The 3.3% formula did not contain the synergist piperonylbutoxid and the solvent N-metylpyrrolidone since the former increases the toxicity to invertebrates, and not to fish (Finlayson et al., 2010). Measurements of rotenone concentration were performed over scattered sites in the surface water and at different depths during and after the treatment. On October 21st 2012 no measurements were below 0.5 ppm CFT-Legumine, and the average concentration was 0.68 ppm. Due to low water temperatures, the breakdown of rotenone was slow. In April 2013, the average concentration of CFT-Legumine was 0.1 ppm and 0.3-0.4 ppm in the surface layers and in the deeper parts (>10m), respectively. In June 2013, the concentration was below 0.1 ppm at all depths. No rotenone was detected in mid-October 2013 (Adolfson et al., 2014).

Fish was reestablished by stocking of Atlantic salmon (*Salmo salar*) in River Fusta and Arctic charr (*Salvelinus alpinus*) and brown trout (*Salmo trutta*) in Lake Fustvatnet. Three thousand Atlantic salmon smolts were released in the river in 2013 (Lo & Holthe, 2014) followed by planting of 187,000 Atlantic salmon eggs in 2014. In both 2015 and 2016, 12,000 Atlantic salmon smolts were released, as well as 384,000 young of

the year (YOY) Atlantic salmon in 2015 and 120,00 YOY in 2016 (Holthe et al. 2018). In Lake Fustvatnet 35,000 Arctic charr, mainly 2-year old, were released during the spring of 2014, as well as 93,000 brown trout YOY (Lo & Holthe, 2014, Holthe et al., 2015).

2.5| Data analyses

Changes in taxa composition between each subsequent sampling period was measured as temporal beta-diversity index (TBI). We calculated two different TBI indices, using the percentage difference from abundance data and the Sorensen index based on presence-absence data. The percentage difference ($D_{\%diff}$) between two sampling periods (t_i and t_{i+1}) was defined as $D_{\%diff} = (B+C)/(2A+B+C)$, where A is the sum of the abundance of all taxa found in both sampling period t_i and t_{i+1} . This represents the unscaled similarity between the two sampling periods. B is the part of abundance of the common taxa that is higher in samples at t_i than at t_{i+1} and is the unscaled sum of taxa *losses* between sampling periods. C is the part of the abundance that is higher t_{i+1} than at t_i and is the unscaled sum of taxa *gains* between the two sampling periods. The equation for the Sorensen index, using presence-absence data, is the same as for the $D_{\%diff}$ outlined above (see Legendre (2019) for more details). Differences in abundances and presence-absence data among subsequent sampling occasions were tested for significance with a paired t-test. The p-values (based upon 9999 random permutations) were controlled for the family-wise error rate using Holm correction. The calculations of the TBI indices and significance tests were done in the R package “adespatial” (Dray et al., 2019).

We analyzed data from riverine and lake habitat separately. Responses to rotenone treatment is expected to vary among habitats. The abundance of each invertebrate taxon is expressed as average number of specimens per kick-sample for each sampling period at each site.

Results

Rivers

The temporal species turnover appeared to co-vary in the treated and untreated river, especially for abundance data (Figure 2). The exception was the period after the lake rotenone treatment in October 2012 to summer 2013 where benthic invertebrate abundances of the rotenone treated River Fusta changed significantly (Figure 2a, Appendix 1), mainly due to losses (Figure 2b). For presence-absence data, there was a significant dissimilarity of species occurrences in the treated river from immediately after the lake treatment in October 2012 until August 2015 (Figure 2d, Appendix 2). The species loss-rate of the treated river was at its highest from October 2012 to June 2013 (Figure 2e) and gain-rate at its highest from June 2013 to August 2013 (Figure 2f, Appendix 2). No significant changes in abundance or species occurrence were found in the untreated river (Figure 2a, b, Appendix 1 and 2).

Figure 2

The total abundance of benthic invertebrates reflects the temporal turnover indices (Figure 3a). There were no clear short-term effects on the total abundance or the abundance on most taxa following the two rotenone treatments of River Fusta (Figure 3). As an exception, the abundance of Plecoptera decreased strongly in the treated river immediately after the first and second river treatment (89% and 92%, respectively), whereas the changes were smaller in the untreated river (+17% and -37%, respectively) (figure 3c).

Figure 3

In contrast, there was a marked increase in the abundance of taxa observed in samples from both the treated and the untreated river in October 2012, followed by a pronounced decline in the treated river in June 2013 (Figure 3a). However, the observed response in total abundance masks distinct differences in response to the lake rotenone treatment of riverine benthic invertebrate taxa.

The abundance of Ephemeroptera in the treated river decreased shortly after the two river treatments (Figure 2b). This was also found in the untreated river after the first river treatment. The Ephemeroptera *Baetis rhodani* and *Baetis* sp. decreased in the treated river after both treatments, and in the untreated river

after the first treatment. Abundances of all taxa from the rivers not presented in the figures are listed in Appendix 3. For other Ephemeroptera, like *Heptagenia dalecarlica* and *Ephemerella* sp., the abundance in both rivers was relatively high shortly after the river treatments. Immediately after the lake treatment in October 2012, *B. rhodani* was the only Ephemeroptera with reduced abundance and was not recorded in the treated river. In June 2013, eight months after the lake treatment, there was a dramatic decline in the Ephemeroptera abundance in the treated river (Figure 3b) with only a few specimens present of *Siphonurus* sp., *B. rhodani* and *H. dalecarlica*. However, a decline was also found in the untreated river, but here most taxa were present in June 2013. In the subsequent years, the Ephemeroptera abundances in the treated river increased, mainly *B. rhodani* and *H. dalecarlica*. A similar increase occurred in untreated river, in addition to an increase in *B. muticus*.

There was a strong decrease in the abundance of Plecoptera in the treated river after both river treatments, whereas there was an increase in the untreated river. Taxa with reduced abundances included *Diura* sp., *Isoperla* sp. and *Leuctra* sp. Shortly after the lake treatment in October 2012, the abundance of Plecoptera in the treated river decreased with 95% compared to September 2012 (Figure 3c), and only three specimens of the genus *Nemoura* were recorded. Most taxa were present in the untreated river and with a relatively high abundance of *Amphinemura borealis*, *Diura* sp. and *Isoperla* sp. In June 2013, eight months after the lake treatment, no Plecoptera were detected in the treated river, while at the same time there was an average of more than 200 specimens of Plecoptera in the untreated river (Figure 3c). Contrary to the river treatments, where the negative effect were minor, Trichoptera abundances decreased in the treated river immediately after the lake treatment but did not change in the untreated river (Figure 3d). Eight months after the lake treatment in June 2013, there was a dramatic decrease (99%) in the abundance of Trichoptera in the treated river, and only a few specimens of *Rhyacophila nubila*, *Hydropsychidae* sp. and *Micrasema* sp. were present. A corresponding but smaller decline (41%) was observed in the untreated river.

Both rivers had high abundance of Chironomidae shortly after the lake treatment (Figure 3e). The abundance of Chironomidae co-varied considerably among rivers. The abundance of Oligochaeta in the treated river decreased shortly after the first river treatment and increased subsequent to the second river treatment and the lake treatment (Figure 3f). In June 2013, eight months after the lake treatment, the abundance of Oligochaeta was at its minimum in the treated river.

There was a low abundance of Hydrachnidia in the treated river and the taxon was absent in 2013 (Appendix 3). In the untreated river, Hydrachnidia was recorded in all sampling occasions and with the highest abundances after the treatments. The abundance of the riffle beetles (Elmidae) decreased in the treated river following the treatment. In the untreated River Drevjaelva, the taxon also decreased in abundance, albeit at a much lower rate than in the treated River Fusta.

Lakes

Temporal species turnover appeared to co-vary in the treated and untreated lakes (Figure 4). The untreated lake tended to have a greater temporal turnover, both measured as dissimilarity in taxa abundances and species occurrences throughout most of the study period (Figure 4a and 4d). The highest losses in terms of species abundance occurred in the untreated lake in August 2012 and for the treated lake in August 2015 (Figure 4b). Gains were generally highest for the treated lake (Figures 4c and 4f), especially for abundance data. No significant changes in abundance or species occurrence indices between sampling periods were found in the treated or untreated lake (Appendix 4 and 5).

Figure 4

The total abundances of benthic invertebrates in the lakes showed major variations even before the rotenone treatment, i.e., the abundances decreased by about 50% from in from August 2011 to August 2012 in both lakes (Figure 5a). In October 2012, shortly after the treatment, the total abundance in both lakes decreased, but stronger in the rotenone treated Lake Fustvatnet. In June 2013, eight months after the treatment, a minor decrease occurred in both lakes. From August 2013 and onwards the variations in abundance differed between the lakes, with higher values in the treated lake in August 2013, August and October 2014, October

2015 and August and October 2016.

Figure 5

The Ephemeroptera abundance varied over time both within and between lakes (Figure 5b). There was a high abundance in both lakes in June 2013, compared to pre-treatment abundances mainly caused by the Ephemeroptera *Siphonurus* sp. Abundances of all taxa from the lakes not presented in the figures are listed in Appendix 6. High abundances were also registered in the treated lake in August 2014 and August 2016. In October 2012, shortly after the lake treatment the abundances was lower in both lakes, compared to samples taken just before the treatment. The abundance of Ephemeroptera taxa decreased immediately after the lake treatment, especially *Centroptilum luteolum* in the treated lake and Leptophlebiidae in both the treated and untreated lakes. Ephemeroptera species with increased abundances in the treated lake after the treatment included *Centroptilum luteolum*, *Caenis horaria* and *Ephemerula vulgata*. However, the abundance of *E. vulgata* decreased strongly in August and October 2015 (73% and 71%, respectively), compared to 2014. Except from *Siphonurus* sp. there was no marked temporal change in the abundances of Ephemeroptera in the untreated lake.

The abundance of Plecoptera in the treated lake was generally low except in October 2012 just before the rotenone treatment (Figure 5c). In the untreated lake, abundances varied but were generally much higher than in the treated lake. In the treated lake shortly after the treatment in October 2012, there was a strong decrease in Plecoptera abundances (Figure 5c), with only a few specimens of the genus *Nemoura* present. The decline was much smaller in the untreated lake. In June 2013, eight months after the treatment, no Plecoptera were recorded in the treated lake, and only a few specimens were observed in the untreated lake. The genus *Capnia* had the most marked decrease in abundance in the treated lake following the treatment, whereas the abundance of *Capnia* peaked in some of the years after the treatment in the untreated lake.

For Trichoptera, the abundances in both lakes varied across sampling occasions (Figure 5d). In October 2012, shortly after the treatment, abundances decreased markedly in the treated lake but did not change in the untreated lake. However, a decreased abundance also occurred before the treatment in both lakes between August 2011 and August 2012. The highest abundances in the treated lake after the treatment were recorded in August 2013 and in August 2014. In the untreated lake, the abundances peaked in 2016. Polycentropodidae had the most pronounced decrease among Trichoptera after the treatment in the treated lake, whereas the family increased slightly during the same period in the untreated lake. A few Trichoptera taxa in the treated lake, all found at low abundances, were only recorded before or after the treatment.

The abundance of Chironomidae showed only minor changes after the treatment in both lakes (Figure 5e). In subsequent years, the abundances of Chironomidae in the treated lake varied compared to pre-treatment levels, whereas it did not change or were higher in the untreated lake.

The abundance of Oligochaeta (Figure 5f) and Hydrachnidia generally varied to a similar extent in both lakes. For Corixidae and Coleoptera, and especially of the genus *Callicorixa* and dytiscid beetles, the abundances increased in the treated Lake Fustvatnet after the treatment. No Corixidae was recorded in untreated Lake Drevvatnet, while Dytiscidae occurred sporadically.

Discussion

The immediate impact of rotenone treatments in August 2011 and 2012 was minor for benthic invertebrates in the rotenone treated River Fusta. Only a few taxa showed a marked decrease in abundance in the treated river compared to the untreated River Drevjaelva, including the Ephemeroptera *Baetis rhodani* and the Plecoptera genera *Diura* and *Isoperla*. Because the upstream Lake Fustvatnet was treated two months after the last river treatment, River Fusta also received rotenone from the lake. Still, the immediate effect of the lake rotenone treatment on benthic invertebrate community in the downstream river was minor. One exception was the Trichoptera *Rhyacophila nubila*, which occurred at high abundances in both rivers, but which disappeared from the River Fusta subsequently to the treatment of the upstream lake. All the above-mentioned taxa are known to be rotenone sensitive (Engstrom-Heg et al., 1978, Arnekleiv et al., 2001, Gladso

& Raddum, 2002) and negative impacts were expected.

The rotenone treatment of Lake Fustvatnet took place in October, just before the lake ice formed. Breakdown of rotenone was thus slowed down due to low temperatures and limited light. Consequently, the treated river was exposed to rotenone from the lake for several months. Even though the toxic effect of rotenone decreases with decreasing water temperatures (Meadows, 1973, Kjaerstad et al. 2015), the long exposure apparently had a pronounced negative effect. In June 2013, eight months after the lake treatment, we recorded the lowest abundance of benthic invertebrates in the treated river. A similar decrease did not occur in the untreated river. The benthic invertebrate community in the treated river at this time were dominated by Chironomidae, Oligochaeta, Nematoda and elmids beetles. Other taxa occurred at very low abundances and several taxa were missing, including rotenone tolerant species like the Ephemeroptera *Ephemerella*. Moreover, despite that the freshwater pearl mussel (*Margaritifera margaritifera*) is known to be rotenone tolerant (Dolmen et al., 1995), all known specimens in the treated river died during the spring of 2013 (Larsen, 2015). The temporal beta diversity confirms that the changes in taxa composition were by far highest, and significant, in the treated river from October 2012 to June 2013. This decline in benthic invertebrates was mainly due to losses both in terms of taxa abundances and presence-absence. This indicates that rivers may receive rotenone for several months following treatment of upstream lakes, which can have major impacts. Most taxa in the treated river, including rotenone tolerant taxa, decreased to very low abundances and some disappeared, like the freshwater pearl mussel.

Despite the strong negative impacts recorded in the treated river eight months after the lake rotenone treatment, recolonization was relatively fast. The highest gains in terms of species occurrence in the treated river occurred in August 2013. Already in August 2014 the total abundances exceeded pre-treatment levels. In August and October 2016, the change in taxa composition, in terms of abundance and presence-absence of taxa between sampling occasions, were rather similar for the treated and the untreated river. This indicates that the benthic fauna of the treated river had recovered to a great extent. Several taxa, like the Ephemeroptera *Heptagenia dalecarlica* and *Ephemerella*, the Plecoptera *Isoperla* and *Amphinemura borealis* and Oligochaeta reached much higher abundances in 2013, as compared to pre-treatment levels. Ephemeroptera and Plecoptera are known to be among the most abundant groups in the drift of swift-flowing temperate streams (Brittain & Eikeland, 1988). These groups likely colonized rapidly from drift in untreated tributary streams, although some may have survived as eggs. Oligochaeta may have avoided rotenone exposure by digging deeper into the substratum. It took almost four years after the last rotenone treatment before most of the taxa had recolonized. The Plecoptera *Taeniopteryx nebulosa* were among the slowest to recolonize and appeared in 2016. Also, two rotenone tolerant taxa, the Ephemeroptera *Ephemerella* and elmids beetles did not reach pre-treatment abundances in the treated river four years after the treatments.

According to Vinson et al. (2010), the overall invertebrate abundances generally return to pre-treatment levels quicker than the biodiversity and taxonomic composition measures. This is in accordance to our findings. However, the recolonization will greatly depend on the taxonomic resolution of the data. Biodiversity measured at family and genera level will recolonize fast, while it may take longer before all species to recolonize. We have identified some taxa to species level, such as Ephemeroptera, Plecoptera and Trichoptera, and some taxa to a higher taxonomic level, such as the speciose Chironomidae and Hydrachnidia. Thus, changes in abundance and recolonization patterns of many species in this study remain unknown.

A reduction of total benthic invertebrate abundance was observed both in the treated lake and untreated lake immediately after the lake treatment, but to a higher extent in the treated lake. This indicates both natural changes and a negative effect of rotenone. Taxa with clearly decreasing abundances shortly after the treatment in Lake Fustvatnet, but not in the untreated Lake Drevvatnet, included the Ephemeroptera *Centroptilum luteolum* and the Plecoptera *Capnia* sp. While *Capnia* is known to be rotenone sensitive, *C. luteolum* is quite tolerant (Kjaerstad & Arnekleiv, 2011). The reason for the seemingly high sensitivity of *C. luteolum* to rotenone in the treated lake is unknown.

The recolonization in terms of abundance of benthic invertebrates in the treated lake was fast, reaching pre-

treatment levels eight months after the treatment. A few increased in abundance after the treatment in the treated lake, but not in the untreated lake. This was particularly evident for the diving beetles (Dytiscidae). Beetles are generally known to be very rotenone tolerant (Engstrom-Heg et al., 1978). Some will also benefit from increased prey availability because of the general increase in invertebrate abundance after the recolonization. Even though most taxa increased in abundance during the years following the treatments, the Plecoptera *Capnia*, the Trichoptera Polycentropodidae and the Gastropoda *Gyraulus acronicus* decreased and had not reached their pre-treatment levels in the treated lake four years after the treatment. Considering that Gastropoda are rotenone tolerant (Holcombe et al., 1987, Arnekleiv et al., 1997), the decline of *G. acronicus* in the treated lake was surprising. Mangun & Madrigal (1999) found that of several taxa missing five years after rotenone treatments, most belonged to rotenone sensitive groups, like Ephemeroptera, Plecoptera and Trichoptera. Taxa that were tolerant to rotenone were identified as "non-missing" or "briefly missing" (Mangun & Madrigal, 1999).

Fish may play an important role in structuring benthic invertebrate communities (e.g. Weyl et al., 2010, Alexiades & Kraft, 2017). Stocking of Atlantic salmon (*S. salar*) took place in the treated river between 2013 and 2016, whereas Arctic charr (*S. alpinus*) and brown trout (*S. trutta*) were stocked in the treated lake in 2014. A few benthic invertebrate taxa did not reach pre-treatment abundance. However, if fish was the main driver, we would expect a decrease in abundance of other common taxa that are eaten by fish in the years after the treatments. Instead, many taxa increased in abundances and exceeded pre-treatment levels, indicating a relatively low predation pressure from fish in the treated river. In the treated lake, contrary to the untreated lake, we observed a strong decrease in total abundances of benthic invertebrates in August 2015. This was most likely due to predation from fish stocked during the previous year. The decrease was particularly high for Chironomidae and Ephemeroptera. Abundance of large sized and important fish prey, like the Ephemeroptera *Ephemerula vulgata*, increased to a high abundance in 2013-2014, but decreased drastically in 2015 in the treated lake, probably due to predation from fish. The species was not found before the treatment, but occurred in all sampling events after the treatment, indicating a lower predation pressure from fish after the treatment. Despite the rotenone treatments and the fish removal/fish stocking, the abundance of many taxa varied to a similar extent in the impact and control sites. As an exception, the effect was more prominent in the impacted treated river than in the untreated river after the lake treatment and subsequent months of rotenone exposure. This illustrates the importance of using control sites when performing time-series experiments.

It is striking to observe the large fluctuations in the abundance and in the presence-absence of benthic invertebrates over time of both treated and untreated localities. Even before the lake treatment, the invertebrate abundances in both lakes were reduced by approximately 50% from August 2011 to August 2012. According to Vinson et al. (2010), the accumulation curve for benthic invertebrate genera showed little inclination for flattening out during a ten-year monthly sampling of a Utah river. We did not see the same pattern in our data, perhaps due to fewer taxa present. Even though it will vary greatly with habitat type, latitude etc., a rising taxa accumulating curve would certainly be found elsewhere and shows how random the community composition may be registered at any given time.

Our study found minor short-term (i.e. < one month) effects of the river rotenone treatments, whereas the lake treatment caused major temporary negative effects on the lotic fauna eight months after the treatment. No major short- or long-term (i.e. four years) effects on lake benthic invertebrate communities were detected. Recovery time of invertebrate abundance after the lake treatment was fast whereas the recovery in terms of presence-absence of species were comparably much longer, lasting for several years. The current study underpins the importance of performing long-term studies during investigation of the effects of rotenone on benthic invertebrates.

References

Adolfson, P., Sandvik, M. & Waaler, T. (2014). Rotenonanalyser og resultater. In: Bekjempelse av *Gyrodactylus salaris* i Vefsnaregionen. (Eds: J. H. Stensli & H. Bardal). Veterinærinstituttets rapportserie 2-2014.

- Alexiades, A.V. & Kraft, C. E. (2017). Effects of stocked trout on stream invertebrate communities. *Journal of Freshwater Ecology*, 32 , 95-102. <https://doi.org/10.1080/02705060.2016.1248502>
- Arnekleiv, J. V., Dolmen, D., Aagaard, K., Bongard, T. & Hanssen, O. (1997). Rotenonbehandlingens effekt på bunndyr i Rauma og Hensvassdraget, More & Romsdal. Del I: Kvalitative undersøkelser. Viten-skapsmuseet Rapport Zoologisk Serie 1997-8.
- Arnekleiv J.V., Dolmen D. & Ronning L. (2001). Effects of rotenone treatment on mayfly drift and standing stocks in two Norwegian rivers. In: E. Dominguez (ed). Trends in Research in Ephemeroptera and Plecoptera. Kluwer Academic/Plenum Publishers.
- Bellingan, T.A., Hugo, S., Woodford, D.J., Gouws, J., Villet, M.H. & Weyl, O.L.F. (2019). Rapid recovery of macroinvertebrates in a South African stream treated with rotenone. *Hydrobiologia*, 834 , 1–11. <https://doi.org/10.1007/s10750-019-3885-z>
- Brittain, J.E. & Eikeland, T.J. (1988). Invertebrate drift – A review. *Hydrobiologia* 166 , 77-93. <https://doi.org/10.1007/BF00017485>
- Brooks, I. C. & Price, R. W. (1961). Studies on the chronic toxicity of Pro-Noxfish(r), a proprietary synergized rotenone fish-toxicant. *Toxicology and Applied Pharmacology*, 3, 49–56. [https://doi.org/10.1016/0041-008X\(61\)90007-2](https://doi.org/10.1016/0041-008X(61)90007-2)
- Caffrey J.M., Baars J.R., Barbour J.H., Boets P., Boon P. & Davenport K. (2014). Tackling invasive alien species in Europe: the top 20 issues. *Management of Biological Invasions*, 5, 1-20. <http://dx.doi.org/10.3391/mbi.2014.5.1.01>
- Chandler, J. H. & Marking, L.L. (1982). Toxicity of Rotenone to Selected Aquatic Invertebrates and Frog Larvae. *The Progressive Fish-Culturist* 44 , 78-80. [https://doi.org/10.1577/1548-8659\(1982\)44\[78:TORTSA\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1982)44[78:TORTSA]2.0.CO;2)
- Cook, S.F. jr. & Moore, R.L. (1969). The effects of rotenone treatment on the insect fauna of a California stream. *Transactions of the American Fisheries Society*, 98 , 539-544. [https://doi.org/10.1577/1548-8659\(1969\)98\[539:TEOART\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1969)98[539:TEOART]2.0.CO;2)
- Dalu, T., Wasserman, R.J., Jordaan, M., Froneman, W.P. & Weyl, O.L.F. (2015). An Assessment of the Effect of Rotenone on Selected Non-Target Aquatic Fauna. – *PloS ONE* (11) : e0142140. <https://doi.org/10.1371/journal.pone.0142140>
- Dar P.A. & Reshi Z.A. (2014). Components, processes and consequences of biotic homogenization: a review. *Contemporary Problems of Ecology* 7, 123–136. <https://doi.org/10.1134/S1995425514020103>
- Dittel, A. & Epifanio, C. (2009). Invasion biology of the Chinese mitten crab *Eriocheir sinensis* : A brief review. *Journal of Experimental Marine Biology and Ecology*, 374 , 79-92. <https://doi.org/10.1016/j.jembe.2009.04.012>
- Dolmen, D., J.V. Arnekleiv, & T. Haukebo. (1995). Rotenone tolerance in the freshwater pearl mussel *Margaritifera margaritifera* . *Nordic Journal of Freshwater Research* 70 , 1-30.
- Dray, S., Bauman, D., Blanchet, G., Borchard, D., Clappe, S., Guenard, G., ... Wagner, H.H. (2019). Adespatial: Multivariate multiscale spatial analysis. R package version 0.3-3. Retrieved from <https://cran.r-project.org/package=adespatial>
- Dudgeon D. (1990). Benthic community structure and the effect of rotenone piscicide on invertebrate drift and standing stocks in two Papua New Guinea streams. *Archives of Hydrobiology* 119 , 35–53.
- Engblom, E. 1996. Ephemeroptera, Mayflies. In: Aquatic insects of North Europe. A taxonomic handbook. Vol 1. (Ed: A. Nilsson). Apollo Books, Stenstrup.

- Engstrom-Heg R., Colesante R.T. & Silco, E. (1978). Rotenone tolerances of stream-bottom insects. *New York Fish Game Journal* 25 , 31–41.
- Eriksen, T. E., Arnekleiv, J. V. & Kjaerstad, G. (2009). Short-Term Effects on Riverine Ephemeroptera, Plecoptera, and Trichoptera of Rotenone and Aluminum Sulfate Treatment to Eradicate *Gyrodactylus salaris*. *Journal of freshwater ecology*, 24 , 507–607. <https://doi.org/10.1080/02705060.2009.9664337>
- Finlayson, B., Somer, W. L. & Vinson, M. R. (2010). Rotenone toxicity to rainbow trout and several mountain stream insects. *North American Journal of Fisheries Management* 30 , 102–111. <https://doi.org/10.1577/M09-078.1>
- Finlayson, B., R. Schnick, D. Skaar, J. Anderson, L. Demong, D. Duffield, W., ...Steinkjer, J. (2010). Planning and standard operating procedures for the use of rotenone in fish management—rotenone SOP manual. American Fisheries Society, Bethesda, Maryland.
- Gladso, J. & Raddum, G.G. (2002). Rotenone treatment of a west Norwegian river: Effects on drift of invertebrates. *Internationale Vereinigung fur Theoretische und Angewandte Limnologie: Verhandlungen*, 28 , 764–769. <https://doi.org/10.1080/03680770.2001.11901816>
- Gloer, P. (2002). Susswassergastropoden Nord- und Mitteleuropas. Bestimmungsschlüssel, Lebensweise, Verbreitung. ConchBooks.
- Hamilton, B.T., Moore, S.E., Williams T.B., Darby, N. & Vinson, M.R. (2009). Comparative effects of rotenone and antimycin on macroinvertebrate diversity in two streams in Great Basin National Park, Nevada. *N. Am. J. Fish Management* 29 , 1620–1635. <https://doi.org/10.1577/M08-178.1>
- Holdich D., Reynolds J., Souty-Grosset C. & Sibley P. (2009). A review of the ever-increasing threat to European crayfish from non-indigenous crayfish species. *Knowledge and Management of Aquatic Ecosystems*, 394–395 , 11. <https://doi.org/10.1051/kmae/2009025>
- Holmen, M. (1987). The Aquatic Adephaga (Coleoptera) of Fennoscandia and Denmark, Volume 1. Gyrinidae, Haliplidae, Hygrobiidae and Noteridae. Fauna entomologic Scandinavica 20.
- Holcombe, G. W., Phipps, G. L., Sulaiman, A. H. & Hoffman, A. D. (1987). Simultaneous multiple species testing: Acute toxicity of 13 chemicals to 12 diverse freshwater amphibian, fish, and invertebrate families. *Archives of Environmental Contamination and Toxicology* 16 , 697–710. <https://doi.org/10.1007/BF01055420>
- Holthe, E., Bjorna, T. & Lo, H. (2015). Reetableringsprosjektet i Vefsnaregionen - Arsrapport for aktiviteten i 2014. Veterinaerinstitutets rapportserie 14-2015.
- Holthe, E. Bjorna, T. & Lo, H. (2018). Reetableringsprosjektet i Vefsnaregionen. Arsrapport 2017. Veterinaerinstitutets rapportserie 11-2018.
- Hulme, P. E. (2009). Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology*, 46 , 10–18. <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- Josefsson, J., Hiron, M., Arlt, D., Auffret, A.G., Berg, A., Chevalier, M., Glimskar, A., ...Part, T. 2020. Improving scientific rigor in conservation evaluations and a plea deal for transparency on potential biases. *Conservation Letters*. 2020; 13:e12726. <https://doi.org/10.1111/conl.12726>
- Kumschick S., Gaertner M., Vila M., Essl, F., Jeschke, J.M., Pysek, P., Ricciardi, A., ...Winter, M. (2015). Ecological impacts of alien species: quantification, scope, caveats, and recommendations. *BioScience*, 65 , 55–63. <https://doi.org/10.1093/biosci/biu193>
- Kjaerstad, G. & Arnekleiv, J. V. (2004). Rotenonbehandling av elver i Rana-regionen i 2003 og 2004: effekter pa bunndyr. NTNU Vitenskapsmuseet zoologisk notat 2004-4.
- Kjaerstad, G. & Arnekleiv, J.V. (2011). Effects of Rotenone Treatment on Lotic Invertebrates. *International Review of Hydrobiology* 96 , 58–71. <https://doi.org/10.1002/iroh.201111261>

- Kjaerstad, G., Arnekleiv & Speed, J.D.M. (2015). Effects of three consecutive rotenone treatments on the benthic macroinvertebrate fauna of the River Ognå, Central Norway. *Rivers Research and Applications*. <https://doi.org/10.1002/rra.2873>
- Koksvik, J.I. & Aagaard, K. (1984). Effects of rotenone on the benthic fauna of a small eutrophic lake. *Internationale Vereinigung für Theoretische und Angewandte Limnologie: Verhandlungen*, 22 , 658-665. <https://doi.org/10.1080/03680770.1983.11897361>
- Larsen, B.M. (2015). Elvemusling i Fusta, Nordland – konsekvenser av rotenonbehandling i vassdraget og tiltak for å sikre bestanden av muslinger. NINA Rapport 1189.
- Legendre, P. (2019). A temporal beta-diversity index to identify sites that have changed in exceptional ways in space-times surveys. *Ecology and Evolution*, 9 , 3500-3514. <https://doi.org/10.1002/ece3.4984>
- Lillehammer, A. (1988.) Stoneflies (Plecoptera) of Fennoscandia and Denmark. E. J. Brill/Scandinavian Science Press Ltd. Leiden, The Netherlands. Vol. 21.
- Ling, N. (2003). Rotenone - review of its toxicity and use for fisheries management. *Science for conservation* 211.
- Lo, H. & Holthe, E. (2014). Bevaring av fiskebestander. In: Bekjempelse av *Gyrodactylus salaris* i Vefsnaregionen. (Eds: J.H. Stensli & H. Bardal). Veterinaer instituttets rapportserie 2-2014.
- Mack, R.N., Simberloff, D., Mark Lonsdale, W., Evans, H., Clout, M. and Bazzaz, F.A. (2000). *Biotic invasions: causes, epidemiology, global consequences, and control. Ecological Applications*, 10 , 689-710. [https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEGC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEGC]2.0.CO;2)
- Mangum, M.A. & Madrigal, J.L. (1999). Rotenone Effects on Aquatic Macroinvertebrates of the Strawberry River, Utah: A Five-Year Summary. *Journal of Freshwater Ecology* 14 , 125-135. <https://doi.org/10.1080/02705060.1999.9663661>
- Meadows, B. S. (1973). Toxicity of rotenone to some species of coarse fish and invertebrates. *Journal of Fish Biology*, 5 , 155– 163. <https://doi.org/10.1111/j.1095-8649.1973.tb04444.x>
- Melaas, C.L., Zimmer, K.D., Butler, M. G. & Hanson, M.A. (2001). Effects of rotenone on aquatic invertebrate communities in prairie wetlands. *Hydrobiologia* 459 , 177-186. <https://doi.org/10.1023/A:1012514124430>
- Moen, A. & Bardal, H. (2014). Innsjøbehandlingen. In: Bekjempelse av *Gyrodactylus salaris* i Vefsnaregionen. (Eds: J. H. Stensli & H. Bardal). Veterinaer instituttets rapportserie 2-2014.
- Nilsson and Holmen (1995). The Aquatic Adephaga (Coleoptera) of Fennoscandia and Denmark, Volume II. Dytiscidae. *Fauna entomologica Scandinavia* 32.
- McClay, W. (2000). Rotenone use in North America (1988-1997). *Fisheries* 20, 15-21. [https://doi.org/10.1577/1548-8446\(2000\)025%3C0015:RUINA%3E2.0.CO;2](https://doi.org/10.1577/1548-8446(2000)025%3C0015:RUINA%3E2.0.CO;2)
- McClay, W. (2005). Rotenone use in North America (1998-2002). *Fisheries* 30, 29-31. [https://doi.org/10.1577/1548-8446\(2005\)30\[29:RUINA\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2005)30[29:RUINA]2.0.CO;2)
- Pham, L., Jarvis, M. G., West, D & Closs, G. P. (2018). Rotenone treatment has a short-term effect on New Zealand stream macroinvertebrate communities. *New Zealand Journal of Marine and Freshwater Research*, 52 , 42-54. <https://doi.org/10.1080/00288330.2017.1330273>
- Rinne, A. & Wiberg-Larsen, P. (2017). Trichoptera Larvae of Finland: Identification Key to the Caddis Larvae of Finland and Nearby Countries. Trificon books.
- Sala O.E., Chapin III F.S., Armesto J.J. & Berlow, E. (2000). Biodiversity: global biodiversity scenarios for the year 2100. *Science* 287 , 1770-1774. <https://doi.org/10.1126/science.287.5459.1770>

Stensli, J.H. & Wist, A.N. (2014). Behandlingene i sone Indre Vefsnfjorden. In: Stensli, J.H. & Bardal, H. (eds.). Bekjempelse av *Gyrodactylus salaris* i Vefsnaregionen. (Eds: J. H. Stensli & H. Bardal). Veterinærinstituttets rapportserie 2-2014.

Vinson, M.R., Dinger, E.C. & Vinson, D.K. (2010). Piscicides and Invertebrates: After 70 Years, Does Anyone Really Know? *Fisheries* 35, 61-71. <https://doi.org/10.1577/1548-8446-35.2.61>

Weyl, P.S.R., de Moor, F. C., Hill, M.P. & Weyl, O.L.F. (2010). The effect of largemouth bass *Micropterus salmoides* on aquatic macroinvertebrate communities in the Wit River, Eastern Cape, South Africa. *African Journal of Aquatic Science* 2010, 35, 273–281. <https://doi.org/10.2989/16085914.2010.540776>

Tables

TABLE 1 Time of sampling (grey) and rotenone treatments (black) in River Fusta, River Drevjaelva, Lake Fustvatnet and Lake Drevvatnet

Hosted file

image1.emf available at <https://authorea.com/users/426555/articles/531027-long-term-responses-of-benthic-invertebrates-to-rotenone-treatment>

Figure captions

FIGURE 1 Overview of the two studied watercourses Fusta and Drevja. Triangles indicate the position of sampling stations. A double black bar indicates the upstream limit of the treatment of River Fusta in August 2011 and August 2012, and a single black bar indicates the upstream limit of the treatment of the watercourse in October 2012.

FIGURE 2 Mean changes in dissimilarity (% difference) for abundance data (a – c) and for presence-absence data (d – f) between sampling occasions (a and d) and its components; losses (b and e) and gains (c and f). Grey: River Fusta (rotenone treated), black: River Drevjaelva (untreated). Dashed vertical lines indicate the timing of the rotenone treatments. Error bars indicate maximum and minimum values of the temporal biodiversity indices at individual sampling stations. Significant differences in diversity indices between sampling periods ($p < 0.05$) in River Fusta are marked with grey asterisk (no significant differences detected in River Drevjaelva).

FIGURE 3 Average abundance (with standard deviation) of benthic invertebrates of River Fusta (rotenone treated) and River Drevjaelva (untreated) in the period 2011-2016. a) total abundance, b) Ephemeroptera, c) Plecoptera, d) Trichoptera, e) Chironomidae and f) Oligochaeta. Timing of rotenone treatments are marked with arrows.

FIGURE 4 Mean changes in dissimilarity (% difference) for abundance data (a – c) and for presence-absence data (d – f) between sampling occasions (a and d) and its components; losses (b and e) and gains (c and f). grey: Lake Fustvatnet (rotenone treated), black: Lake Drevvatnet (untreated). Dashed vertical lines indicate the timing of the rotenone treatment of the lake. Error bars indicate maximum and minimum values of the temporal biodiversity indices at individual sampling stations.

FIGURE 5 Average abundance (with standard deviation) of benthic invertebrates of Lake Fustvatnet (rotenone treated) and Lake Drevvatnet (untreated) in the period 2011-2016. A) Total abundance, b) Ephemeroptera, c) Plecoptera, d) Trichoptera, e) Chironomidae and f) Oligochaeta. Timing of the rotenone treatment is marked with an arrow.







