

Antiviral effects of propolis against SARS-COV 2. A mini review article

Rim Harfouch¹

¹Affiliation not available

January 4, 2022

Rim M. Harfouch,

Department of microbiology and biochemistry, Faculty of pharmacy, Al Sham private university, Latakia, Syria. Correspondence Autho's E-mail: r.h.foph.lat@aspu.edu.sy

Abstract:

Propolis, which is produced by honey bees from plant exudates, has long been used in traditional herbal medicine and is widely consumed as an immune system enhancer. The COVID-19 pandemic has renewed interest in propolis products worldwide; fortunately, various aspects of the SARS-CoV-2 infection mechanism are potential targets for propolis components. In pre-clinical studies, propolis promoted immune-regulation of pro-inflammatory cytokines, including reduction in IL-6, IL-1 beta and TNF- α . Propolis has also shown promise as an aid in the treatment of various of the comorbidities that are particularly dangerous in COVID-19 patients, including respiratory diseases, hypertension, diabetes, and cancer. In this mini review, we summarize the potential role of COVID-19 treatment and prevention.

Keywords:

COVID-19, propolis, antiviral activity.

Introduction:

Natural products are among the options being considered as an adjuvant treatment for SARS-CoV-2 infection, because they are generally inexpensive, widely available, and rarely have undesirable side effects. Some have proven antiviral activity. An important advantage of using natural remedies is that people who have other health problems or have mild flu-related symptoms could take simple and inexpensive measures to help reduce the impact of infection with SARS-CoV-2 [1]. Among these natural remedies is propolis. Propolis is a mixture of beeswax and resins collected by the honeybee from different plant buds, leaves and exudates. Bees use propolis not only as a building material but also as a means of maintaining low levels of bacterial and fungal concentration in the hive. More than 150 components such as polyphenols, phenolic aldehydes, quinones, coumarins, amino acids, steroids and inorganic components have been identified in propolis samples [2].

Infection by SARS-CoV-2, the virus that causes COVID-19, is characterized by binding between viral spike proteins and angiotensin-converting enzyme 2 (ACE2) [3]. SARS-CoV-2 infection is associated with increased levels of chemokines and activated pro-inflammatory cytokines that lead to the development of atypical pneumonia, with rapid respiratory impairment and pulmonary failure. Immunological/inflammatory phenomena have been shown to be important in the spectrum of SARS-CoV-2 infection. These mechanisms are associated with organ dysfunction more than the viral load [4].

Anti-COVID-19 Effects of Propolis:

In silico studies have investigated the use of flavonoids as effective therapeutic candidates against COVID-19 by targeting S protein cleavage by host-cell proteases, S protein binding to cell surface receptors such as ACE-II, inhibiting S protein, and S protein binding to the inflammatory B56 unit in PP2A as well as by interfering with NSPs of SARS-CoV-2 in order to hamper viral replication [5, 6]. A Brazilian Randomized control trial reported that treating hospitalized COVID-19 patients with a single oral daily dose of non-alcoholic preparation non-alcoholic preparation was associated with significant reductions in the length of hospital stay (LOS) and renal injury. Propolis treatment was not associated with a decrease in the need for oxygen therapy [7].

Identification of the most potent compounds is necessary for effective drug development; however, it is necessary to note that the pharmacological effects of supercritical extracts of propolis are superior to its single components (e.g., artemillin C and its precursor coumaric acid) indicating synergetic interaction of different compounds in propolis [8].

Conclusion:

The current emergency caused by the COVID-19 pandemic and limited therapeutic options, propolis is presented as a promising and relevant therapeutic option that is safe, easy to administrate orally and is readily available as a natural supplement and functional food.

References:

1. Berretta AA, Silveira MAD, C ndor Capcha JM, De Jong D. Propolis and its potential against SARS-CoV-2 infection mechanisms and COVID-19 disease: Running title: Propolis against SARS-CoV-2 infection and COVID-19. *Biomed Pharmacother.* 2020 Nov;131:110622. doi: 10.1016/j.biopha.2020.110622. Epub 2020 Aug 17. PMID: 32890967; PMCID: PMC7430291.
2. Harfouch R.M., Mohammad R., Suliman H. Antibacterial activity of Syrian propolis extract against several strains of bacteria in vitro. *World J. Pharm. Pharmaceuti. Sci.*, 6 (2016), pp. 42-46
3. Harfouch RM, Moualla YM, Shahoud SM, Belal LA, Mohammad BS, Mustafa NM, et al. Epidemiology of COVID-19 in the Most Pandemic Countries. *Ann Clin Cases.* 2020;1(4):1021.
4. Harfouch RM, Cytokine Storm Syndrome in COVID-19 Patients: Characteristics and Diagnosis. *Ann Clin Med Case Rep.* 2021; V7(18): 1-3.
5. da Silva, F.M.; da Silva, K.P.; de Oliveira, L.P.; Costa, E.V.; Koolen, H.H.; Pinheiro, M.L.; de Souza, A.Q.; de Souza, A.D. Flavonoid glycosides and their putative human metabolites as potential inhibitors of the SARS-CoV-2 main protease (Mpro) and RNA-dependent RNA polymerase (RdRp). *Mem. Inst. Oswaldo Cruz* 2020, 115, e200207.
6. Refaat, H.; Mady, F.M.; Sarhan, H.A.; Rateb, H.S.; Alaaeldin, E. Optimization and evaluation of propolis liposomes as a promising therapeutic approach for COVID-19. *Int. J. Pharm.* 2020, 592, 120028.
7. Duarte Silveira, M.A.; De Jong, D.; dos Santos Galv o, E.B.; Ribeiro, J.C.; Silva, T.C.; Berretta, A.A.; Amorim, T.C.; San Martin, R.L.A.; da Rebelo Concei o, L.F.M.; Dantas Gomes, M.M.; et al. Efficacy of propolis as an adjunct treatment for hospitalized COVID-19 patients: A randomized, controlled clinical trial. *medRxiv* 2021, 20248932.
8. Ali AM, Kunugi H. Propolis, Bee Honey, and Their Components Protect against Coronavirus Disease 2019 (COVID-19): A Review of In Silico, In Vitro, and Clinical Studies. *Molecules.* 2021; 26(5):1232. <https://doi.org/10.3390/molecules26051232>