

# Combining in vitro and in vivo screening to identify efficient *Pseudomonas* biocontrol strains against the phytopathogenic bacterium *Ralstonia solanacearum*

Sophie Clough<sup>1</sup>, Alexandre Jousset<sup>2</sup>, John Elphinstone<sup>3</sup>, and Ville-Petri Friman<sup>1</sup>

<sup>1</sup>University of York

<sup>2</sup>Utrecht University

<sup>3</sup>FERA

February 22, 2024

## Abstract

While plant pathogens are traditionally controlled using synthetic agrochemicals the availability of commercial bactericides is still limited. One potential control strategy could be the use of plant-growth-promoting bacteria (PGPBs) to suppress pathogens via resource competition or the production of antimicrobial compounds. This study aimed to conduct in vitro and in vivo screening of eight *Pseudomonas* strains against *Ralstonia solanacearum* (the causative agent of bacterial wilt) and to investigate underlying mechanisms of potential pathogen suppression. We found that inhibitory effects were *Pseudomonas* strain-specific, with strain CHA0 showing the highest pathogen suppression. Genomic screening identified 2, 4-diacetylphloroglucinol (DAPG), pyoluteorin, and orfamides A and B secondary metabolite clusters in the genomes of the most inhibitory strains, which were investigated further. While all these compounds suppressed *R. solanacearum* growth, only Orfamide A was produced in the growth media based on mass spectrometry. Moreover, orfamide variants extracted from *Pseudomonas* cultures showed high pathogen suppression. Using the Micro Tom tomato cultivar, it was found that CHA0 could reduce bacterial wilt disease incidence with one of the two tested pathogen strains. Together, these findings suggest that a better understanding of *Pseudomonas*-*Ralstonia* interactions in the rhizosphere is required to successfully translate in vitro findings into agricultural applications.

## Combining *in vitro* and *in vivo* screening to identify efficient *Pseudomonas* biocontrol strains against the phytopathogenic bacterium *Ralstonia solanacearum*

Clough, S.E.<sup>1,2</sup>, Jousset, A.<sup>3</sup>, Elphinstone, J.G.<sup>4</sup> and Friman, V-P.<sup>1</sup>

<sup>1</sup>University of York (Dept. Of Biology, Wentworth Way, University of York, York, YO10 5DD, U.K)

<sup>2</sup>Durham University (Dept. Of Chemistry, South Road, Durham University, Durham, DH1 3LE, U.K)

<sup>3</sup>Utrecht University (Biology Department, Institute of Environmental Biology, Ecology and Biodiversity Group, Padualaan 8, 3584CH Utrecht, Netherlands)

<sup>4</sup>Fera Science Ltd (National Agri-Food Innovation Campus, Sand Hutton, York, U.K)

### Abstract

While plant pathogens are traditionally controlled using synthetic agrochemicals the availability of commercial bactericides is still limited. One potential control strategy could be the use of plant-growth-promoting bacteria (PGPBs) to suppress pathogens via resource competition or the production of antimicrobial compounds. This study aimed to conduct *in vitro* and *in vivo* screening of eight *Pseudomonas* strains against *Ralstonia solanacearum* (the causative agent of bacterial wilt) and to investigate underlying mechanisms of potential pathogen suppression. We found that inhibitory effects were *Pseudomonas* strain-specific, with strain CHA0 showing the highest pathogen suppression. Genomic screening identified 2, 4-diacetylphloroglucinol (DAPG), pyoluteorin, and orfamides A and B secondary metabolite clusters in the genomes of the most inhibitory strains, which were investigated further. While all these compounds suppressed *R. solanacearum* growth, only Orfamide A was produced in the growth media based on mass spectrometry. Moreover, orfamide variants extracted from *Pseudomonas* cultures showed high pathogen suppression. Using the 'Micro Tom' tomato cultivar, it was found that CHA0 could reduce bacterial wilt disease incidence with one of the two tested pathogen strains. Together, these findings suggest that a better understanding of *Pseudomonas-Ralstonia* interactions in the rhizosphere is required to successfully translate *in vitro* findings into agricultural applications.

## Introduction

The use of chemical pesticides has declined in recent years due to elevated costs, environmental toxicity, and stricter legislation (Chen *et al.*, 2016). As a result, new methods and approaches are required to control plant pathogens and to ensure future food security in the face of expanding human population (Hayward, 1991; Kaczmarek *et al.*, 2014; Wang *et al.*, 2019). One alternative to traditional agrochemicals is biocontrol, which depends on using natural microbial competitors against a variety of pathogens to restrict pathogens' growth and survival. Several biocontrol agents have been identified and found to show a broad range of activity against bacterial, fungal, viral, and nematode pathogens (Sofrata *et al.*, 2011; Hansen and Keinath, 2013; Wang *et al.*, 2019; Nagachandrabose, 2020). Plant growth-promoting bacteria (PGPB) are one example of potentially highly effective biocontrol agents, with a broad-range biocontrol activity (Pierson *et al.*, 1998). PGPB biocontrol outcomes are however often variable in field conditions, despite clear inhibition observed under laboratory conditions (Compant *et al.*, 2005). One reason for this is that lab conditions fail to replicate actual field conditions in terms of nutrient availability, competition with rhizospheric bacteria, degradation, and adsorption of secreted antimicrobials, and the likelihood of successful colonization in natural environments, these can all affect the efficacy of disease suppression (Weller, 2007). Additionally, even though biocontrol agents can show a broad range of effects against different types of pathogens, they might not be able to inhibit different genotypes of one given pathogen species (Xue *et al.*, 2013). Considering pathogen-biocontrol agent interactions at the genotype level, and the mechanisms by which inhibition is mediated in more realistic environmental conditions, are therefore required to develop functionally robust biocontrol methods.

Plant pathogenic bacteria belonging to the *R. solanacearum* species complex (RSSC) cause bacterial wilt disease in many wild and cultivated plants. It is ranked the second most important bacterial plant pathogen globally and possesses numerous virulence mechanisms that allow it to infect more than 250 plant species from 54 plant families (Hayward, 1991; Genin and Denny, 2012; Mansfield *et al.*, 2012; Nion and Toyota, 2015). RSSC has spread globally across the world and has a quarantine status in many countries meaning it is monitored regularly to prevent further spread (EPPO, 2021). In addition to the trade of infected plant materials, RSSC global distribution is likely explained by its high genetic variability and ability to rapidly adapt to surrounding environmental conditions (Genin, 2010; Genin and Denny, 2012). Between-strain differences include variation in pathogenicity-related genes, such as the presence of type III secretion effectors and genes involved in chemotaxis, adherence, secondary metabolic pathways, biosynthesis of phytohormones, and the detoxification of various antimicrobial compounds which are important in determining *R. solanacearum* host range (Genin, 2010; Genin and Denny, 2012). This variation can be generated via accumulation of mutations (Guidot *et al.*, 2014; Gopalan-Nair *et al.*, 2021) or horizontal gene

transfer between different *R. solanacearum* strains, which can result in an exchange of virulence traits between different genotypes (Guidot *et al.*, 2009). While *R. solanacearum* strains in Europe are considered to be clonal, belonging to the phylotype IIB group (Castillo and Greenberg., 2007; Clarke *et al.*, 2015), some strain variation exists at the local scale within countries (Parkinson *et al.*, 2013; Caruso *et al.*, 2017). To what extent this variation affects the pathogen response to biocontrol PGPBs is however currently unknown.

Traditionally, the development of a biocontrol method begins *in vitro* in the lab by identifying bacterial strains with potential biocontrol activity (Fravel, 2005). Biocontrol efficiency within *Pseudomonas* plant growth-promoting species is often mediated by a variety of secreted secondary metabolites, including several antimicrobials and iron-scavenging siderophores (Becker *et al.*, 2012). Some biocontrol characteristics are more common amongst these strains, including the ability to produce 2, 4-diacetylphloroglucinol (DAPG) and hydrogen cyanide (HCN) - which are effective antimicrobial against bacterial, fungal, and nematode plant pathogens (Cronin *et al.*, 1997; Compant *et al.*, 2005; Haas and Keel, 2003; Haas and Défago, 2005; Humair *et al.*, 2009, Almario *et al.*, 2017). In contrast, other biocontrol properties are strain-specific, such as the production of cyclic orfamide lipopeptides by certain subgroups of *Pseudomonas fluorescens* such as *P. protegens* strains: CHA0 and Pf-5 (Ma *et al.*, 2016). Methods for screening bacterial biocontrol agents against *R. solanacearum* include exposing pathogenic bacterial strains to biocontrol bacteria directly in co-cultures and measuring pathogen growth or survival as a function of time (Fravel, 2005). Supernatant assays are also often used to test if secretions in the media suppress the growth of the pathogens (Kaur *et al.*, 2019; Yang *et al.*, 2019). However, the successful identification of biocontrol strains *in vitro* does not always translate to successful biocontrol outcomes *in vivo* and *Pseudomonas* biocontrol outcomes are often variable in field conditions despite clear inhibition observed in laboratory conditions. For example, *Pseudomonas* strains have been shown to unsuccessfully replicate *in vitro* protection against *R. solanacearum* during *in vivo* experiments with *Eucalyptus* trees, and it was suspected this may have been due to low-level expression of genes responsible for biocontrol activity (Ran *et al.*, 2005). The success of biocontrol is further influenced by competition with native bacteria in the rhizosphere, and it has also been shown that the effectiveness of their secondary metabolites, such as DAPG, can be influenced by the age and species of the host plant is has been selected to protect (Notz *et al.*, 2001; Siddiqui and Shaikat, 2003). Soil properties such as moisture, temperature, clay content, and pH can also influence the success of biocontrol (Van der Putten *et al.*, 2006), while common bacterial feeders, such as protozoa and nematodes, could also predate on the biocontrol bacteria (Pedersen *et al.*, 2009). Therefore, it is vital that *in vitro* screening is coupled with *in vivo* validation of biocontrol effectiveness with plants to develop successful biocontrol applications with translational potential.

Here we used such an approach to screen and identify efficient *Pseudomonas* biocontrol bacterial strains against six UK and one Polish *R. solanacearum* isolates that belong to an economically important potato-specific RSSC lineage (Phylotype IIB sequevar 1; formerly known as race3 biovar2 strains). We first tested direct and indirect interactions between *R. solanacearum* strains and eight plant growth-promoting *Pseudomonas* strains *in vitro*, which were chosen for their previously characterized antimicrobial activity against a Chinese *R. solanacearum* isolate belonging to Phylotype I (Hu *et al.*, 2016). We then conducted genomic screening of *Pseudomonas* strains to identify potential antimicrobial secondary metabolite clusters and directly tested whether these compounds were produced and if they inhibited *R. solanacearum* under laboratory conditions. Finally, the biocontrol potential of the *P. protegens* CHA0 strain was tested against two *R. solanacearum* strains in the tomato rhizosphere. Results revealed that antimicrobial effects of *Pseudomonas* were strain-specific, and the strongest inhibitory effects were mediated by *P. protegens* strain CHA0. Of the tested compounds, the Pyoluteorin antibiotic had the strongest inhibitory effect followed by DAPG and Orfamides A and B. While CHA0 also showed biocontrol efficiency with tomato, this was clear only with one of the tested UK *R. solanacearum* isolates. Together, these results show that while *in vitro* screening can be used to identify potential *Pseudomonas* biocontrol strains against *R. solanacearum*, more work is needed to understand their biocontrol activity *in vivo* in the plant rhizosphere.

## Methods

### **(a) Culturing and maintenance of bacterial strains**

Eight fluorescent *Pseudomonas* strains (CHA0, Pf-5, Q2-87, Q8R1-96, 1M1-96, MVP1-4, F113, and Ph11C2) which have been studied extensively previously in biocontrol (Hu *et al.*, 2016), and seven *Ralstonia solanacearum* strains (#1-7), which were isolated as a part of annual river sampling survey in England and Wales by Fera Science Ltd, were used in the experiments (listed in Tables A1-A2). All bacteria were stored in 20% glycerol stocks at -80 °C. Prior to experiments, bacterial starting cultures were prepared as follows: frozen samples were inoculated in 5 mL of LB, NB, or CPG broth (media recipes described in Table A3) and incubated with shaking at 200 rpm at 28°C for 24 hours. Rich media preparations were used throughout all experiments to ensure the efficient growth of both bacterial genera. Bacterial cultures were prepared similarly throughout all experiments unless stated otherwise.

### **(b) Measuring *R. solanacearum* inhibition by *Pseudomonas* strains in direct contact**

Soft agar overlay assays were used to test direct inhibition of *R. solanacearum* by *Pseudomonas* strains. *Pseudomonas* strains were grown for 24 h in 5 mL of 100% LB broth at 28 °C with shaking

at 200 rpm and *R. solanacearum* strains were grown for 30 h under the same conditions. Soft agar overlay plates were prepared by first filling sterile 90 mm Petri dishes with a layer of hard LB which was left to solidify. 200  $\mu$ L of each *R. solanacearum* culture was then mixed with 20 mL of cooled (below 55°C) liquid soft agar and poured on top of the hard agar layer. As *R. solanacearum* grows more slowly than *Pseudomonas*, plates were left to incubate for ~ 8 hours prior to spotting the *Pseudomonas* cultures on top of the soft agar overlay as follows. Each plate was divided evenly into quarters and 2  $\mu$ L of a *Pseudomonas* strain (approximately  $1.0 \times 10^6$  CFU/mL) was spotted on the center of each quarter of the plate (example shown in Figure A1). Plates were then incubated upside down at 28 °C and zones of inhibition were recorded after 96 hours. The distance of the inhibition zones was measured from the outer edge of the *Pseudomonas* spot to the *R. solanacearum* lawn (in millimeters with a ruler). Each strain combination treatment was carried out in triplicate.

#### **(c) Measuring *R. solanacearum* inhibition by *Pseudomonas* strains indirectly using supernatant assays**

To investigate *R. solanacearum* inhibition by *Pseudomonas* in the absence of direct contact, we exposed *R. solanacearum* strains to supernatants of each *Pseudomonas* species in pairwise supernatant cultures. The *Pseudomonas* supernatants included all secondary metabolites excreted when *Pseudomonas* strains were grown alone in LB media and were prepared as follows. All *Pseudomonas* strains were first grown individually in 20 mL of LB broth (Table A3) for 24 h with shaking at 200 rpm. The supernatant was then prepared by centrifuging cultures for 10 minutes at 4000 g before separating bacterial cells and fragments from soluble material including secondary metabolites using 0.2  $\mu$ m filters. The inhibition was measured using flat bottomed 96-well plates in 50:50 *Pseudomonas* supernatant to LB mixtures. As a negative control *R. solanacearum* was grown in 50:50 LB in the sterile water mixture. At the start of the experiment, every supernatant mixture was inoculated with 2  $\mu$ L of each *R. solanacearum* strain (approximately  $1.0 \times 10^6$  CFU/mL), and microplates were then incubated at 28 °C for three days and their bacterial densities were recorded as optical density at 24 and 72 h (OD 600 nm; Tecan Infinite spectrophotometer). All pairwise combinations were replicated four times and control treatments three times.

#### **(d) Genome sequencing of *Pseudomonas* strains**

Single colonies of each *Pseudomonas* strain were inoculated in NB broth and grown for 12 h at 30 °C with shaking at 170 rpm. Genomic DNA was extracted using the QIAGEN blood and cell culture DNA kit (catalog No. QIAGEN Genomic-tip 100/G, Midi 13343) following the manufacturer's protocol, and DNA quality was tested using nanodrop. All strains were sequenced

using the Miseq platform (2\*300 bp) at the Utrecht Sequencing Facilities. To obtain more accurate genome assemblies, reads were pre-processed as follows. We first removed adapter sequences, read shorter than 50 bp, and low-quality nucleotides using a Phred quality score threshold of < 20. Genome assemblies were carried out in two steps. First, we used SOAPdenovo v2.04 to assemble reads into contigs and scaffolds based on K-mer size (available at <http://soap.genomics.org.cn/>). Second, GapCloser v1.12 was used to close gaps emerging during the scaffolding process by SOAPdenovo (Li *et al.*, 2010).

#### ***(e) Investigating Pseudomonas secondary metabolite biosynthesis gene clusters using AntiSmash5.0***

The *de novo* assembled draft genomes of the eight *Pseudomonas* strains were analyzed to identify potential secondary metabolic clusters linked with antibiosis using the antiSMASH 5.0 pipeline (<https://antismash.secondarymetabolites.org>, Blin, *et al.*, 2017, 2019), which allows rapid genome-wide identification and in-depth analysis of secondary metabolite biosynthesis gene clusters based on several open-source databases. Fasta files were uploaded to AntiSMASH5.0 bacterial version with detection strictness set to “relaxed” and all search features included to maximize potential metabolic clusters identified. Based on this analysis, we identified four antimicrobial compounds which could potentially be produced by our *Pseudomonas* strains. One of these compounds, 2, 4-diacetylphloroglucinol (DAPG), is a common antimicrobial produced by all eight fluorescent *Pseudomonas* strains, while three specific gene clusters for Pyoluteorin (an antimicrobial with unknown mechanisms of action; Kidarsa *et al.*, 2011) and Orfamides A and B (cyclic lipopeptides that can cause membrane pore formation; Ma *et al.*, 2016) were found only in CHA0 and Pf-5 *Pseudomonas* genomes.

#### ***(f) Determining the effects of identified Pseudomonas metabolites on R. solanacearum growth in single-compound and multi-compound mixtures***

Pure compounds of Orfamide A (Santa Cruz Biotechnology), Orfamide B (Santa Cruz Biotechnology), Pyoluteorin (Santa Cruz Biotechnology), and 2,4-diacetylphloroglucinol (DAPG) (Santa Cruz Biotechnology) were purchased to test their efficacy against *R. solanacearum* strains. 10 mM stocks were made in 100% DMSO and stored at -20 °C, except for DAPG, for which a 100 mM stock was prepared in 100% methanol and stored at -20 °C. The effect of DAPG was measured by inoculating 2 µL of each *R. solanacearum* strain (approximately 1.0x10<sup>6</sup> CFU/mL) to 100% LB broth with DAPG at the following concentrations: 1000, 500, 100, 50, and 0 µM (negative control). Due to the relatively high costs of chemicals, only two *R. solanacearum* strains (#1 and #7) were tested for their susceptibility to Pyoluteorin (100 µM concentration only). Similarly, only strain #1

susceptibility was tested to Orfamides A and B (100  $\mu$ M concentration only) following the same methods as with DAPG. Bacterial densities were recorded as optical density at 0 h and 24, 48, and 72 h after inoculation with a spectrophotometer (OD 600 nm; Tecan Sunrise spectrophotometer).

***(g) Confirmation of the production of secondary metabolites in the *Pseudomonas* supernatant using mass spectrometry***

Mass spectrometry was used to identify and quantify any secondary metabolites produced by CHA0 and Pf-5 in the growth conditions they were exposed to *in vitro* (compounds were identified against chemical standards). We also used untargeted analysis to identify any novel antimicrobials produced by comparing peaks against existing databases.

The strains were grown in triplicate in LB broth with 200 rpm shaking at 28 °C for 24 h. Following incubation, bacterial densities were normalized to an optical density of 0.1 (OD 600nm), and the cultures were centrifuged for 10 minutes at 4000 g. The supernatant was then filtered using 0.2  $\mu$ m filters to separate bacterial cells and fragments from secondary metabolites present in the supernatants. A total of 6 samples were produced (two *Pseudomonas* strains in one growth condition in triplicate). 200  $\mu$ L of each sample was provided to the Centre of Excellence in Mass Spectrometry (CoEMS) at the University of York for mass spectrometry analysis. The LC separation was performed by an Acquity UPLC I class system (Waters, Elstree, UK), on a BEH C18 100x2.1, 1.7 $\mu$  column (Waters). The sample injection volume was 7.5  $\mu$ L. The MS end was a Synapt G2S-Si QTOF (Waters) mass spectrometer which was operated in positive ESI, resolution mode, using the HDMS<sup>E</sup> acquisition technique (alternating scans of MS and MS<sup>2</sup> acquisitions (fragmentation by CID) alongside traveling wave ion mobility separation). Voltages in the low energy function (MS) were 4 V both in the trap and transfer cell; and in the high energy function (MS<sup>2</sup>), the voltage was 4 V in the trap and a 20 -100 V ramp in the transfer cell. The source was operated with the capillary set to 3 kV, source temperature 150 °C, desolvation temperature 450 °C, cone gas flow 20 L/h, desolvation gas flow 450 L/h, and nebulizer 6.5 bar. Trap gas flow was 2 mL/min, helium cell flow 180 ml/min, IMS gas flow 80 mL/min. Mass range was 100-1000m/z, scan time 0.2 sec, lock mass was leucine-enkephalin at m/z 556.2771.

Data was collected using MassLynx Software Version V4.2 SNC983 (Waters) and analyzed using UNIFI 1.9 (Waters) for semi-quantitative analysis of target analytes (Orfamide A, Orfamide B, Pyoluteorin, and DAPG). Calibration curves were constructed for target analytes at 10, 50, 100, 300, and 500 nM in LB media; the MS response for all analytes was linear in this range. Analytes were quantified using the calibration curve in the appropriate medium. Blanks, standards, and samples were injected in technical triplicates from the same vial. Progenesis QI v.2.0 (Waters) was used for the untargeted analysis of metabolomics data: putative compound identification was based

on a score comprising MS match of exact mass of the compound precursor (ChemSpider database search with 10 ppm mass tolerance) and a match to *in silico* MS<sup>2</sup> fragmentation pattern. Further data processing and downstream analysis were performed using Bioconductor package XCMS in R, combined with a database (LipidMaps) search.

Untargeted analyses were also conducted by comparing the collected dataset with the Knapsack database (<http://www.knapsackfamily.com/KNAPSAcK/>) Briefly, Waters.raw files were converted to zlib compressed 32-bit precision .mzML files using ProteoWizard MSConvert [1] version 3.0.19172, using the combineIonMobilitySpectra filter, and spectra centroided using the qtofpeakpicker with resolution set to 20000 and threshold 1. Using custom scripts in R 3.0.0 operating in a Linux 64 bit environment, .mzML files were further processed using the mzR package [1] to lockmass correct spectra against leucine enkephalin ( $[M+H]^+ = 556.27568$ ,  $[M-H]^- = 554.26202$ ). Lockmasses were identified within a  $\pm 0.1$  Da  $m/z$  window, and the running mean of lockmasses in a 60 s moving window was used to adjust the  $m/z$  values for every scan. Across files, feature detection was achieved using the `xcmsSet()` function from the `xcms` package [2], using the following parameters: `method = 'centWaveWithPredictedIsotopeROIs'`, `ppm = 10`, `snthresh = 10`, `peakwidth = c(3, 20)`, `prefilter=c(3, 1000)`, `integrate = 2`, `mzdiff = -0.1`, `firstBaselineCheck = FALSE`. Custom scripts were used to retain gaussian features, which were then aligned and missing values recalculated across samples using the `xcms group()` and `fillPeaks()` functions, respectively. Feature areas were adjusted by subtracting the mean + 3 x the standard deviation of blank samples, with adjusted values < 0 being set to 0. The feature list was then filtered to retain only the most intense monoisotope belonging to a single compound as identified by CAMERA [3], with areas above 0 in at least 90% of blank-subtracted samples. These filtered features were reported as `masstags` (unique  $m/z$  and retention time pairs) and annotated where possible against authentic standards or putative compounds from literature or databases based on exact mass. The Progenesis QI v.2.0 (Waters) was also used to search for compounds by their monoisotopic mass which have been reported to be present in *Pseudomonas* genomes from the literature. The potential metabolites were identified if their observed monoisotopic mass matched with the accurate known ppm within a range of  $\pm 10$  ppm. However, without a chemically verified standard for definitive identity matches, this can only be considered speculative.

#### ***(h) Extraction and isolation of Pseudomonas orfamide antimicrobials and testing their effects on R. solanacearum growth***

A protocol was adapted from Ma *et al.* (2016) to extract cyclic lipopeptide orfamides from the CHA0 *Pseudomonas* strain. Briefly, a starting culture of CHA0 was grown in a 250 mL flask containing 50 mL of King's Broth growth medium at 28 °C with shaking at 200 rpm. This was then

inoculated into a 2 L flask containing 500 mL of liquid media and shaken at 150 rpm for 48 h. *Pseudomonas* supernatant was collected via centrifugation at 10,000 *g* for 10 minutes (J25 XP series centrifuge). The supernatant was then acidified to a pH of 2.0 using 8 M HCl and stored overnight at 4 °C. The precipitate was collected after centrifugation at 10,000 *g* for 20 minutes and extracted with 1 mL of 100% methanol. At this stage in the centrifuge tube, there were two distinguishable precipitates, one at the bottom of the tube and one close to the top. It was decided to collect these separately and call them 'A' and 'B' as it was unknown if they would have different or similar properties. The organic phase was collected by spinning samples until dry in a vacuum concentrator (Savant svc SpeedVac 100h concentrator) and samples were then dissolved in 300 µL of 100% DMSO to make orfamide variant stocks. Due to a limited amount of these orfamide variant stocks, only *R. solanacearum* strains #1 and #7 were used for inhibition testing. Inoculant cultures were grown in 5 mL of LB broth overnight and 2 µL of each *R. solanacearum* strain was inoculated with 1% orfamide variants in LB broth (200 µL final volume). Bacterial densities were recorded at 24, 48, and 72 h (OD 600 nm; Tecan Infinite spectrophotometer), and each treatment was carried out in triplicate.

**(i) Testing *Pseudomonas* biocontrol efficiency in vivo using a tomato plant model**

To explore the translational potential of *Pseudomonas* CHA0 strain *in vivo*, we tested if any of the inhibitory effects observed *in vitro* could protect plants from *R. solanacearum* infections using the Micro Tom tomato cultivar. This cultivar of tomato was selected as it is well established and often used when studying *R. solanacearum* infections (Gu *et al.*, 2016). The infectivity of two *R. solanacearum* UK strains (#1 and #7) was tested in the absence and presence of CHA0 (a negative water control treatment was included). Tomato (*Solanum lycopersicum* 'Micro Tom') plants were grown in an incubated light chamber at 28 °C with 16:8 h light:dark conditions with regular watering. Seeds were sown in 7 cm seedling trays in 35 g of autoclaved compost (John Innes #2) where they remained for the entire experiment. Experiments were conducted between July and September 2019 in temperature-controlled plant growth chambers (Sanyo MLR-352) at the University of York. To test the biocontrol efficacy of CHA0, the *Pseudomonas* culture was prepared by inoculating 100 mL of CPG broth with 100 µL of frozen stock bacterial culture in 300 mL glass culture flasks and the strain was grown for 24 h at 28 °C with shaking at 200 rpm. Bacterial culture was then washed of nutrient media by centrifuging at 4000 *g* for 10 minutes and resuspending in sterile dH<sub>2</sub>O. Seeds were germinated and sown in 7 cm seedling trays with 35 g of autoclaved compost and regularly watered using sterile water (for growth conditions see earlier section). *Pseudomonas* strain was inoculated onto the soil of 4-week old seedlings one week prior to *R. solanacearum* strains to allow time for the biocontrol bacteria to effectively colonize the soil and

roots. Before inoculation, *Pseudomonas* cell densities were adjusted to an OD 600 nm of 0.25 (approximately  $1.0 \times 10^9$  CFU/mL) and 6 mL was poured into each pot, soaking through the soil and roots, and any excess culture remained in the saucer below. Roots of tomato seedlings were cut prior to *R. solanacearum* inoculation using a sterile scalpel to mimic natural pathogen entry points in the field. *R. solanacearum* strains #1 and #7 were cultured in 30 mL of CPG broth (Table A3) for 48 h with shaking at 200 rpm at 28 °C and adjusted to an OD600 nm of 0.25. A total of 1 mL of each *R. solanacearum* strain was inoculated at the base of the visible stems one week after *Pseudomonas* inoculation. Three replicates were used for each treatment and each replicate consisted of nine individual plants. All plants were watered one hour prior to monitoring infection to be certain that wilting was not due to dehydration. Bacterial wilting symptoms were recorded daily by scoring the plants as “uninfected” or “infected” when plants showed clear wilting symptoms. The experiment was terminated three weeks after pathogen inoculation and above-ground dry weight was recorded for each replicate plant.

### **(j) Statistical analyses**

We used repeated measures ANOVA to analyze mean differences between pathogen densities for all experiments including temporal data such as *R. solanacearum* exposure to different secondary metabolites. Two-way ANOVA was used to analyze the mean differences between treatments when only a one-time point was used for analysis, followed by pairwise post hoc Tukey tests with 95% confidence levels and Bonferroni-corrected p-values. These analyses were used when interpreting the direct and indirect soft agar and bacterial supernatant interactions between *R. solanacearum* and *Pseudomonas*. Poisson glm models with chi-squared tests were used to analyze the binomial *in vivo* tomato infection data. All statistical analyses and graphs were produced using R (R Foundation for Statistical Computing, R Studio Version (3. 4. 4), Packages: ggplot, tidyverse, nlme, rcompanion).

## **Results**

### **(a) Measuring the direct inhibition of *R. solanacearum* by *Pseudomonas* strains using soft agar assays**

Clear inhibition zones were detected in all pathogen strain treatments indicating that each *Pseudomonas* strain could inhibit the growth of *R. solanacearum* when in direct contact. However, inhibition zone sizes differed depending on the identities of interacting *Pseudomonas* and *R. solanacearum* strains (*Ralstonia* × *Pseudomonas*:  $F_{42, 112} = 3.877$ ,  $P < 0.0001$ , Figure 1A). This variation appeared to be mainly driven by *Pseudomonas* strains, as post hoc analyses revealed only a small and non-significant variation between different *R. solanacearum* strains (*Ralstonia*:  $F_{6, 161} = 0.6364$ ,  $P = 0.7009$ , Figure 1A). In contrast, much higher variation was observed between the

inhibitory activity of *Pseudomonas* strains (*Pseudomonas*:  $F_{7, 160} = 60.81$ ,  $P = 0.0001$ , Figure 1A), and post hoc analyses revealed that CHA0 was the most inhibitory strain, followed by Q8R1-96 and MVP1-4, which showed modest levels of inhibition (Tukey:  $P < 0.05$ ). The strains 1M1-96 and Ph11C2 caused the smallest zones of inhibition on *R. solanacearum* lawns overall (Tukey:  $P < 0.05$ ). These results suggest that all *Pseudomonas* strains inhibited the growth of *R. solanacearum* in direct contact, which varied between different *Pseudomonas* strains.

### **(b) Measuring the indirect inhibition of *R. solanacearum* by *Pseudomonas* strains using supernatant assays**

Almost all *Pseudomonas* supernatants suppressed the growth of *R. solanacearum* compared to the no-supernatant LB control treatments (*Supernatant treatment*:  $F_{8, 236} = 3.697$ ,  $P = 0.01567$ , Figure 1B). In contrast to agar plate assays, *R. solanacearum* strain variation was more evident and the growth reduction of strains #2 and #3 was significantly lower compared to other strains (*Ralstonia*:  $F_{6, 217} = 19.1102$ ,  $P < 0.001$ , Figure 1B). The growth of strains #1, #6, and #7 was reduced most by the *Pseudomonas* supernatants overall. In line with direct inhibition assays, the *Pseudomonas* CHA0 strain caused the greatest reduction in *R. solanacearum* growth, followed by strains Ph11C2 and MVP1-4, which showed intermediate growth reduction (*Pseudomonas strain*:  $F_{7, 216} = 11.3595$ ,  $P < 0.001$ , Figure 1B). *Pseudomonas* strains 1M1-96 and Q8R1-96 showed relatively small growth reduction. Together, these findings are qualitatively similar to the direct inhibition assay (Figure 1A), highlighting the high efficacy of the *Pseudomonas* CHA0 strain in suppressing the growth of *R. solanacearum*.

### **(c) Comparative genomic analysis reveals variation in the presence of secondary metabolic clusters between different *Pseudomonas* strains**

Comparative genomic analysis of secondary metabolite clusters based on the AntiSMASH output identified between 11-17 metabolic clusters in each of the eight *Pseudomonas* genomes (Tables A4-A11). The type and diversity of recognized clusters are shown in Table 1. Non-ribosomal peptide synthetases (NRPS) were the most abundant secondary metabolite clusters, which was expected as these account for several common antimicrobials produced by fluorescent *Pseudomonas* strains (Hesse *et al.*, 2018; Girard, L., Höfte, M. and Mot., 2020). Similarly, DAPG metabolite (belonging to the T3PKS cluster), as well as the pyoverdine siderophore (NRPS cluster) metabolite clusters, were found in all strains. Overall, the highest number of clusters were detected in CHA0 and Pf-5 strains (17). These strains also harbored some unique metabolite clusters such as the T1PKS metabolic cluster, which encodes Pyoluteorin antimicrobial, and the CDPS cluster, which encodes unknown metabolites. CHA0 and Pf-5 also had the greatest number of NRPS clusters and were the

only two strains capable of producing the cyclic lipopeptides known as orfamides. Other *Pseudomonas* strains also possessed some unique clusters such as the ectoine metabolic cluster found in the Q2-87 strain. A more in-depth insight into the clusters and the percentage similarity with various *Pseudomonas* and other bacterial genomes based on the AntiSMASH database is shown in Tables A4-A11. The all clusters which were investigated further had 95-100% similarity with already characterized secondary metabolite clusters, increasing the certainty of their predicted identity and functioning.

#### **(d) Confirming the production of *Pseudomonas* secondary metabolites using mass spectrometry**

Our genome screening results suggest that *Pseudomonas* inhibitory activity could have been due to the presence of DAPG or certain, less common, secondary metabolism clusters encoding Pyoluteorin and Orfamides that were identified only in CHA0 and Pf-5 strains. Due to the presence of these two unique clusters, CHA0 and Pf-5 strains were chosen for a more detailed study. To verify the production of these compounds in LB media, the supernatants of CHA0 and Pf-5 were analyzed using mass spectrometry against chemical standards for DAPG, pyoluteorin, orfamide A, and orfamide B. Based on matching monoisotopic masses to a database of *Pseudomonas* metabolites, a list of putative metabolites present in the *Pseudomonas* supernatants of CHA0 and Pf-5 was determined using Progenesis QI v.2.0 (Waters) analysis (Table A12). These included compounds such as enantiopyochelin, rhizoxin, and indole-3-acetic acid which have different metabolic functions that can enhance bacterial survival (Table A12). For example, pyochelin is a siderophore that can aid iron acquisition when *Pseudomonas* may be in an iron-limited environment to improve its survival chances and outcompete other bacterial competitors (Duffy and Defago, 1999). Based on a comparison with four available standards, only Orfamide A production by CHA0 strain was detected in LB supernatant with an average concentration of 7.5 mg/mL (Figure A2). In contrast, none of the other four candidate compounds were detected in Pf-5 supernatant in LB media. These results suggest that only Orfamide A was produced by CHA0 in the conditions used in the supernatant inhibition assays. However, approximately 70 unidentified metabolites were detected in the samples, which could also have contributed to *R. solanacearum* inhibition.

#### **(e) Testing the inhibitory effects of identified *Pseudomonas* secondary metabolites on *R. solanacearum* growth**

The inhibitory effects of DAPG, pyoluteorin, and orfamides A and B were tested individually against *R. solanacearum* strains using commercially available chemical standards. The effect of DAPG was tested against all *R. solanacearum* strains, while the effect of Orfamides and

Pyoluteorin were tested against a subset of strains due to the high costs of chemical standards (*Ralstonia* strain #1 for Orfamides, and *R. solanacearum* strains #1 and #7 for Pyoluteorin). It was found that DAPG suppressed every *R. solanacearum* strain in a concentration-dependent manner (Concentration:  $F_{4, 200} = 238.2599$ ,  $P < 0.001$ , Figure 2A), and all tested strains were unable to grow at the two highest concentrations (500  $\mu\text{M}$  and 1000  $\mu\text{M}$ ). When all concentrations were included in the analysis, the effect of DAPG was independent of the *R. solanacearum* strain identity (*Ralstonia*:  $F_{6, 98} = 0.724$ ,  $P = 0.6305$ , Figure 2A). However, excluding the two highest concentrations, where all growth was inhibited, some *R. solanacearum* strain differences were revealed (*Ralstonia*:  $F_{6, 56} = 3.355$ ,  $P = 0.0068$ , Figure 2A): Strain #4 was the least susceptible to DAPG, while strains #3, #5 and #7 were relatively more susceptible. We also considered growth reduction by DAPG with growth in the absence of DAPG at the final time point for all *R. solanacearum* strains. Growth reduction was greater in 100  $\mu\text{M}$  DAPG than 50  $\mu\text{M}$  (Concentration:  $F_{1, 124} = 46.8$ ,  $P < 0.001$ , Figure 2A) and between strain variation was evident at both concentrations (*Ralstonia* at 100  $\mu\text{M}$ :  $F_{6, 56} = 5.671$ ,  $P < 0.001$ , Figure 2A; *Ralstonia* at 50  $\mu\text{M}$ :  $F_{6, 56} = 3.856$ ,  $P = 0.00273$ , Figure 2A). Post hoc analyses revealed that strains #5 and #6 were the least susceptible to DAPG, while strains #2 and #3 were relatively more susceptible. Both orfamides A and B reduced the growth of *R. solanacearum* strain #1 and this effect became clearer over time (Metabolite:  $F_{2, 6} = 460.7$ ,  $P < 0.001$ ; Metabolite  $\times$  Time:  $F_{4, 12} = 45.806$ ,  $P = 0.0359$ , Figure 2B). However, no difference was found between orfamides A and B ( $P = 0.11$ ). No visible growth was observed when either *R. solanacearum* strains #1 or #7 were exposed to pyoluteorin (*Ralstonia*:  $F_{2, 16} = 48.596$ ,  $P < 0.001$ , Figure 2C), suggesting that both strains were highly susceptible to this compound.

#### **(f) Testing the inhibitory effects of orfamide variants isolated from *Pseudomonas* CHA0 on *R. solanacearum* growth**

Orfamide variants 'A' and 'B' were isolated from the *Pseudomonas* CHA0 strain and their effects tested only against *R. solanacearum* strains #1 and #7 due to limited quantities of extracted compounds (Figure 3). The growth of both *Ralstonia* strains was constrained by both orfamide variants (*Ralstonia*  $\times$  Treatment:  $F_{2, 12} = 0.0476$ ,  $P = 0.9537$ , Figure 3) and the growth suppression by the orfamide variants became visible only after 48 h during the assays (Treatment  $\times$  Time:  $F_{4, 30} = 75.652$ ,  $P < 0.0001$ , Figure 3). At the final time point, there were clear differences between orfamide variants on both strain #1 (Treatment:  $F_{2, 10} = 681.3$ ,  $P < 0.001$ , Figure 3) and strain #7 (Treatment:  $F_{4, 10} = 65.77$ ,  $P < 0.001$ , Figure 3), with orfamide variant 'B' showing a slightly higher pathogen growth suppression. Together these results suggest that the *Pseudomonas* CHA0 strain produces pathogen-suppressing orfamide variants *in vitro*.

### (g) Testing *Pseudomonas* CHA0 biocontrol efficacy *in vivo* in a tomato system

The biocontrol potential of CHA0 was tested using the Micro Tom tomato cultivar infected by the UK *R. solanacearum* strains #1 and #7. In the absence of CHA0, the levels of disease incidence were higher when tomatoes were infected with strain #7 compared to strain #1 (*Ralstonia*:  $F_{1,34} = 48.788$ ,  $P = 0.315$ , Figure 4A). The CHA0 treatment did not affect wilting incidence compared to the water control treatment in the case of *Ralstonia* strain #1 (*Pseudomonas*:  $F_{1,16} = 24.731$ ,  $P = 1$ , Figure 4A). However, CHA0 reduced wilting incidence of *R. solanacearum* strain #7 (*Pseudomonas*:  $F_{1,16} = 17.736$ ,  $P = 0.01114$ , Figure 4A). The presence of CHA0 also increased the plant dry weight overall (*Pseudomonas*:  $F_{1,34} = 4.623$ ,  $P = 0.0387$ , Figure 4B) and this effect was the same for both *R. solanacearum* strains (*Ralstonia*:  $F_{1,34} = 1.668$ ,  $P = 0.205$ , Figure 4B). Together, these results suggest that CHA0 biocontrol efficacy was dependent on the *R. solanacearum* strain.

## Discussion

This study aimed to screen and identify effective *Pseudomonas* bacterial biocontrol strains against *Ralstonia solanacearum* species complex, understand potential underlying mechanisms of inhibition, and validate their efficacy *in vivo* in the tomato rhizosphere. *Pseudomonas* strain CHA0 was the most suppressive strain in both direct and indirect *R. solanacearum* inhibition assays. The comparative genomic analysis highlighted that metabolite clusters encoding DAPG, Pyoluteorin, and Orfamides A and B were unique for the most suppressive CHA0 *Pseudomonas* strain, and while only Orfamide A production was detected through mass spectrometry, extracted orfamide variants showed high inhibitory activity against *R. solanacearum*. *In vivo* tests revealed that CHA0 was effective at reducing bacterial wilt incidence. Interestingly, plant protection depended on the *R. solanacearum* strain identity and was only observed with one of the two tested strains. Despite the successful identification of potential *Pseudomonas* biocontrol species, more work is needed to harness their biocontrol activity in the plant rhizosphere.

When screening for the most effective *Pseudomonas* strains against *R. solanacearum* through direct and indirect assays, *P. protegens* strain CHA0 showed the most suppressive activity against all the tested *R. solanacearum* strains. The strong inhibition effects of CHA0 were observed through both direct and indirect assays. This result is in line with previous studies demonstrating the high biocontrol activity of CHA0 against various plant pathogens (Hu *et al.*, 2016). Although multiple *Pseudomonas* strains have been described in the literature with biocontrol abilities, CHA0 is one of the most well-established, studied, and successful biocontrol agents against plant-parasitic nematodes, fungal and bacterial pathogens (Siddiqui and Shaukat, 2003; Siddiqui and Shaukat, 2004; Jousset *et al.*, 2006; Humair *et al.*, 2009; Jamali *et al.*, 2009; Neidig *et al.*, 2011; Flury *et al.*,

2017). Interestingly, some less studied strains, such as MVP1-4 and Ph11C2, showed also good biocontrol potential and warrant more study in the future. Overall, all *Pseudomonas* strains suppressed all tested *R. solanacearum* strains to some extent *in vitro*. The likely explanation for this is that European *R. solanacearum* strains are likely to be very similar as they belong to a clonal lineage of Phylotype 2b strains that are adapted to grow in cold climates (Hayward, 1991). In contrast, biocontrol strains belonged to different *P. fluorescens* subgroups (e.g., *P. protegens* and *P. corrugata*) and originated from different countries across Europe and America. Hence, *Pseudomonas* strains were likely genetically more dissimilar compared to *R. solanacearum* strains. A more detailed analysis of European *R. solanacearum* strains is however required to link observed phenotypic similarities (equal susceptibility) and differences (infectivity *in vivo*) with underlying genetic differences

Comparative genomic analyses revealed that *Pseudomonas* strains CHA0 and Pf-5 (which belong to the *P. protegens* subgroup of *P. fluorescens*) had the greatest number of secondary metabolite clusters (17), while the other six *Pseudomonas* strains (the majority of which belong to the *P. corrugata* subgroup of *P. fluorescens*) harbored between 11 to 15 metabolic clusters. Based on previous literature, CHA0 and Pf-5 can produce a similar range of secondary metabolites such as pyoluteorin and cyclic lipopeptides, which could potentially explain why CHA0 exhibited the greatest inhibitory effects (Haas and Keel, 2003; Loper and Gross, 2007; Ma *et al.*, 2016). When testing these candidate metabolites on *R. solanacearum* growth, pyoluteorin stood out as the most suppressive compound, leading to the poorest *R. solanacearum* growth. Only *Pseudomonas* strains CHA0 and Pf-5 were found to harbor the T1PKS metabolic cluster, which encodes pyoluteorin production. However, no pyoluteorin was produced *in vitro* by either of the strains in the growth conditions used. Pyoluteorin inactivity in our experiments was potentially due to the use of a rich growth medium (Jamali *et al.*, 2009; Heidari-Tajabadi *et al.*, 2011; Yang *et al.*, 2018). Pyoluteorin is a chlorinated polyketide antibiotic and its production is regulated by the DAPG precursor, monoacetylphloroglucinol (Kidarsa *et al.*, 2011). It is thus possible that efficient production of pyoluteorin limits the production of DAPG, and that these compounds are not produced simultaneously. While CHA0 and Pf-5 strains harbored the highest amount of NRPS clusters (8 and 7, respectively), they were also associated with a wide variety of other putative metabolites that could have been linked with pathogen suppression (e.g., enantiopyochelin, rhizomide, and pyoverdine). Furthermore, we also isolated uncharacterized orfamamide variants from the CHA0 supernatant and observed highly inhibitory effects against the tested *R. solanacearum* strain. This suggests that orfamides at least partly explain the inhibitory capacity of the CHA0 strain in lab conditions. A wealth of other potential antimicrobial compounds was also identified in the CHA0 supernatant using non-targeted analysis. In the future, these secondary metabolites could be isolated

and tested in more detailed studies to identify potentially novel antimicrobials (Geudens and Martins, 2018). For example, Rose *et al.* (2021) have shown that Pf-5 has notable anti-algal properties when interacting with microalga *Chlamydomonas reinhardtii* through the production of rhizoxins and they were also able to detect the production of DAPG, pyrrolnitrin, pyoluteorin and orfamide A in their experiments. A key difference between ours and their study is that Pf-5 was grown in a TAP medium under continuous illumination which could have changed the Pf-5 metabolism and antimicrobial production (Rose *et al.*, 2021). It is thus possible that the use of rich culture media constrained the production of certain secondary metabolites in our assays. In the future, it will be important to characterize the secondary metabolite gene expression and metabolite production across different environments and to identify metabolite production potential in more realistic biocontrol conditions (Deveau *et al.*, 2016; Köhl, Kolnaar, and Ravensberg, 2019).

To validate *Pseudomonas* efficacy *in vivo*, we first tested that both selected *R. solanacearum* strains could infect tomato plants and found that they both caused around 50% bacterial wilt disease incidence. Such variation in disease incidence is typical for *R. solanacearum* as its virulence is determined by a combination of host immunity and environmental conditions that can vary considerably even in controlled greenhouse experiments (Hu *et al.*, 2016; Wei *et al.*, 2019). We found that bacterial wilt incidence was clearly reduced in the presence of CHA0 but only in the case of *R. solanacearum* strain #7. These results contrast with the *in vitro* results as *R. solanacearum* strains #1 and #7 did not show any differences in their susceptibility to DAPG, Pyoluteorin, or orfamide variants produced by CHA0. There are multiple potential biological explanations for this result. For example, it is possible that *Ralstonia* strain #1, was less able to colonize tomato roots, failed to express genes required for protection against *Pseudomonas* inhibition (e.g. efflux pumps or other deactivation of antimicrobials), or was not able to activate virulence gene expression (Ran *et al.*, 2005) in the presence of CHA0. Alternatively, the conditions used in tomato experiments may not have been optimal for *R. solanacearum* survival. Moreover, autoclaving of the soil could have influenced soil properties and microbial metabolism as heating soil over 120°C can result in increased levels of ammonium and nitrogen (Serrasolsas and Khanna, 1995). Together, these findings demonstrate the challenges of translating *in vitro* results to *in vivo* applications, highlighting the importance of studying biocontrol effects in more realistic *in vivo* conditions.

In conclusion, here we show that a combination of *in silico*, *in vitro*, and *in planta* approaches can be used to identify and validate effective biocontrol agents against *R. solanacearum*. By using simple microbiological assays based on direct and indirect interactions with the pathogen we were able to identify *P. protegens* CHA0 as the most inhibitory biocontrol strain. Comparative genomics was used to identify potential secondary metabolite clusters responsible for the inhibition and efficacy of identified compounds and their production was

validated in additional experiments. In addition to identified clusters, orfamide variants were also found to be highly inhibitory which could be characterized in the future (Keel *et al.*, 1992; Yasmin *et al.*, 2017). Based on these results, screening for Pyoluteorin and Orfamide secondary clusters could be used as a rapid way to identify effective biocontrol strains in the future. Such screening methods have been previously used to identify siderophore-producing biocontrol agents against rice fungal pathogens (Chaiharn, Chunnaleuchanon, and Lumyong, 2009). Despite the ability to produce a repertoire of secondary metabolites, CHA0 was able to reduce disease incidence with only one of the two tested *R. solanacearum* strains *in vivo*. Biocontrol effects should thus be studied in more realistic *in vivo* conditions in the future to test which mechanisms are expressed and active in the plant rhizosphere and if the natural plant microbiota is also affected by or affects the activity of identified biocontrol bacterial strains. Together, our results suggest that while the developed low-cost *in vitro* screening process can be used to identify effective biocontrol agents based on their secondary metabolism, further optimization is needed to predict the function of different strains in the plant rhizosphere.

### Author contributions

**Sophie Clough:** Conceptualization-Equal, Data curation-Equal, Formal analysis-Equal, Investigation-Equal, Methodology-Equal, Validation-Equal, Visualization-Equal, Writing – original draft-Equal, Writing – review & editing-Equal; **Alexandre Jousset:** Data curation-Equal, Investigation-Equal, Methodology-Equal, Writing – review & editing-Equal; **John Elphinstone:** Investigation-Equal, Supervision-Equal, Writing – review & editing-Equal; **Ville-Petri Friman:** Conceptualization-Equal, Formal analysis-Equal, Funding acquisition-Equal, Investigation-Equal, Project administration-Equal, Supervision-Equal, Writing – review & editing-Equal

### Acknowledgments

We thank Tony Larson and Swen Wagner of the CoEMS group for performing *Pseudomonas* metabolite identification and semi-quantification of select target compounds by ultra-performance liquid chromatography-mass spectrometry (UPLC-MS) analysis. S.C was funded by the White Rose Ph.D. Studentship Network and N8 AgriFood pump-priming grant at the University of Leeds. V-P.V. is funded by the Royal Society (grant nos. RSG\R1\180213 and CHL\R1\180031) and jointly by a grant from UKRI, Defra, and the Scottish Government, under the Strategic Priorities Fund Plant Bacterial Diseases program (BB/T010606/1).

### Conflict of interest

None declared.

## Data availability statement

Raw reads for all *Pseudomonas* strains are available at the NCBI Sequence Read Archive under BioProject accession number: PRJNA784243: <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA784243>

## Ethics statement

None required

## References

1. Almario, J., Bruto, M., Vacheron, J., Prigent-Combaret, C., Moëgne-Loccoz, Y. and Muller, D. (2017) Distribution of 2, 4-diacetylphloroglucinol biosynthetic genes among the *Pseudomonas* spp. reveals unexpected polyphyletism. *Frontiers in microbiology*, 8, p.1218.
2. Bangera, M. G. and Thomashow, L. S. (1999) Identification and characterization of a gene cluster for synthesis of the polyketide antibiotic 2, 4-diacetylphloroglucinol from *Pseudomonas fluorescens* Q2-87, *Journal of bacteriology*, 181(10), pp. 3155–3163.
3. Becker, J., Eisenhauer, N., Scheu, S. and Jousset, A. (2012) Increasing antagonistic interactions cause bacterial communities to collapse at high diversity, *Ecology Letters*, 15(5), pp.468-474, doi: 10.1111/j.1461-0248.2012.01759.
4. Blin, K., Wolf, T., Chevrette, M.G., Lu, X., Schwalen, C.J., Kautsar, S.A., Suarez Duran, H.G., de Los Santos, E.L., Kim, H.U., Nave, M. and Dickschat, J.S. (2017) antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification, *Nucleic acids research*, 45(W1), pp. W36–W41.
5. Blin, K., Shaw, S., Steinke, K., Villebro, R., Ziemert, N., Lee, S.Y., Medema, M.H. and Weber, T. (2019) antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline, *Nucleic acids research*, 47(W1), pp. W81–W87.
6. Caruso, P., Biosca, E.G., Bertolini, E., Marco-Noales, E., Gorris, M.T., Licciardello, C. and López, M.M. (2017) Genetic diversity reflects geographical origin of *Ralstonia solanacearum* strains isolated from plant and water sources in Spain. *International Microbiology*, 20(4), pp.155-164.
7. Castillo, J.A. and Greenberg, J.T. (2007) Evolutionary dynamics of *Ralstonia solanacearum*. *Applied and Environmental Microbiology*, 73(4), pp.1225-1238.
8. Chaiarn, M., Chunchaleuchanon, S. and Lumyong, S. (2009) Screening siderophore producing bacteria as potential biological control agent for fungal rice pathogens in Thailand, *World Journal of Microbiology and Biotechnology*, 25(11), pp. 1919–1928.
9. Chen, J., Yu, Y., Li, S. and Ding, W. (2016) Resveratrol and Coumarin: Novel Agricultural Antibacterial Agent against *Ralstonia solanacearum* In Vitro and In Vivo, *Molecules*, 21(11), p.1501,doi: 10.3390/molecules21111501
10. Clarke, C.R., Studholme, D.J., Hayes, B., Runde, B., Weisberg, A., Cai, R., Wroblewski, T., Daunay, M.C., Wicker, E., Castillo, J.A. and Vinatzer, B.A. (2015) Genome-enabled phylogeographic investigation of the quarantine pathogen *Ralstonia solanacearum* race 3 biovar 2 and screening for sources of resistance against its core effectors. *Phytopathology*, 105(5), pp.597-607.
11. Compant, S., Duffy, B., Nowak, J., Clément, C. and Barka, E.A. (2005) Use of Plant Growth-Promoting Bacteria for Biocontrol of Plant Diseases: Principles, Mechanisms of Action, and Future Prospects, *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 71(9), pp. 4951–4959. doi: 10.1128/AEM.71.9.4951–4959.2005
12. Cronin, D., Moenne-Loccoz, Y., Fenton, A., Dunne, C., Dowling, D.N. and O'gara, F.

- (1997) Role of 2,4-Diacetylphloroglucinol in the Interactions of the Biocontrol Pseudomonad Strain F113 with the Potato Cyst Nematode *Globodera rostochiensis*, *Applied and Environmental Microbiology*, 63(4), pp. 1357–1361.
13. Deveau, A., Gross, H., Palin, B., Mehnaz, S., Schnepf, M., Leblond, P., Dorrestein, P.C. and Aigle, B. (2016) Role of secondary metabolites in the interaction between *Pseudomonas fluorescens* and soil microorganisms under iron-limited conditions, *FEMS microbiology ecology*, 92(8).
  14. EPPO (2021) *Ralstonia solanacearum* species complex. EPPO datasheets on pests recommended for regulation. Available online. <https://gd.eppo.int>
  15. Flury, P., Vesga, P., Péchy-Tarr, M., Aellen, N., Dennert, F., Hofer, N., Kupferschmied, K.P., Kupferschmied, P., Metla, Z., Ma, Z. and Siegfried, S. (2017) Antimicrobial and insecticidal: cyclic lipopeptides and hydrogen cyanide produced by plant-beneficial *Pseudomonas* strains CHA0, CMR12a, and PCL1391 contribute to insect killing. *Frontiers in microbiology*, 8, p.100.
  16. Fravel, D.R. (2005) Commercialization and implementation of biocontrol. *Annu. Rev. Phytopathol.*, 43, pp.337-359.
  17. Genin, S. (2010) Molecular traits controlling host range and adaptation to plants in *Ralstonia solanacearum*, *New Phytologist*, pp. 920–928. doi: 10.1111/j.1469-8137.2010.03397.x.
  18. Genin, S. and Denny, T. P. (2012) Pathogenomics of the *Ralstonia solanacearum* Species Complex, *Annu. Rev. Phytopathol*, 50, pp. 67–89. doi: 10.1146/annurev-phyto-081211-173000.
  19. Geudens, N. and Martins, J.C., (2018) Cyclic lipodepsipeptides from *Pseudomonas* spp.–biological Swiss-army knives. *Frontiers in microbiology*, 9, p.1867.
  20. Gopalan-Nair, R., Jardinaud, M.F., Legrand, L., Landry, D., Barlet, X., Lopez-Roques, C., Vandecasteele, C., Bouchez, O., Genin, S. and Guidot, A. (2021) Convergent rewiring of the virulence regulatory network promotes adaptation of *Ralstonia solanacearum* on resistant tomato. *Molecular Biology and Evolution*, 38(5), pp.1792-1808.
  21. Govaerts, B., Fuentes, M., Mezzalama, M., Nicol, J.M., Deckers, J., Etchevers, J.D., Figueroa-Sandoval, B. and Sayre, K.D. (2007) Infiltration, soil moisture, root rot and nematode populations after 12 years of different tillage, residue and crop rotation managements, *Soil and Tillage Research*, 94(1), pp. 209–219. doi: 10.1016/j.still.2006.07.013.
  22. Gu, Y., Wei, Z., Wang, X., Friman, V.P., Huang, J., Wang, X., Mei, X., Xu, Y., Shen, Q. and Jousset, A. (2016) Pathogen invasion indirectly changes the composition of soil microbiome via shifts in root exudation profile. *Biology and Fertility of Soils*, 52(7), pp.997-1005.
  23. Guidot, A., Coupat, B., Fall, S., Prior, P. and Bertolla, F. (2009) Horizontal gene transfer between *Ralstonia solanacearum* strains detected by comparative genomic hybridization on microarrays. *The ISME journal*, 3(5), pp.549-562.
  24. Guidot, A., Jiang, W., Ferdy, J.B., Thébaud, C., Barberis, P., Gouzy, J. and Genin, S. (2014) Multihost experimental evolution of the pathogen *Ralstonia solanacearum* unveils genes involved in adaptation to plants. *Molecular biology and evolution*, 31(11), pp.2913-2928.
  25. Haas, D. and Défago, G. (2005) Biological control of soil-borne pathogens by fluorescent pseudomonads, *Nature Reviews Microbiology*, pp. 307–319. doi: 10.1038/nrmicro1129.
  26. Haas, D. and Keel, C. (2003) ‘Regulation of antiobiotic production in root-colonizing *Pseudomonas* spp. And relevance for biological control of plant disease, *Annu. Rev. Phytopathol*, 41, pp. 117–53. doi: 10.1146/annurev.phyto.41.052002.095656.
  27. Hansen, Z. R. and Keinath, A. P. (2013) Increased pepper yields following incorporation of biofumigation cover crops and the effects on soilborne pathogen populations and pepper diseases, *Applied soil ecology*, 63, pp. 67–77.

28. Hayward, A.C. (1991) Biology and epidemiology of bacterial wilt caused by *Pseudomonas solanacearum*. *Annual review of phytopathology*, 29, pp.65-87.
29. Heidari-Tajabadi, F., Ahmadzadeh, M., Moinzadeh, A. and Khezri, M. (2011) Influence of some culture media on antifungal activity of *Pseudomonas fluorescens* UTPF61 against the *Sclerotinia sclerotiorum*. *African Journal of Agricultural Research*, 6(30), pp.6340-6347.
30. Howell, C. R. and Stipanovic, R. D. (1979) Control of *Rhizoctonia solani* on cotton seedlings with *Pseudomonas fluorescens* and with an antibiotic produced by the bacterium, *Phytopathology*, 69(5), pp. 480–482.
31. Hu, J., Wei, Z., Friman, V.P., Gu, S.H., Wang, X.F., Eisenhauer, N., Yang, T.J., Ma, J., Shen, Q.R., Xu, Y.C. and Jousset, A. (2016) Probiotic diversity enhances rhizosphere microbiome function and plant disease suppression. *MBio*, 7(6).
32. Humair, B., González, N., Mossialos, D., Reimann, C. and Haas, D. (2009) Temperature-responsive sensing regulates biocontrol factor expression in *Pseudomonas fluorescens* CHA0. *The ISME journal*, 3(8), pp.955-965.
33. Jamali, F., Sharifi-Tehrani, A., Lutz, M.P. and Maurhofer, M. (2009) Influence of host plant genotype, presence of a pathogen, and coinoculation with *Pseudomonas fluorescens* strains on the rhizosphere expression of hydrogen cyanide-and 2, 4-diacetylphloroglucinol biosynthetic genes in *P. fluorescens* biocontrol strain CHA0. *Microbial ecology*, 57(2), pp.267-275.
34. Jousset, A., Lara, E., Wall, L.G. and Valverde, C. (2006) Secondary metabolites help biocontrol strain *Pseudomonas fluorescens* CHA0 to escape protozoan grazing. *Applied and Environmental Microbiology*, 72(11), pp.7083-7090.
35. Kaczmarek, A., Mackenzie, K., Kettle, H. and Blok, V.C. (2014) Influence of soil temperature on *Globodera rostochiensis* and *Globodera pallida*. *Phytopathologia Mediterranea*, pp.396-405.
36. Kaur, T., Rani, R. and Manhas, R.K. (2019) Biocontrol and plant growth promoting potential of phylogenetically new *Streptomyces* sp. MR14 of rhizospheric origin. *AMB Express*, 9(1), pp.1-14.
37. Keel, C., Schnider, U., Maurhofer, M., Voisard, C., Laville, J., Burger, U., Wirthner, P.J., Haas, D. and Défago, G. (1992) Suppression of root diseases by *Pseudomonas fluorescens* CHA0: importance of the bacterial secondary metabolite 2, 4-diacetylphloroglucinol. *Molecular Plant-Microbe Interactions*, 5(1), pp.4-13.
38. Kidarsa, T.A., Goebel, N.C., Zabriskie, T.M. and Loper, J.E. (2011) Phloroglucinol mediates cross-talk between the pyoluteorin and 2, 4-diacetylphloroglucinol biosynthetic pathways in *Pseudomonas fluorescens* Pf-5. *Molecular microbiology*, 81(2), pp.395-414.
39. Köhl, J., Kolnaar, R. and Ravensberg, W. J. (2019) Mode of Action of Microbial Biological Control Agents Against Plant Diseases: Relevance Beyond Efficacy, *Frontiers in Plant Science*, 10. doi: 10.3389/fpls.2019.00845.
40. Landa, B.B., Mavrodi, O.V., Raaijmakers, J.M., Gardener, B.B.M., Thomashow, L.S. and Weller, D.M. (2002) Differential ability of genotypes of 2, 4-diacetylphloroglucinol-producing *Pseudomonas fluorescens* strains to colonize the roots of pea plants. *Applied and Environmental Microbiology*, 68(7), pp.3226-3237.
41. Loper, J. E. and Gross, H. (2007) Genomic analysis of antifungal metabolite production by *Pseudomonas fluorescens* Pf-5, *European Journal of Plant Pathology*. pp. 9179-8, doi: 10.1007/s10658-007-.
42. Li, Y., Hu, Y., Bolund, L. and Wang, J. (2010) State of the art de novo assembly of human genomes from massively parallel sequencing data. *Human genomics*, 4(4), pp.1-7.
43. Ma, Z., Geudens, N., Kieu, N.P., Sinnaeve, D., Ongena, M., Martins, J.C. and Höfte, M. (2016) Biosynthesis, chemical structure, and structure-activity relationship of orfamide lipopeptides produced by *Pseudomonas protegens* and related species. *Frontiers in microbiology*, 7, p.382.
44. Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., Dow,

- M.A.X., Verdier, V., Beer, S.V., Machado, M.A. and Toth, I.A.N. (2012) Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular plant pathology*, 13(6), pp.614-629.
45. Nion, Y. A. and Toyota, K. (2015) Recent trends in control methods for bacterial wilt diseases caused by *Ralstonia solanacearum*, *Microbes and environments*. Japanese Society of Microbial Ecology, p. ME14144.
  46. Notz, R., Maurhofer, M., Schnider-Keel, U., Duffy, B., Haas, D. and Défago, G. (2001) Biotic factors affecting expression of the 2, 4-diacetylphloroglucinol biosynthesis gene phlA in *Pseudomonas fluorescens* biocontrol strain CHA0 in the rhizosphere. *Phytopathology*, 91(9), pp.873-881.
  47. Nagachandrabose, S. (2020) Management of Potato Cyst Nematodes Using Liquid Bioformulations of *Pseudomonas fluorescens*, *Purpureocillium lilacinum* and *Trichoderma viride*. *Potato Research*, pp.1-18.
  48. Natsch, A., Keel, C., Pfirter, H.A., Haas, D. and Défago, G. (1994) Contribution of the global regulator gene *gacA* to persistence and dissemination of *Pseudomonas fluorescens* biocontrol strain CHA0 introduced into soil microcosms. *Applied and Environmental Microbiology*, 60(7), pp.2553-2560
  49. Neidig, N., Paul, R.J., Scheu, S. and Jousset, A. (2011) Secondary metabolites of *Pseudomonas fluorescens* CHA0 drive complex non-trophic interactions with bacterivorous nematodes. *Microbial ecology*, 61(4), pp.853-859.
  50. Parkinson, N., Bryant, R., Bew, J., Conyers, C., Stones, R., Alcock, M. and Elphinstone, J. (2013) Application of variable-number tandem-repeat typing to discriminate *Ralstonia solanacearum* strains associated with English watercourses and disease outbreaks. *Applied and Environmental Microbiology*, 79(19), pp.6016-6022.
  51. Pedersen, A.L., Nybroe, O., Winding, A., Ekelund, F. and Bjørnlund, L. (2009) Bacterial feeders, the nematode *Caenorhabditis elegans* and the flagellate *Cercomonas longicauda*, have different effects on outcome of competition among the *Pseudomonas* biocontrol strains CHA0 and DSS73. *Microbial ecology*, 57(3), pp.501-509.
  52. Pierson III, L.S., Wood, D.W., Pierson, E.A. and Chancey, S.T. (1998) N-acyl-homoserine lactone-mediated gene regulation in biological control by fluorescent pseudomonads: Current knowledge and future work. *European journal of plant pathology*, 104(1), pp.1-9.
  53. Raaijmakers, J. M. and Weller, D. M. (1998) Natural plant protection by 2, 4-diacetylphloroglucinol-producing *Pseudomonas* spp. in take-all decline soils, *Molecular Plant-Microbe Interactions*, 11(2), pp. 144–152.
  54. Raaijmakers, J. M. and Weller, D. M. (2001) Exploiting Genotypic Diversity of 2, 4-Diacetylphloroglucinol-Producing *Pseudomonas* spp.: Characterization of Superior Root-Colonizing *P. fluorescens* Strain Q8r1-96, *Applied and Environmental Microbiology*, 67(6), pp. 2545–2554.
  55. Ran, L.X., Liu, C.Y., Wu, G.J., Van Loon, L.C. and Bakker, P.A.H.M. (2005) Suppression of bacterial wilt in *Eucalyptus urophylla* by fluorescent *Pseudomonas* spp. in China. *Biological Control*, 32(1), pp.111-120.
  56. Rose, M.M., Scheer, D., Hou, Y., Hotter, V.S., Komor, A.J., Aiyar, P., Scherlach, K., Vergara, F., Yan, Q., Loper, J.E. and Jakob, T. (2021) The bacterium *Pseudomonas protegens* antagonizes the microalga *Chlamydomonas reinhardtii* using a blend of toxins. *Environmental Microbiology*, 23(9), pp.5525-5540.
  57. Serrasolsas, I. and Khanna, P.K. (1995) Changes in heated and autoclaved forest soils of SE Australia. I. Carbon and nitrogen. *Biogeochemistry*, 29(1), pp.3-24.
  58. Shanahan, P., O'Sullivan, D.J., Simpson, P., Glennon, J.D. and O'Gara, F. (1992) Isolation of 2, 4-diacetylphloroglucinol from a fluorescent pseudomonad and investigation of physiological parameters influencing its production. *Applied and environmental microbiology*, 58(1), pp.353-358.
  59. Siddiqui, I. A. and Shaukat, S. S. (2003) Plant species, host age and host genotype effects on

- Meloidogyne incognita* biocontrol by *Pseudomonas fluorescens* strain CHA0 and its genetically-modified derivatives, *Journal of Phytopathology*, 151(4), pp. 231–238. doi: 10.1046/j.1439-0434.2003.00716.x.
60. Siddiqui, I. A. and Shaikat, S. S. (2004) Systemic Resistance in Tomato Induced by Biocontrol Bacteria Against the Root-Knot Nematode, *Meloidogyne javanica* is Independent of Salicylic Acid Production, *Journal of Phytopathology*, 152(1), pp. 48–54. doi: 10.1046/j.1439-0434.2003.00800.x.
61. Sofrata, A., Santangelo, E.M., Azeem, M., Borg-Karlson, A.K., Gustafsson, A. and Pütsep, K. (2011) Benzyl isothiocyanate, a major component from the roots of *Salvadora persica* is highly active against Gram-negative bacteria. *PLoS One*, 6(8), p.e23045.
62. Van der Putten, W.H., Cook, R., Costa, S., Davies, K.G., Fargette, M., Freitas, H., Hol, W.G., Kerry, B.R., Maher, N., Mateille, T. and Moens, M. (2006) Nematode interactions in nature: models for sustainable control of nematode pests of crop plants. *Advances in Agronomy*, 89, pp.227-260.
63. Wang, X., Wei, Z., Yang, K., Wang, J., Jousset, A., Xu, Y., Shen, Q. and Friman, V.P. (2019) Phage combination therapies for bacterial wilt disease in tomato. *Nature Biotechnology*, 37(12), pp.1513-1520.
64. Wei, Z., Gu, Y., Friman, V.P., Kowalchuk, G.A., Xu, Y., Shen, Q. and Jousset, A. (2019) Initial soil microbiome composition and functioning predetermine future plant health. *Science advances*, 5(9).
65. Weller, D.M. (2007) *Pseudomonas* biocontrol agents of soilborne pathogens: looking back over 30 years. *Phytopathology*, 97(2), pp.250-256.
66. Xue, Q.Y., Ding, G.C., Li, S.M., Yang, Y., Lan, C.Z., Guo, J.H. and Smalla, K. (2013) Rhizocompetence and antagonistic activity towards genetically diverse *Ralstonia solanacearum* strains—an improved strategy for selecting biocontrol agents. *Applied microbiology and biotechnology*, 97(3), pp.1361-1371.
67. Yang, C., Dong, Y., Friman, V.P., Jousset, A., Wei, Z., Xu, Y. and Shen, Q. (2019) Carbon resource richness shapes bacterial competitive interactions by alleviating growth-antibiosis trade-off. *Functional Ecology*, 33(5), pp.868-875.
68. Yang, T., Han, G., Yang, Q., Friman, V.P., Gu, S., Wei, Z., Kowalchuk, G.A., Xu, Y., Shen, Q. and Jousset, A. (2018) Resource stoichiometry shapes community invasion resistance via productivity-mediated species identity effects. *Proceedings of the Royal Society B*, 285(1893), p.20182035.
69. Yasmin, S., Hafeez, F.Y., Mirza, M.S., Rasul, M., Arshad, H.M., Zubair, M. and Iqbal, M. (2017) Biocontrol of bacterial leaf blight of rice and profiling of secondary metabolites produced by rhizospheric *Pseudomonas aeruginosa* BRp3. *Frontiers in Microbiology*, 8, p.1895.

## Figure legends

**Figure 1. Direct and indirect inhibition of *R. solanacearum* by *Pseudomonas* strains.** Panel (a) direct inhibition on soft agar assays, where each interaction represents a different *Pseudomonas* strain and the average diameter of the inhibition zones (mm) against different *R. solanacearum* strains after 96 h growth on soft agar lawns (N = 3). Panel (b) displays indirect inhibition of *R. solanacearum* by *Pseudomonas* strains' supernatant-LB mixes (set up in 96-well plates containing a 50:50 ratio of SN and 100% LB broth). The shade of red represents the growth reduction in *R. solanacearum* densities compared to the control (100% LB broth) treatment after 72 h of growth (N=3).

**Figure 2. Testing the inhibitory effects of identified *Pseudomonas* secondary metabolites on *R. solanacearum* growth.** Panel (a) shows the growth of seven *R. solanacearum* strains (in different panels) in various DAPG concentrations over time (1000  $\mu\text{M}$  – red, 500  $\mu\text{M}$  – blue, 100  $\mu\text{M}$  – green, 50  $\mu\text{M}$  – purple, and 0  $\mu\text{M}$  LB broth (0  $\mu\text{M}$  - control) - black). Panel (b) shows *R. solanacearum* strain #1 growth in the absence and presence of 100  $\mu\text{M}$  of orfamides A and B. Panel (c) shows the growth of *R. solanacearum* strains #1 and #7 in the absence and presence of 100  $\mu\text{M}$  of pyoluteorin. In all panels, bars show the standard error of the mean ( $\pm$ 1 SEM) based on 3 replicates.

**Figure 3. Testing the inhibitory effects of orfamide variants isolated from *Pseudomonas* CHA0 strain on *R. solanacearum* growth.** The black lines denote *R. solanacearum* density in the absence of the orfamide variants over time (control), while colored lines show the growth in the presence of orfamide variants isolated from CHA0. Orfamide variants 'A' and 'B' refer to two different fractions isolated separately during the extraction process. All error bars show the standard error of the mean ( $\pm$ 1 SEM) based on 3 replicates.

**Figure 4. Tomato plant infections by *R. solanacearum* strains #1 and #7 in the absence and presence of *Pseudomonas* strains.** Panel (a) displays the percentage of wilted plants (mean wilting incidence) and panel (b) the mean aboveground dry weight of individual plants at the end of the experiment for all treatments. All error bars show the standard error of the mean ( $\pm$ 1 SEM) based on 3 replicates (every replicate consisting of nine plants).

**Table 1. AntiSMASH5.0 cluster types recognized in eight *Pseudomonas* genomes used in this study including a brief definition of their function**

<i>Pseudomonas</i> Strain	Total no. of clusters	Cluster types	Description of metabolic clusters and examples of metabolites
CHA0	17	NRPS (8), Bacteriocin (2), CDPS (1), T1PKS (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Other (1)	- NRPS (Non-ribosomal peptide synthetase cluster) e.g. Orfamide, lipopeptide, viscosin, rhizomide, pyoverdine, enantiopyochelin
Pf-5	17	NRPS (7), NRPS-like (1), Bacteriocin (2), CDPS (1), T1PKS (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Betalactone (1), Other (1)	- NRPS-like (NRPS-like fragment) e.g. Mangotoxin, lankacidin - Bacteriocin (Unspecified ribosomally synthesized and post-translationally modified peptide product (RiPP) cluster)
Q2-87	15	NRPS (4), NRPS-like (1), Bacteriocin (3), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1), Ectoine (1), Lanthipeptide (1)	e.g. unknown - CDPS (tRNA-dependent cyclodipeptide synthases) e.g. Unknown
Q8R1-96	13	NRPS (6), NRPS-like (1), Bacteriocin (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1)	- T1PKS (Type I Polyketide synthase) e.g. Pyoluteorin - T3PKS (Type III Polyketide synthase) e.g. Unknown - NAGGN (N-acetylglutaminyglutamine amide)
IM1-96	12	NRPS (3), NRPS-like (1), Bacteriocin (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1), Lanthipeptide (2)	e.g. Unknown - Arylpolyene (Aryl polyene cluster) e.g. APE Vf - Butyrolactone (Butyrolactone cluster)
MVP1-4	13	NRPS (4), NRPS-like (1), Bacteriocin (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1), Lanthipeptide (2), LAP (1)	e.g. Unknown - Betalactone (Beta-lactone containing protease inhibitor) - Ectoine (Ectoine cluster) e.g. Unknown - Lanthipeptide (Lanthipeptide cluster)
F113	11	NRPS (2), NRPS-like (2), Bacteriocin (1), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1), Lanthipeptide (1)	e.g. Putative Class II - LAP (Linear azol(in)e-containing peptides) e.g. Unknown - Other (Unknown)
Ph11C2	14	NRPS (4), NRPS-like (1), Bacteriocin (3), T3PKS (1), NAGGN (1), Arylpolyene (1), Butyrolactone (1), Betalactone (1), Lanthipeptide (1)	e.g. Pyrrolnitrin

## APPENDIX

**Figure A1** Experimental design quantifying direct inhibition of *R. solanacearum* strain #1 by 4 types of *Pseudomonas* strains using soft agar overlay assay

**Figure A2** Mass spectrometry analysis indicating the presence of *Pseudomonas* metabolite orfamide A in the supernatant of strain CHA0. Orfamide A is identified with a monoisotopic mass ( $m/z$ ) of 1317.826  $[M+Na]^+$  and a retention time ( $t_R$ ) of 9.61 seconds at 15 ppm. This spectrum represents the replicate with the strongest peak out of the three replicates

**Table A1** *Pseudomonas* strains used in these experiments, and their geographical origin.

<i>Pseudomonas</i> strains	<i>Pseudomonas fluorescens</i> subgroup	<i>Pseudomonas</i> strains' origin
CHA0	( <i>P. protegens</i> )	Tobacco, Switzerland (Natsch <i>et al.</i> , 1994)
Pf-5	( <i>P. protegens</i> )	Cotton, USA (Howell and Stipanovic, 1979)
Q2-87	( <i>P. corrugata</i> )	Wheat, USA (Bangera and Thomashow, 1999)
Q8R1-96	( <i>P. corrugata</i> )	Wheat, USA (Raaijmakers and Weller, 1998)
1M1-96 ( <i>P. fluorescens</i> )	Not reported	Wheat, USA (Raaijmakers and Weller, 2001)
MVP1-4 ( <i>P. fluorescens</i> )	Not reported	Pea, USA (Landa <i>et al.</i> , 2002)
F113	( <i>P. corrugata</i> )	Sugar Beet, Ireland (Shanahan <i>et al.</i> , 1992)
Ph11C2 ( <i>P. fluorescens</i> )	Not reported	Tomato, France (Govaerts <i>et al.</i> , 2007)

**Table A2** *Ralstonia solanacearum* strains used in these experiments, and their geographical origin.

<i>Ralstonia</i> strains	York sample collection number	<i>Ralstonia</i> strains' origin
#1	YO352	River water, UK (2014)
#2	YO354	River water, UK (2015)
#3	YO355	River water, UK (2015)
#4	YO356	River water, UK (2013)
#5	YO162	Commercial culture, Poland (used for verification tests) (2014)
#6	YO351	River water, UK (2013)
#7	YO353	River water, UK (2014)

**Table A3 Media used for bacterial cultures.**

Media recipes	
LB Broth (1 Litre)	10 g Tryptone (pancreatic digest of casein) 5 g Yeast extract 5 g NaCl (+15 g agar for solid media)
CPG Broth (1 Litre)	1 g Casamino acids (casein hydrolysate) 10 g Peptone 5 g Glucose (+17 g agar for solid media)
NB Broth (1 Litre)	10 g Glucose 5 g Tryptone 3 g Beef extract 0.5 g Yeast extract
Kings Medium B Broth (1 Litre)	20 g Proteose peptone 1.5 g K <sub>2</sub> HPO <sub>4</sub> 1.5 g MgSO <sub>4</sub> •7H <sub>2</sub> O 10 mL Glycerol
Cryomedia for freezing bacterial stocks	50% v/v glycerol

**Table A4 AntiSMASH5.0 cluster types recognized in *Pseudomonas* CHA0 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
1.1	Bacteriocin	
1.2	CDPS	
1.3	NRPS	Orfamide (94%)
2.1	other	Pyrrrolnitrin (100%)
2.2	NRPS	Pyochelin (100%)
2.3	T1PKS	Pyoluteorin (100%)
3.1	NRPS	Lipopeptide (6%)
3.2	NAGGN	
3.3	NRPS	Pyoverdine (16%)
4.1	T3PKS	DAPG (100%)
4.2	Bacteriocin	
5.1	Arylpolyene	APE Vf (40%)
6.1	NRPS	Viscosin (31%) Pyoverdine (14%)
6.2	Betalactone	
8.1	NRPS	Pyoverdine (19%)
10.1	NRPS	Pyoverdine (3%)
13.1	NRPS	

**Table A5 AntiSMASH5.0 cluster types recognized in *Pseudomonas* Pf-5 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
1.1	NRPS	Pyoverdine (14%)
1.2	Betalactone	
1.3	Other	Pyrrolnitrin (100%)
1.4	NRPS	Pyochelin 100%
1.5	NRPS-like	Rhizoxin (100%)
1.6	T1PKS	Pyoluteorin (100%)
2.1	Bacteriocin	
2.2	CDPS	
2.3	NRPS	Orfamide (94%)
3.1	NRPS	Pyoverdine (21%)
3.2	NAGGN	
3.3	NRPS	Lipopeptide (6%)
4.1	T3PKS	DAPG (100%)
4.2	Bacteriocin	APE Vf (40%)
5.1	Arylpolyene	
7.1	NRPS	Pyoverdine (19%)
9.1	NRPS	Rhizomide (100%)

**Table A6 AntiSMASH5.0 cluster types recognized in *Pseudomonas* Q2-87 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
1.1	NAGGN	
1.2	NRPS	Pyoverdine (10%)
1.3	Betalactone	
2.1	Butyrolactone	
4.1	T3pks	DAPG (100%)
4.2	Bacteriocin	
5.1	Bacteriocin	
6.1	NRPS	Cupriachelin (35%)
7.1	Arylpolyene	APE vf (40%)
8.1	NRPS	Ishigamide (11%)
9.1	NRPS	Pyoverdine (19%)
9.2	Ectoine	
15.1	Bacteriocin	
19.1	Lanthipeptide	

**Table A7 AntiSMASH5.0 cluster types recognized in *Pseudomonas* Q8R196 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
1.1	NRPS	Pyoverdine (10%)
1.2	Betalactone	
1.3	T3PKS	DAPG (100%)
1.4	NRPS	Cupriachelin (11%)
1.5	Butyrolactone	
4.1	Arylpolyene	APE Vf (40%)
6.1	NAGGN	Taiwachelin (11%)
9.1	NRPS	Viscosin (31%) Pyoverdine (10%)
12.1	Bacteriocin	
20.1	NRPS-Like	Mangotoxin (71%)
23.1	NRPS	Pyoverdine (3%)
24.1	NRPS	Taiwachelin (11%) Pyoverdine (3%)
25.1	NRPS	

**Table A8 AntiSMASH5.0 cluster types recognized in *Pseudomonas* 1M1-96 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
2.1	T3PKS	DAPG (100%)
2.2	NRPS	Cupriachelin (11%)
2.3	Butyrolactone	
3.1	NAGGN	
4.1	NRPS	Serobactin (23%) Pyoverdine (10%)
8.1	Betalactone	
9.1	Bacteriocin	
17.1	Arylpolene	APE Vf (40%)
32.1	Lanthipeptide	Putative class II
44.1	NRPS-like	Mangotoxin (71%)
49.1	Lanthipeptide	Putative class II
62.1	NRPS	Viscosin (25%) Pyoverdine (6%)

**Table A9 AntiSMASH5.0 cluster types recognized in *Pseudomonas* MVP1-4 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
2.1	Betalactone	
2.2	LAP	
4.1	NRPS	Cupriachelin (11%)
4.2	T3PKS	DAPG (100%)
6.1	Arylpolyene	APE Vf (40%)
9.1	NRPS-like	Mangotoxin (71%)
10.1	Bacteriocin	
13.1	NAGGN	
24.1	Lanthipeptide	Putative class II
30.1	Butyrolactone	
32.1	NRPS	Viscosin (31%) Pyoverdine (10%)
35.1	NRPS	Pyoverdine (10%)
36.1	NRPS	Pyoverdine (8%) Serobactin (15%) Taiwachelin (11%)

**Table A10 AntiSMASH5.0 cluster types recognized in *Pseudomonas* F113 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
2.1	Arylpolyene	APE Vf (40%)
4.1	NAGGN	
4.2	NRPS	Pyoverdine (10%) Serobactins (23%)
5.1	NRPS-like	Lankacidin (26%)
6.1	Bacteriocin	
11.1	Betalactone	
12.1	T3PKS	DAPG (100%)
16.1	NRPS	Viscosin (31%) Pyoverdine (11%)
19.1	Butyrolactone	
20.1	Lantipeptide	

**Table A11 AntiSMASH5.0 cluster types recognized in *Pseudomonas* Ph11C2 genome including the types of clusters and % hit in the database with the most similar known cluster.**

Cluster	Type	Most similar known cluster
1.1	NAGGN	
1.2	NRPS	Serobactins (23%) Pyoverdine (10%)
1.3	Betalactone	
2.1	NRPS	
2.2	Bacteriocin	
2.3	Butyrolactone	
4.1	Bacteriocin	
5.1	NRPS-like	Mangotoxin (71%)
5.2	Bacteriocin	Pyoverdine (1%)
5.3	Arylpolyene	APE Vf (40%)
6.1	NRPS	WLIP (28%) Cupriachelin (11%) Pyoverdine (2%)
6.2	T3PKS	DAPG (100%)
7.1	NRPS	Viscosin (31%) Taiwachelin (27%) Pyoverdine (19%)
12.1	Lanthipeptide	Putative class II

**Table A12 Putative *Pseudomonas* metabolites produced in LB media identified using Progenesis software.**

Candidate compound	Metabolite function	Present in CHA0 supernatant	Present in Pf-5 supernatant
Indole-3-acetic acid	Auxin	YES	YES
Pyochelin	Siderophore	YES	
Rhizoxin	Interferes with mitosis by binding to <i>Beta</i> -tubulin affecting microtubule dynamics	YES	YES




Figure A1







Figure A2



## Inhibitory effect of DAPG

(b) Inhibitory effect of Orfamides (100 $\mu\text{M}$ )(c) Inhibitory effect of Pyoluteorin (100 $\mu\text{M}$ )