High prevalence of IgE sensitization to inactivated influenza vaccines, yet robust IgG4 responses, in a healthy pediatric population

Takao Fujisawa¹, Prince Baffour Tonto¹, Mizuho Nagao¹, Shigeru Suga¹, Kiyosu Taniguchi¹, Masahiro Hirayama², Tetsuo Nakayama³, and Takuji Kumagai⁴

¹Kokuritsu Byoin Kiko Mie Byoin ²Mie Daigaku ³Kitasato Daigaku Kansen Seigyo Kenkyu Kiko ⁴Kumagai Naika Shonika Iin

April 13, 2022

Abstract

Background: Anaphylaxis following influenza vaccination is a rare but serious problem. The underlying immune responses are not well understood. **Objective:** To elucidate IgE and IgG antibody responses in healthy children and adolescents following inactivated influenza vaccines (IIVs). **Methods:** The efficacy and safety of quadrivalent IIV (QIV) and trivalent IIV (TIV) were compared in healthy subjects aged 0-18 years. Serum IIV-specific IgE, IgG and IgG4 levels (sIgE, sIgG, sIgG4) were measured with ImmunoCAP. Hemagglutinin inhibition (HI) assay was performed for each influenza virus subtype. Sera from earlier patients who developed anaphylaxis to different IIVs were similarly tested. **Results**: A total of 393 subjects were enrolled: 96 were 6 months -2 years old, 100 were 3-5 years old, 100 were 6-12 years old, and 97 were 13-18 years old . No anaphylaxis was observed. Generally, QIV and TIV induced similar antibody responses. IIV-sIgE levels rose significantly after vaccination in the 6m-2y and 3-5y groups, did not change in the 6-12y group, and decreased in the 13-18y group. In contrast, the IIV-sIgG4/sIgE ratio increased significantly after vaccination in all age groups. Sensitized subjects had significantly higher HI titers and IIV-sIgG4/sIgE ratio in 5 patients with anaphylaxis was significantly lower than in age-matched healthy subjects. **Conclusion:** IIVs induce IgE sensitization in healthy children, but also robust IgG4 responses that may protect them from anaphylaxis. (239 words)

Hosted file

IV-IgE_Manucsript_Allergy_cleanCopy.docx available at https://authorea.com/users/382919/ articles/565205-high-prevalence-of-ige-sensitization-to-inactivated-influenza-vaccinesyet-robust-igg4-responses-in-a-healthy-pediatric-population




Figure 1_Tonto et al.




Figure 2_Tonto et al.




Figure 3_Tonto et al.




Figure 4_Tonto et al.




Figure 5_Tonto et al.