

Analysis of fifty-seven cases of nasal non-Hodgkin's lymphoma: a retrospective study

Yongya Du¹, Guotao Jia¹, Maocai Li¹, Yu Feng¹, Lili Gong¹, and Lianqing Li¹

¹Liaocheng People's Hospital

December 5, 2022

Abstract

*Primary nasal lymphoma is mainly non-Hodgkin lymphoma; NK/T-cell lymphoma is the most common type. *Nasal non-Hodgkin's lymphoma mainly occurs in middle-aged and older men. *Clinical symptoms of patients with nasal NK/T-cell lymphoma are unspecific and the early symptoms are atypical. The main signs, changes in nasal microbiota, nasal mucosa, and surrounding organs, are easily misdiagnosed as sinusitis and nasal polyps, complicating patient awareness at the early stage. *Nasal NK/T-cell lymphoma is rare and clinically highly aggressive with a high recurrence rate and poor prognosis. Therefore, early detection and diagnosis are crucial. *Clinicopathological manifestations of nasal non-Hodgkin lymphoma are complex, and the final pathological classification needs to be confirmed using pathology and immunohistochemistry.

Hosted file

Analysis of fifty-seven cases of nasal non-Hodgkin's lymphoma a retrospective study.docx available at <https://authorea.com/users/562915/articles/610156-analysis-of-fifty-seven-cases-of-nasal-non-hodgkin-s-lymphoma-a-retrospective-study>

Hosted file

Figure 1.docx available at <https://authorea.com/users/562915/articles/610156-analysis-of-fifty-seven-cases-of-nasal-non-hodgkin-s-lymphoma-a-retrospective-study>

Hosted file

Figure 2.docx available at <https://authorea.com/users/562915/articles/610156-analysis-of-fifty-seven-cases-of-nasal-non-hodgkin-s-lymphoma-a-retrospective-study>

Hosted file

Figure 3.docx available at <https://authorea.com/users/562915/articles/610156-analysis-of-fifty-seven-cases-of-nasal-non-hodgkin-s-lymphoma-a-retrospective-study>

Hosted file

Table 1.docx available at <https://authorea.com/users/562915/articles/610156-analysis-of-fifty-seven-cases-of-nasal-non-hodgkin-s-lymphoma-a-retrospective-study>