

# COVID-19 Pneumonia in Pregnancy: A Retrospective Study on Maternal and Neonatal Outcomes

Abdelhamid Benlghazi<sup>1</sup>, Belouad Moad<sup>1</sup>, Hanane Dabdi<sup>1</sup>, Bouhtouri Yassine<sup>1</sup>, Messaoudi Hamza<sup>1</sup>, Benali Saad<sup>1</sup>, Ait Bouhou rachid<sup>1</sup>, El mangoub Fatima<sup>1</sup>, Elhassani Mly El Mehdi<sup>1</sup>, and Kouach Jaouad<sup>1</sup>

<sup>1</sup>Universite Mohammed V de Rabat Faculte de Medecine et de Pharmacie de Rabat

July 31, 2023

## Abstract

**Objective:** To identify risk factors among pregnant with COVID- 19 for adverse outcomes related to disease severity, maternal morbidities, neonatal mortality and morbidity, and adverse birth outcomes. **Materials and methods:** In this retrospective study, 45 pregnant patients with COVID-19 pneumonia confirmed by RT-PCR were evaluated at HMIMV. The inclusion criteria were pregnant patients diagnosed with COVID-19 confirmed by RT-PCR and hospitalized in the gynecology-obstetrics and intensive care unit at Mohammed V Military Training Hospital (HMIMV). Exclusion criteria were non-pregnant patients and pneumonia cases with unconfirmed COVID-19 cause. Data was collected from patient records and telephone calls. The study used SPSS software to analyze the data. **Results:** Our study recorded 45 cases of SARS-CoV-2 infection in pregnant women over 2.5 years in the gynecology-obstetrics department at HMIMV. The age group most affected was 20-35 years, with 75% of cases. 57% of patients had no known comorbidities. 88.8% of patients were symptomatic at diagnosis, with fever, asthenia, headache, myalgia, anosmia, and ageusia being the most common symptoms. Almost 30% of patients required admission to the ICU, with 60% requiring oxygen supplementation. The study recorded 36 live births (80%), of which 26 cases (72.2%) required no further care and had a favorable outcome. **Conclusion:** Severe acute respiratory distress syndrome, which can be fatal, is a significant risk associated with COVID-19. Pregnant women with underlying medical conditions are more vulnerable to this complication and may benefit from preventive measures, such as lockdowns, to reduce their risk of infection. The severity of maternal disease appears to have an impact on obstetrical and neonatal outcomes, with potential complications including cesarean sections, neonatal prematurity, and admission to the intensive care unit .

## Introduction:

Since December 2019, cases of pneumonia have been reported in Wuhan, China. The number of cases increased rapidly in January 2020, leading the WHO to declare a public health emergency of international concern. In March 2020, due to the international spread and severity of the cases, the WHO qualified the situation as a COVID-19 pandemic (1).

Confirmed cases of COVID-19 vary clinically, from asymptomatic to severe. Pregnancy has been classified as a risk factor predisposing to critical forms of the disease (2).

Since the beginning of the pandemic, considerable effort has been devoted to the subject, in order to bring out the epidemiological and clinical data on maternal-fetal evolution, as well as the protocols adapted to the different forms of the disease in this category of patients. The aim of this study is to clarify the association between COVID-19 and pregnancy through a series of 45 cases recorded in the gynecology-obstetrics department at the Mohammed V Military Training Hospital (HMIMV) and to review the literature on the subject.

## Materials and methods:

We conducted a retrospective descriptive and analytical study of 45 patients who were registered in the gynecology-obstetrics department at HMIMV between April 1, 2020, and August 1, 2022. These pregnant patients were managed for COVID-19 pneumonia confirmed by real-time polymerase chain reaction (RT-PCR). The data to be analyzed were recorded on a data sheet, and we identified 45 files that met our inclusion criteria. For some patients, we made telephone calls to obtain the remaining information.

The inclusion criteria for our study were pregnant patients who had been diagnosed with COVID-19 infection confirmed by RT-PCR on nasopharyngeal swab samples and were hospitalized in the gynecology-obstetrics and intensive care unit at HMIMV. We only included cases that met these criteria in our evaluation form, which was filled in for each case while ensuring anonymity and encryption of the data.

Our exclusion criteria were non-pregnant patients and pneumonia cases with COVID-19 cause not confirmed by RT-PCR test. After collecting all data, we processed it using SPSS software and analyzed it using headcounts and percentages. This allowed us to draw meaningful conclusions from the data and better understand the impact of COVID-19 on pregnant patients

## Results:

Our study recorded 45 cases of SARS-CoV-2 infection in pregnant women over 2.5 years in the gynecology-obstetrics department at HMIMV. The average age of the patients was 36, and the age group most affected was 20-35 years, accounting for 75% of cases. In terms of comorbidities, 57% of patients had no known comorbidities, 15% had diabetes, and 11% had obesity. Gestational age at the time of COVID-19 infection was over 28 SA in 35 cases, accounting for 78%, 9 cases were between 28 and 14 SA, accounting for 20%, and only one case was less than 14 SA. Additionally, 37% of patients had received vaccination against COVID-19.(Table 1)

In terms of clinical symptoms, 88.8% of patients were symptomatic at the time of diagnosis, with general symptoms including fever (55.5%), asthenia (49%), headache (35.5%), myalgia (35.5%), anosmia (28%), and ageusia (12%). Cough was present in 40% of patients, sore throat in 13%, and dyspnea in 8.85%. On admission, vital signs were assessed for all patients, and hemodynamically, all patients had a correct mean arterial pressure, although tachycardia was noted in 12%. Respiratory dyspnea with desaturation to room air (Saturation <93%) was noted in 17.5%. The Glasgow score was 15/15 in all patients.(table 2)

Paraclinical findings from our study revealed that 15.6% of cases exhibited hyperleukocytosis (WBC > 15000 cells/ $\mu$ L), anemia was detected in 15% of cases, whereas lymphopenia and thrombocytopenia in two cases each. CRP was positive (> 10 mg/L) in 36% of patients, and LDH > 250 IU/l was noted in 22% of cases. Additionally, creatine kinase levels greater than 170 IU/l were detected in 4.5% of cases, and ferritin levels greater than 120 mg/l in 6.7% of cases.

All patients with signs of clinical or biological severity underwent a thoracic CT scan, with eight patients receiving this evaluation. Of those, two patients were classified as Corads 2, two as Corads 3, one as Corads 4, and three as Corads 5.

In terms of management and evolution, the study found that almost 30% of patients required admission to the intensive care unit, with 60% requiring oxygen supplementation. Of those patients, 49% received oxygen via high-concentration mask, 7% via non-invasive ventilation, and 4% via invasive ventilation. Azithromycin was used as a preventive antibiotic therapy in 65% of cases, while other antibiotics were prescribed in the event of signs of superinfection, including amoxicillin-clavulanic acid (20%), ceftriaxone (13%), and tavanic (2.2%).

Low-molecular-weight heparin was used as anticoagulation therapy in 51.2% of patients in a preventive dose and 15.5% in a curative dose, while 33.3% did not receive anticoagulation. The study found that 43 patients had a favorable outcome, but unfortunately, two deaths were noted. Additionally, two cases did not result in delivery, one due to abortion and the other due to maternal death before delivery. Of the deliveries, 20

were Caesarean sections, and 23 were vaginal deliveries. Over 50% of Caesarean sections were performed due to acute fetal distress.

Regarding pregnancy outcomes, the study recorded 36 live births (80%), of which 26 cases (72.2%) required no further care and had a favorable outcome. Ten cases (27.7%) were hospitalized in a neonatal unit, but also had a favorable outcome. Unfortunately, there were six cases (13.3%) of fetal death in utero (FIDU), one case of fetal death following maternal death, one case of death in a premature newborn, and one case of early abortion.

Low weight for gestational age was noted in almost 25% of cases. (Table 3)

## Discussion:

The COVID-19 pandemic has created an unprecedented global health crisis, with the rapid spread of the infection and the uncertainty surrounding its impact on pregnancy due to the lack of scientific data. Obstetricians have had to adapt their practices based on pragmatic approaches in response to this challenging situation (1).

Pregnancy triggers physiological changes that primarily affect the cardiorespiratory and immune systems. These changes include Th2 immunotolerance and alterations in the major histocompatibility complex expression at the maternal-fetal interface, making pregnant women more vulnerable to viral infections such as influenza or SARS-CoV-2 (2).

Pregnant women are at a heightened risk of developing respiratory pathologies and severe pneumonia due to their immunosuppressed state and adaptive physiological changes during pregnancy. These changes can make them more intolerant to hypoxia, putting them at higher risk during pandemics and other respiratory outbreaks.

Historical evidence, such as the influenza pandemic of 1918, highlights the increased vulnerability of pregnant women during pandemics. During that pandemic, pregnant women had a mortality rate of 2-6%, compared to the general population. Similarly, during the H1N1 2009 pandemic influenza virus outbreak, pregnant women faced a relative risk of 3-4 for complications from infection and were more than four times as likely as the general population to be admitted to the hospital.

A multicenter study conducted in France since the start of the COVID-19 pandemic examined data from 33 maternity clinics and found that 617 pregnant women tested positive for SARS-CoV-2, with the most commonly reported symptoms being cough (62.2%), fever (46.2%), anosmia (27.2%), dyspnea (26.7%), and diarrhea (8.8%). Of the pregnant women, 20.7% required respiratory assistance, with 4.7% requiring mechanical ventilation, 1% needing extracorporeal membrane oxygenation (ECMO), and 0.2% died. Risk factors for needing respiratory assistance included age over 35, high pre-pregnancy body mass index ( $>30 \text{ kg/m}^2$ ), pre-existing diabetes or a history of pre-eclampsia, and a current diagnosis of gestational hypertension or pre-eclampsia. These risk factors are similar to those found in the general population.

A meta-analysis published in September 2020 that included 77 studies found that pregnant women with COVID-19 had a 40% incidence of fever and a 39% incidence of cough. They were more likely to require intensive care unit admission and invasive ventilation. The analysis found 73 deaths (0.1%). Risk factors for severe disease included age, high body mass index, chronic hypertension, and diabetes. Women with pre-existing comorbidities had a higher risk of ICU admission (OR = 4.21) (5).

In a case-control study comparing data from 8,207 pregnant women and 83,205 non-pregnant women with confirmed SARS-CoV-2 infection, it was found that pregnant women had a higher risk of intensive care hospitalization (RR 1.2) and mechanical ventilation (RR 1.9) compared to their non-pregnant counterparts. However, there was no significant difference in mortality risk between pregnant and non-pregnant women with the virus.

Various biological abnormalities have been observed in individuals with COVID-19, as per existing literature. One such abnormality is lymphopenia, which can be identified in patients as early as the viral disease phase.

During the first week of illness, elevated levels of transaminases may also be observed. In the second week, which is considered the inflammatory response phase, markers such as CRP, procalcitonin, and ferritin may increase (7).

Special attention is required in managing pregnant women with COVID-19, considering the potential impact of pregnancy on the respiratory and cardiovascular systems. Close monitoring is particularly crucial during the first two weeks of the disease's evolution, as rapid deterioration can occur during this period.

Hospitalization of pregnant women with confirmed or suspected COVID-19 is not always required, except in cases where there is a need for oxygen support. In situations where symptoms are not severe, ambulatory management with symptomatic treatment is generally advised. However, it is crucial to educate the woman about the symptoms that require medical reassessment, such as fever, cough, and dyspnea. Self-medication should be avoided.

Hospitalization is necessary in cases where severe symptoms develop rapidly, such as a respiratory rate of 25 cycles per minute or higher and oxygen saturation levels of less than 95% in ambient air. Management in the hospital may involve a thoracic CT scan, oxygen therapy, invasive ventilation, and corticosteroid therapy to reduce inflammation when C-reactive protein (CRP) levels are greater than 50(2).

Oxygen therapy is initiated based on respiratory rate and saturation levels. The threshold for initiating oxygen therapy is when SpO<sub>2</sub> falls below 95%.

According to the latest recommendations, antibiotic therapy is not recommended for COVID-19 unless there is a documented co-infection (7).

SARS-CoV-2 infection increases the risk of thrombosis in patients, and therefore, it is crucial to systematically assess the thrombotic risk in all patients with confirmed infection. The completion of ongoing studies, including the FREEDOM COVID-19 trial, is necessary to determine if therapeutic-dose anticoagulation provides additional efficacy in reducing thrombotic events, preventing intubation, or improving survival compared to prophylactic-dose anticoagulation in hospitalized patients (9).

Vitamin D supplementation may be beneficial for COVID-19 patients with vitamin D deficiency or insufficiency, but there is no evidence to support its use in the prevention or reduction of disease severity in individuals with normal blood vitamin D levels (10). Beigmohammadi et al. found that vitamin supplementation led to significant changes in serum levels of various vitamins and inflammatory markers, and reduced the rate of hospitalization lasting more than 7 days, but did not have a significant effect on mortality (11).

According to the PAN-COVID study, which is an international multicenter study conducted in 43 centers across 18 different countries to evaluate the pregnancy and neonatal outcomes of women with COVID-19. The study found that infection during pregnancy is associated with preterm birth, primarily due to fetal distress. Although the incidence of small for gestational age and fetal growth restriction was not higher than expected, there was a significant difference in the proportion of participants affected by stillbirth and FGR, depending on whether they delivered within two weeks or after. To address this, clinicians should have a low threshold for delivery if they detect concerns with fetal movements or heart rate monitoring during this period.

The study also found that SARS-CoV-2 infection during pregnancy does not seem to affect birth weight or increase the risk of congenital malformations. However, the effect of infection on miscarriage was not determined. The rate of pre-eclampsia among the study participants was not higher than expected, and neonatal infection was uncommon(12).

### **Conclusion:**

Even though pregnant women are already recognized as a high-risk group by the WHO and should have equal access to safe and effective preventive and therapeutic measures, pregnant women with additional risk factors require special attention. Evidence suggests that advanced maternal age and high body mass index are associated with a greater risk of severe COVID-19 in pregnant women. Furthermore, pregnant women

with COVID-19 have been found to have higher rates of preterm birth compared to those without the disease. Therefore, it is crucial that healthcare providers closely monitor pregnant women with these risk factors and provide appropriate care to mitigate the risk of adverse outcomes.

### Références :

1. Benlghazi A, Benali S, Bouhtouri Y, Belouad M, Massoudi H, Kouach J. Infection SARS-CoV-2 chez la femme enceinte; profil épidémiologique, clinique, biologique et évolutifs, à propos de 16 cas: expérience de l'Hôpital Militaire Marocain COVID-19 de Benslimane. *The Pan African Medical Journal*. 2021;38.
2. Bertholom C. Covid-19 et grossesse. *Option/Bio*. 2022;32(653):15.
3. Jamieson DJ, Honein MA, Rasmussen SA, Williams JL, Swerdlow DL, Biggerstaff MS, et al. H1N1 2009 influenza virus infection during pregnancy in the USA. *The Lancet*. 2009;374(9688):451-8.
4. Kayem G, Lecarpentier E, Deruelle P, Bretelle F, Azria E, Blanc J, et al. A snapshot of the Covid-19 pandemic among pregnant women in France. *Journal of gynecology obstetrics and human reproduction*. 2020;49(7):101826.
5. Allotey J, Stallings E, Bonet M, Yap M, Chatterjee S, Kew T, et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *bmj*. 2020;370.
6. Ellington S, Strid P, Tong VT, Woodworth K, Galang RR, Zambrano LD, et al. Characteristics of women of reproductive age with laboratory-confirmed SARS-CoV-2 infection by pregnancy status—United States, January 22–June 7, 2020. *Morbidity and Mortality Weekly Report*. 2020;69(25):769.
7. Moretto F, Sixt T, Abdallahoui M, Devilliers H, Chavanet P, Catherine F, et al. Intérêt des antibiotiques au cours de la COVID-19. *Médecine et Maladies Infectieuses*. 2020;50(6):S93.
8. Peyronnet V, Sibiude J, Huissoud C, Lescure F-X, Lucet J-C, Mandelbrot L, et al. Infection par le SARS-CoV-2 chez les femmes enceintes. Actualisation de l'état des connaissances et de la proposition de prise en charge. *CNGOF. Gynécologie Obstétrique Fertilité & Sénologie*. 2020;48(12):858-70.
9. Farkouh ME, Stone GW, Lala A, Bagiella E, Moreno PR, Nadkarni GN, et al. Anticoagulation in patients with COVID-19: JACC Review Topic of the Week. *Journal of the American College of Cardiology*. 2022;79(9):917-28.
10. Pereira M, Dantas Damascena A, Galvão Azevedo LM, de Almeida Oliveira T, da Mota Santana J. Vitamin D deficiency aggravates COVID-19: systematic review and meta-analysis. *Critical reviews in food science and nutrition*. 2022;62(5):1308-16.
11. Beigmohammadi MT, Bitarafan S, Hoseindokht A, Abdollahi A, Amoozadeh L, Soltani D. The effect of supplementation with vitamins A, B, C, D, and E on disease severity and inflammatory responses in patients with COVID-19: a randomized clinical trial. *Trials*. 2021;22(1):1-9.
12. Mullins E, Perry A, Banerjee J, Townson J, Grozeva D, Milton R, et al. Pregnancy and neonatal outcomes of COVID-19: The PAN-COVID study. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2022;276:161-7.
13. Smith ER, Oakley E, Grandner GW, Rukundo G, Farooq F, Ferguson K, et al. Clinical risk factors of adverse outcomes among women with COVID-19 in the pregnancy and postpartum period: a sequential, prospective meta-analysis. *American journal of obstetrics and gynecology*. 2023;228(2):161-77.

### Hosted file

Table 1 Maternal comorbidities.docx available at <https://authorea.com/users/644676/articles/657470-covid-19-pneumonia-in-pregnancy-a-retrospective-study-on-maternal-and-neonatal-outcomes>

### Hosted file

Table 2 Maternal symptoms at presentation.docx available at <https://authorea.com/users/644676/articles/657470-covid-19-pneumonia-in-pregnancy-a-retrospective-study-on-maternal-and-neonatal-outcomes>

### Hosted file

Table 3 -Key perinatal outcomes .docx available at <https://authorea.com/users/644676/articles/657470-covid-19-pneumonia-in-pregnancy-a-retrospective-study-on-maternal-and-neonatal-outcomes>