# How Fern and Fern Allies Respond to Heterogeneous Habitat— A Case in Yuanjiang Dry-Hot Valley

Feng-Chun Yang<sup>1</sup>, Chaya Sarathchandra<sup>2</sup>, Jing-Xin Liu<sup>3</sup>, Hua-Ping Huang<sup>4</sup>, Jian-Yong Gou<sup>5</sup>, Suthathong Homya<sup>6</sup>, Kritana Prueksakorn<sup>7</sup>, Jun Zhao<sup>1</sup>, and Ming-Fu Yang<sup>8</sup>

<sup>1</sup>Yibin Vocational and Technical College
<sup>2</sup>Research Centre for Eco-Environmental Sciences Chinese Academy of Sciences
<sup>3</sup>Environmental Education Center, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences
<sup>4</sup>Chinese Academy of Tropical Agricultural Sciences
<sup>5</sup>Honghe Meteorological Bureau
<sup>6</sup>Department of Environmental Science, Faculty of Science and Technology, Phuket Rajabhat University
<sup>7</sup>Faculty of Environment and Resource Studies, Mahidol University
<sup>8</sup>Yanyuan Vocational Middle School

April 05, 2024

#### Abstract

The Yuanjiang dry-hot valley features hot and dry climate, low vegetation and soil degradation. It had lush vegetation in the past, but has become degraded in recent decades. Understanding the interrelationship between species and the habitat is necessary to explain this change. In this study, a link between fern and fern allies - a group that is hypersensitive to environmental factors and their circumstances is constructed. Intensive transects and plots were designed to be proxies for extant fern and fern allies, and their habitats. Fifty years of meteorological records of precipitation and temperature along altitude and river running direction (latitudinal) were employed. Alpha and beta diversity are used to access diversity. Species\_estimated, Singletons, Uniques, ACE, ICE, and Chao2, which associate to abundance and rarity, are subscribed to the correlation between fern and fern allies, and their ecosystem. Eight species, Selaginella pseudopaleifera, Aleuritopteris squamosa, Adiantum malesianum, Pteris vittata, Davallia trichomanoides, Sinephropteris delavayi, Selaginella jugorum, and Lygodium japonicum are used as indicators of a typical xeric and sun-drying habitat. The results indicate (1) accompanied by dramatically shrinking habitats, fern and fern allies are in very low diversity and abundance, whereas the rarity is relatively high; (2) for fern and fern allies, environmental factors are positive when altitude goes up; and (3) eight indicator species are latitudinally correlated with fern and fern allies along the river running direction.

How Fern and Fern Allies Respond to Heterogeneous Habitat— A Case in Yuanjiang Dry-Hot Valley

Feng-Chun Yang  $^{1, 2, 3}$ , Chaya Sarathchandra $^{4, 5}$ , Jing-Xin Liu $^6$ , Hua-Ping Huang<sup>7</sup>, Jian-Yong Gou $^8$ , Ye Li<sup>7</sup>, Xiao-Ye Mao $^1$ , Hui-Ting Wen<sup>1</sup>, Suthathong Homya $^9$ , Kritana Prueksakorn $^{10}$ \*, Jun Zhao $^{1, 3}$  and Ming-Fu Yang $^{3, 11}$ 

<sup>1</sup> Yibin Vocational and Technical College, Yibin, Sichuan, 644000, China

 $^2$ School of Chinese Medicinal Material Resources, Guang<br/>dong Pharmaceutical University, Guangzhou, 510006, China

 $^3$  Yi Minority Culture Research Center of the Key Research Base of Philosophy and Social Sciences of Sichuan Province

<sup>4</sup> State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

<sup>5</sup> Department of Biological Science, Faculty of Applied Sciences, Rajarata University of Sri Lanka, Mihintale, 50300, Sri Lanka

<sup>6</sup> Environmental Education Center, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla Yunnan, 666100, China; xtbg\_liujx@qq.com

<sup>7</sup> Environment and Plant Protection Research Institute, Chinese Academy of Tropical Agricultural Sciences, Haikou, 570100, China; hhp18@163.com

<sup>8</sup> Honghe Meteorological Bureau, Yunnan, 661100, China; 125529676@qq.com

<sup>9</sup> Department of Environmental Science, Faculty of Science and Technology, Phuket Rajabhat University, 83120, Thailand; suthathong.h@pkru.ac.th

<sup>10</sup> Faculty of Environment and Resource Studies, Mahidol University, Nakhon Phathom, 73170, Thailand

<sup>11</sup> Yanyuan Vocational Middle School of Liangshan Yi Autonomous Prefecture, Sichuan, 615500, China

First author: yangfengchun@gdpu.edu.cn

Co-first author: chayasarathchandra@gmail.com

Correspondence: kritana.pru@mahidol.ac.th

**Abstract:** The Yuanjiang dry-hot valley features hot and dry climate, low vegetation and soil degradation. It had lush vegetation in the past, but has become degraded in recent decades. Understanding the interrelationship between species and the habitat is necessary to explain this change. In this study, a link between fern and fern allies - a group that is hypersensitive to environmental factors and their circumstances is constructed. Intensive transects and plots were designed to be proxies for extant fern and fern allies, and their habitats. Fifty years of meteorological records of precipitation and temperature along altitude and river running direction (latitudinal) were employed. Alpha and beta diversity are used to access diversity. Species\_estimated, Singletons, Uniques, ACE, ICE, and Chao2, which associate to abundance and rarity, are subscribed to the correlation between fern and fern allies, and their ecosystem. Eight species, *Selaginella pseudopaleifera, Aleuritopteris squamosa, Adiantum malesianum, Pteris vittata, Davallia trichomanoides, Sinephropteris delavayi, Selaginella jugorum,* and *Lygodium japonicum* are used as indicators of a typical xeric and sun-drying habitat. The results indicate (1) accompanied by dramatically shrinking habitats, fern and fern allies are in very low diversity and abundance, whereas the rarity is relatively high; (2) for fern and fern allies, environmental factors are positive when altitude goes up; and (3) eight indicator species are latitudinally correlated with fern and fern allies along the river running direction.

Keywords: diversity; dry-hot valley; fern; indicator; rarity

## 1. Introduction

The dry-hot valley is represented by high temperature and dry air throughout the year, and it is one of the arid ecosystems in the world influenced by climate change (Dong *et al*., 2014; Wang *et al*., 2016; Srivastava *et al*., 2018). Climate change has resulted in extensive alteration of terrestrial ecosystems, including the change in biogeographic distribution and biological diversity, and the interaction between organisms and their living circumstances (Hea *et al*., 2018). Understanding the interrelationship between plants and their environmental conditions in arid ecosystems is a preliminary step to interpret the mechanism of adaptation, or giving a feasible solution for saving earthbound environment (Jina *et al*., 2019).

Yuanjiang dry-hot valley is located in the upper and middle region of Yuanjiang-Red River, an international river shared by China and Vietnam (6. GMS Operations Center, 2017). In the valley, the nature of soil, water and fertilizer are not in good condition (Su *et al*., 2015; Yang *et al*., 2018). The vegetation has changed gradually in the past 500 years, but has been degraded significantly in recent decades causing a dry and hot tropical ecosystem (Xu *et al*., 1985). Broadleaf and evergreen plants have been shrinking, while it is currently dominated by montane savannah, covered with few trees and high grasses (Zhang *et al*., 2009). To elucidate this change, it is necessary to understand a corresponding pattern between plants and environmental factors such as humidity, temperature and water availability. Fern and fern allies are traditionally considered hypersusceptible to environmental changes; their occurrence or disappearance is closely correlated to their living place (Abotsi *et al*., 2020). In this study, the relationship between fern and fern allies and environments in the Yuanjiang dry-hot valley is investigated together with species population and environment factors.

#### 2. Materials & Methods

## 2.1. Transect and plot designing

The research site is the core area of Yuanjiang dry-hot valley with a length of 105.26 km along the river channel, oriented northwest to southeast (Figure 1, A). Six transects were arranged randomly (Figure 1, B) with a concept of no interference by human activities, where three transects were set up altitudinally along a mountain slope, and the other three latitudinally along the riverside (Table 1). Altitudinal transects were in height vertically, with ten plots arranged equidistantly 20 m apart from the lower plot to the higher. The latitudinal transects were in length along the river channel, and ten plots were spaced at a distance of 100 m evenly (Figure 2). Plots were 2 m  $\times$  2 m squared. All fern species in the sampling sites were investigated and nomenclature following *Flora Yunnanica* (Wuet al., 2006).

Table 1. Location, altitude and circumstance, and orient of each transect

Tr.	Location	Habitat	Orient
A	23°28'05.22"N, 102deg10'43.32"E	$360\pm15$ m <i>a.s.l.</i> , open river terrace	Latitudinal
В	23°19'13.98"N, 102deg32'45.12"E	490~690  m a.s.l., closed ravine	Altitudinal
$\mathbf{C}$	23°17'17.03"N, 102deg39'59.50"E	440~640  m a.s.l., open ravine	Altitudinal
D	$23^{\circ}11'51.97"$ N, $102 deg 53'57.82"$ E	$260\pm15$ m <i>a.s.l.</i> , closed river terrace	Latitudinal
$\mathbf{E}$	$23^{\circ}04'29.90"$ N, $103$ deg $11'40.64"$ E	360~560  m a.s.l., closed ridge	Altitudinal
F	$23^{\circ}03'20.57"$ N, $103$ deg $16'38.56"$ E	$230\pm15$ m <i>a.s.l.</i> , closed river terrace	Latitudinal

## (Tr.: Transect).



**FIGURE 1** Watershed of Yuanjiang-Red River and transects A-F in the dry-hot valley.**FIGURE 2** Transects designing along altitude and latitude.

2.2. Meteorological data collecting

Meteorological records, primarily annual temperature and precipitation, were collected from long-time meteorology stations in the Yuanjiang dry-hot valley at 312 m to 1758 m a.s.l. (Supplementary data: Table A). Temperature and precipitation are considered the main environmental factors correlated to fern and fern allies growth and distribution in the dry-hot valley. Linear regression is conducted based on twenty stations.

## 2.3. Statistical analyses

Alpha and beta biodiversity (a - and  $\beta$  -index) are both employed. The a -index, associating to species, is commonly represented by Shannon-Wiener index (H), Evenness (J), and Dominance. The $\beta$  -index, which is based on not only species occurrence but also abundance analytically, and associated to living habitat, is represented by *Bray-Curtis* (Magurran, 2004; Chao & Chiu, 2016; Ricotta & Podani, 2017). When comparing *turnover* and *nestedness* components (Baselga, 2017), it is more approachable to use *Bray-Curtis* similarity index for analysis of abundance and incidence, following the strategy of Diserud and Ødegaard (Diserud & Ødegaard, 2007).

Additionally, there are approaches to assess the difference in plots and transects using the program EstimateS Version 9.1 (Colwell, 2019). Species\_estimated, Singletons, Uniques, ACE, ICE, and Chao2 are prevalently accepted in diverse studies, such as savanna ants of Australia (Andersen et al., 2015), the elevational richness of Colorado Mountain (Szewczyk & McCain, 2019) and many different areas (Rajakaruna et al., 2016; Chao et al., 2017; Girardello, et al., 2019), with details as follows: 1) Species\_estimated indicates the expected number of species in t pooled plots. It is not a real but a calculated index. Increase of species approaches zero early or late indicating different habitats; 2) Singletons indicates the number of species with only one individual in t pooled plots. It is associated with the species rarity; 3) Uniques indicates the number of species richness; 5) ICE is an incidence coverage-based estimator of species richness; 5) ICE is an incidence coverage-based estimator of species richness; 5) ICE is an incidence coverage-based estimator of species richness; 6) Chao2 is a species richness estimator (Chao, 1984, 1987; Rajakaruna et al., 2016). These six indices are applied in this study, and their model inferences are tested on Chi-square statistic ( $\chi^2$ ) at 0.05 significance level. Residuals are evaluated for normality using the Shapiro-Wilk test (normality was assumed when P [?] 0.5) (Shapiro & Wilk, 1965). With Tukey's method, multiple comparisons are used to test the disparity between habitats at 0.05 significant levels.

Therefore, to cope with the issues mentioned above, two kinds of species matrices are required. The first matrix is individual-based, while the second matrix is occurrence-based ignoring the number of individuals. All matrices are computed 999 times in EstimateS (version 9.10) and then the selected index is rearranged and analysed with packages vegan(Oksanen *et al.*, 2011), labdsv (Roberts, 2018) and lme4 (Pinheiro, 2011) in R (v.2.15.2) (R Development Core Team, 2012). Indicator value function (IndVal) in package indicspecies (Cáceres & Legendre, 2009) is also employed.IndVal is a random variable that takes "1" to represent a happened event and "0" for nothing. After that, multilevel pattern analysis is conducted at significance level P = 0.05 in the pooled species.

#### 3. Results

The analytical results are from 2016 fern individuals belonging to 17 genera and 13 families (Supplementary data: Table B).

3.1. Spatial patterns of thermal and moisture in Yuanjiang dry-hot valley

Temperature and precipitation are the two main climatic factors being considered in this study. According to a 57 year (1962-2018) data profile, the temperature decreases when altitude increases (Figure 3). The maximum annual temperature was 24.5 at 401 m *a.s.l*. while the minimum was 16.3 at 1,635 m *a.s.l*. (Supplementary data: Table A). On the contrary, precipitation rose when altitude increased (Figure 3). The maximum annual precipitation was 2,159 mm at 1,263 m*a.s.l*. while the minimum was 738 mm at 401 m *a.s.l*. (Supplementary data: Table A).

Rainfall in Yuanjiang dry-hot valley depends on two factors. The first is the process of wet air sinking, pressurizing and heating-up when warm and humid airflow moves down to the valley, which weakens the

uplifting of water vapor in the precipitation system and reduces water vapor condensation (Zhang *et al* ., 2009). Secondly, typical valley wind is generated by consistent sunshine, high temperature and rapid evaporation. Water vapor is taken to a relatively high altitude by the up-valley wind and then freezes and falls to the ground in the forms of droplets (Giovannini *et al* ., 2017). Therefore, the valley's weather condition is cool and humid at the relatively high altitude, but it is reversed in the lower place. Fifty year records in the Jinsha River valley display a similar statistical profile (Zhang *et al* ., 2009). Dry air and high temperature prevail as typical environmental features in the dry-hot valley.

FIGURE 3 Correlation of temperature and precipitation in altitudal gradients.

3.2. Shannon-Wiener index (H) and diversity variation along mountain slope and riverine separately

Shannon-Wiener index among plots in each transect are displayed separately in Figure 4 with a general linear regression. The changes along plots can be noticed, and the habitat heterogeneity can also be identified based on these dispersing values.

## (A) Altitudinal transects (mountain slope, Figure 4, B/C/E)

Plots in transect are labelled 1-10 from the bottom to the top of the mountain. Transect B was a closed ravine while C was open. Transect E was a mountain ridge. In general, there was a positive correlation between the H index and the altitude. It clearly shows a tendency that fern and fern allies became more diverse as the altitude increased.

Transect B had a significant correlation (P < 0.05). It was a ravine with a closed habitat and vegetative plants. The humidity inside the ravine was relatively high due to well irrigating from the upper site. Most fern and fern allies grew well in this circumstance.

Transect C was also a ravine but aggregated by fragment rocks, and poor vegetation. It could not keep groundwater, neither a close and humid habitat. Fern and fern allies were rarely discovered in this site.

Transect E was a mountain ridge which could not maintain water consistently, nor construct a humid habitat. Xerophytes bushes were well developed on the ridge compared to fern and fern allies.

## (B) Latitudinal transects (riverside, Figure 4, A/D/F)

Plots in transects were labelled 1-10 from the upper to the lower across 900 meters. Transect A was an open river terrace while D and F were closed. The H index was positively and significantly correlated along the river (P < 0.05). Compared to altitudinal transects, the water supply of transects A/D/F seemed sufficient because it was close to the river channel. Simultaneously, closed habitats were constructed in some areas in transect D and F due to developed shrubs and trees. However, transect A did not develop flourishing riverine vegetation. The main reason would be ascribed to irregular river tides.

**FIGURE 4** The variance of Shannon-Wiener index (H) in transects. (Al. = Altitudinal, La. = Latitudinal)

3.3.  $\beta$ -diversity and the habitat heterogeneity in the valley

## (A) In transects

Although all the six transects existed in a narrow valley of 105 km in length, the *Bray-Curtis* index showed a deviation altitudinally and latitudinally (Table 2). The maximum was 0.292 in latitudinal transect D and F which defined similar living circumstances of fern and fern allies. The minimum was 0 in A and D, A and F, and A and E which defined different circumstances.

 Table 2. Similarity in transects.

	$\mathbf{A}$	В	С	D	Ε	$\mathbf{F}$
Α		0.148	0.092	0	0	0
В			0.059	0.108	0.01	0.11

	Α	в	$\mathbf{C}$	D	$\mathbf{E}$	$\mathbf{F}$
$\overline{\mathbf{C}}$				0.125	0.222	0.039
D					0.073	0.292
$\mathbf{E}$						0.02
$\mathbf{F}$						

(A, D, and F, latitudinal transects; B, C, and E, altitudinal transects).

## (B) In plots

To specify the microhabitat in the plots of each transect, *Bray-Curtis* indices were calculated. The values are listed separately (Table 3). Transects A, B, D, and F were similar in the plots with *Bray-Curtis*, and the living circumstance of fern and fern allies was continuous. In contrast, Bray-Curtis in the plots of transects C and E was approaching 0. This is due to the poor records of target species and the same low humidity in these two places. The results match well with the field investigation.

 Table 3. Similarity in plots

Transect A	Transect A	Transect A	Transect A	Transect A	Transect A	Transect A	Transect A
1 2 3 4 5 6 7 8 9 10	1	<b>2</b> 0.8	<b>3</b> 0.59 0.44	<b>4</b> 0.55 0.41 0.82	<b>5</b> 0.53 0.39 0.65 0.82	<b>6</b> 0.57 0.42 0.68 0.87 0.95	7 0.59 0.44 0.67 0.82 0.92 0.93
Transect C 1 2 3 4 5 6 7 8 9 10	Transect C 1	Transect C 2 0	Transect C 3 0 0.73	Transect C 4 0 0	<b>Transect C</b> <b>5</b> 0 0 0 0	<b>Transect C</b> <b>6</b> 0 0 0 0 0	Transect C 7 0 0 0 0 0 0
Transect E 1 2 3 4 5 6 7	Transect E 1	Transect E 2 0	Transect E 3 0 0	<b>Transect E 4</b> 0 0 0	<b>Transect E 5</b> 0 0 0 0 0	<b>Transect E 6</b> 0 0 0 0 0 0 0 0	<b>Transect E</b> 7 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0

Transect A	$\mathbf{Transect}~\mathbf{A}$	${\rm Transect}~{\bf A}$	${\rm Transect}~{\bf A}$	${\bf Transect}~{\bf A}$	${\rm Transect}~{\bf A}$	$\mathbf{Transect}~\mathbf{A}$	Transect A
8							
9							
10							

## 3.4. The variation of Species\_estimated, Singletons, Uniques, ACE, ICE, and Chao2

 $Species\_estimated$  (Figure 5): Transects A and C approached a gentle level earlier than others. The estimated species curve reaches zero prematurely indicating a homogeneous habitat among the plots investigated. There were three species in transect A: *Aleuritopteris squamosa*, *Selaginella stauntoniana*, and *S. pseudopaleifera*; while there were two species in transect C, *Pteris Vittata* and *A. squamosa*. Species increase in transect A and C stopped at the plots 3 and 2 separately. There were no new species discovered in the following plots. In contrast, species accumulation curves in the other four transects keep going up from plot 1 to plot 10, showing a high possibility to find new species in the next plot. This indicates a heterogeneity of habitat among the plots.

Singletons (Figure 5): It was zero in transects A and C because no species grew as a single individual. In transect F, the Singletons approached 1, and there was only one species (*Histiopteris incise*) found in plot 4. Unlike transects A, C, and F, Singletons increased quickly without a turnover in transect B, D, and E. The maximum of Singletons happens in transect B bearing approximately nine species, *Pteris vittata, P. ensiformis ,P. excelsa , P. fauriei , Lygodium japonicum, Selaginella delicatula , Adiantum philippense , Microlepia speluncae, and Phymatosorus scolopendria .* These species are generally distributed worldwide in a broad niche. However, none of them colonize the whole plot. Instead, other species take advantage. Extreme water reliance takes charge of this. Water suspended in the air and absorbed in the living substance is vital to the growing and regeneration process.

Uniques (Figure 5): According to the curves, transects A and C were in low Unique . Transect A was riverine; plots in this transect had similar habitats. Transect C was a ravine along a mountain slope, and it featured fragmented rocks and bare surface soil from plot 1 at the bottom to plot 10 at the top. Plots in this transect had similar habitats too. Therefore, there were very few species found in these two transects, and these species were presented in the plots from 1-10 evenly. In opposite to transects A and C, Unique increased in the plots of transects B, D, E, and F, indicating that more species and habitats were expected in the next plot. Transects D and F, for example, were riverine transects, the same with transect A, but the growing substance and canopy coverage made the habitat more suitable for fern and fern allies. When the investigated plots increased continuously, new records were found occupying a single plot or sharing a plot with other species, e.g., Selaginella helferi, S. biformis, S. doederleinii, Histiopteris incisa, and Equisetum hyemale.

ACE, ICE, and Chao2 (Figure 5): They verified each other by displaying similar curve shapes. Species richness in transect B, D, and F were much higher than other transects. The maximum richness was 22 species detected in transect B whereas only two species were detected in transect C.

**FIGURE 5** The variance of estimated species, rarity, and abundance in plots 1-10 of each transect. (Red = transect A; blue = transect B; black = transect C; green = transect D; orange = transect E; purple = transect F)

## 3.5. Indicator species

Eight species are screened out to represent the Yuanjiang dry-hot valley as shown in Table 4. All indicators perched at the altitudes  $490 \degree 690$  m although some of them distributed down to 260 m. All indicators require a humid environment although they grow in different river segments (Figure 6).

Table 4. Indicator candidates in Yuanjiang dry-hot valley.

Code	Species	Altitude and habitat	Transect	p-value	
1	Selaginella pseudopaleifera	$360^{\circ}690 \text{m} a.s.l.$ , hot and wet	1,2	0.001 ***	
2	Aleuritopteris squamosa	$360^{\circ}690 \text{m} a.s.l.$ , hot and dry	1,2,3	0.001 ***	
3	Adiantum malesianum	$260^{\circ}690m \ a.s.l.$ , hot, limestone	2,4,5,6	0.001 ***	
4	Pteris vittata	$260^{\circ}690\text{m} a.s.l.$ , acidic soil	2,3,4,5,6	0.001 ***	
5	$Davallia\ trichomanoides$	$330^{\circ}690 \text{m} a.s.l.$ , hot and wet	6	0.001 ***	
6	Sinephropteris delavayi	$490^{\circ}690 \text{m} a.s.l.$ , hot and wet	2	0.002 **	
7	Selaginella jugorum	$490^{\sim}690$ m <i>a.s.l.</i> , hot and wet	2	0.003 **	
8	Lygodium japonicum	490~690 m $\mathit{a.s.l.},$ dry acidic soil	2	0.018 *	

**FIGURE 6** Indicator species in different segments of Yuanjiang dry-hot valley. (A) Upper stream, transect A; (B) Middle stream, transect B; (C) Downstream, transect F; (D) Aleuritopteris squamosa indicator of transect A; (E) Sinephropteris delavayindicator of transect B; (F) Selaginella pseudopaleiferaindicator of transect F.

## 4. Discussion

#### 4.1. Variation of thermal and moisture altitudinally

The importance of altitudes in plant distribution is usually ignored. However, some studies have paid attention to altitudinal influence on vascular plants. In the tropical area, studies reveal a hump-shaped pattern with the highest diversity at mid-altitudes and then decrease toward both high and low altitudes (Hernández-Rojas *et al.*, 2018). The altitude where maximum fern and fern allies diversity occurs differs somehow among mountain ranges. For example, maximum diversity occurs around 1,800 m in both Costa Rica (Kluge *et al.*, 2006) and Mount Kinabalu, Borneo (Kessler *et al.*, 2001), 2,000 m in Bolivia (Salazar *et al.*, 2015), and 2,400 m on Mount Kilimanjaro, Tanzania (Kessler *et al.*, 2011). Climatically, these gradients corresponded to the upper parts of tropical gradients where richness also declined. Species were comparable between these data sets at the same mean annual temperature. In temperate regions, richness was reported to decline continuously with elevation or remains roughly constant, such as New Zealand or North America (Brock *et al.*, 2016). In subtropical regions, a study from the Fanjingshan of Guizhou, China revealed a strong negative correlation between altitude and species (Wang *et al.*, 2008).

The altitudinal transects along mountain slope were located from  $360^{-}690 \text{ m } a.s.l$ . (table 1). The annual temperature was over 22.3 <sup>O</sup>C, and annual precipitation was less than 984 mm (Supplementary data: Table A). Species regeneration was difficult for most species. Generally, fern and fern allies were rarely discovered. in which two species were discovered in transect C, and three species were in transect E. Transect B was different from C and E. In contrast, it was rich in fern and fern allies with 22 species and 971 individuals (Supplementary data: Table B). Such a difference could be explained with topographic features. The altitude of transects C and E was around 970 m a.s.l., where annual temperature and precipitation were 20.8 <sup>O</sup>C and 785 mm, respectively. In contrast, the altitude of transect B was around 1750 m a.s.l., where annual temperature and precipitation were 16.8  $\sim$  17.3 °C and 1211  $\sim$  1312 mm, respectively (Supplementary data: Table A). Rainfall drained into the ravine and supports the plants inside. Water supply in transect B was better than others because there was plenty of water supply coming from the higher elevation. Relative humidity was recorded at above 90% in transect B, which was more humid than other transects (around 72%). Comparing to the drought in transects C and E, much more shrubs (Salix myrtillacea, Ilex cornuta, Buddleja officinalis, Solanum verbascifolium etc.) and trees (Phyllanthus emblica, Trema tomentosa, Grewia biloba var. parviflora, Broussonetia papyrifera etc.) were developed. A closed canopy was composed in densely vegetates. Diversity in the closed ravine was higher than an open ravine or a ridge.

4.2. Water condition determines the distribution of fern and fern allies

The population of species acquiring high water supply had shrunk in Yuanjiang dry-hot valley for centuries (Xu, 1985). Undoubtedly, fern populations were distributed broadly in the past compared to the mosaic

habitats nowadays. Water condition has traditionally been considered a decisive factor for fern and fern allies (Kluge *et al*., 2006; Yang *et al*., 2011). The fern and fern allies flourishment is interpreted as a reflection of environmental humidity (Yang *et al*., 2011) (or an optimal combination of mild temperature and humidity (Kluge *et al*., 2006). Compared to ground-living fern and fern allies, epiphytic species strongly rely on water air in the closed-canopy (Kreft *et al*., 2010). In this study, ground-living fern and fern allies did not display a different water requirement. Instead, both of them performed similar positive correlations to the water condition. It turns out that 1980 individuals (98%) were growing well in humid transects, i.e., A, B, D, and F (A, D, F benifited from river flow, while transect B benifited from inner cycling precipitation), whereas 36 individuals (2%) lived in dry transects. Considering transect B, the plot at the bottom preserved 54 individuals of 3 species. It increased to 364 individuals of 19 species such as *Pteris vittata* and*Adiantum malesianum*. However, it was altered to uncommonly seen species when the altitude went up, where the dominant species were*Sinephropteris delavayi* and *Phymatosorus cuspidatus*. The reason was ascribed to inner cycling precipitation at high altitude.

## 4.3. Heterogeneous habitat and fern and fern allies distribution

Generally, to maintain plants growing and population size, fern and fern allies have to absorb more water from the circumstance surroundings, e.g., Pteris vittata, Lygodium japonicum, and Microlepia speluncae. However, other species, such as SelaginellaS delicatula, S. uncinata, Adiantum capillus-veneris, and A. ma*lesianum*, survive in a unique approach, *e.q.*, slender plant size and vegetative reproduction by rhizomes. Their rarity in the plots is ascribed to frequent hot and dry air and habitat deterioration such as substance rocks fragmentation and surface soil erosion. It becomes worse in recent decades and has resulted in a dramatically declined environment. Reproductive organisms, e.g., spores, gemmae, and slender rhizomes, suffer and regeneration cannot proceed under such environmental stress. Habitat determines which plants grow. while environmental factors define a habitat. According to the habitat heterogeneity hypothesis (Tuanmu & Jetz, 2015; Hamm & Drossel, 2017), the living requirement differs in species, and each species has their living requirements exclusively. Their requirements include living substances, water, temperature, and nutrition, to name a few. A complicated and diverse habitat supports more species while a unitary habitat supports less. Diversity increases in more heterogeneous habitats (Negrão et al., 2017). For an arid ecosystem, disturbances with varying intensities and spatial scales are responsible for habitat patterns (Torimaru et al ., 2018). As regards the fern and fern allies distribution in Yunnan, China, heterogeneity was reported to be critical to the diversity of species and to be related to biogeographic zonation (Zhang et al., 2017). In Yuanjiang dry-hot valley, Bray-Curtis index revealed that transects C and E were unitary habitats, while transects A, B, D, and F were more complicated and diverse. In the field survey, only three species were recorded in the unitary transects C and E, whereas 33 species were found in heterogeneous transects A, B, D, and F (Supplementary data: Table B). A Topographic variation on altitudinal and latitudinal gradients and divergent temperature and water condition generated a mosaic microclimate on the slopes in the valley.

#### 4.4. Fern and fern allies indicators

Indicator species are correlated to specific environmental factors. The population sizes usually increase or decrease due to the change of one or several factors. Their presence or absence is mostly determined by an environmental factor. They are optimal to indicate the environmental evolution process regarding their diverse habitat requirements in species (Karst *et al*., 2005; Yang & Grote, 2018; Yang *et al*., 2019). Most fern and fern allies are strictly limited in their habitats because of their intrinsic sensitivity to environmental change. For example, *Gonocormus minutus* live on water due to their fragile mesophyll tissue, while *Selaginella tamariscina* can resist extreme arid at Yang tribute due to its resuscitation, etc. They are considered key indicators of the environment (Silva *et al*., 2018).

In this study, screened indicators fit well with the hot-dry or hot-wet environment (Table 4). Three species were worldwide distributed with broad ecological amplitude, i.e., *Pteris vittata*, *Lygodium japonicum*, and *Adiantum malesianum*.

However, five other species were limited in distribution and confined to narrow amplitude, i.e., *Selaginella pseudopaleifera*, *S. jugorum*, *Sinephropteris delavayi*, *Davallia trichomanoides*, and *Aleuritopteris squamosa*. Although growing in different river segments from upper to lower, they all occurred in relatively humid areas and disappeared in hot and dry places. Similar results revealed the indicator function in these selected genera, e.g., endemic species of genus *Selaginella* in Philippine are recognised as indicators of refuge in the geological past (Tan, 2013) and *Davallia mariesii* is used as an indicator of landforms and vegetation types (Park *et al.*, 2019). *A. squamosa* is endemic in the upper and middle segment of Yuanjiang-Red River, which strongly associates to dry-hot climate (Wu, 1981). Meanwhile, a cryptic species *A. argentea*, is reported as a pyrophytic species and *Sinephropteris* is monotypic confined to karst in southwest China, northeast India, and north Burma (Pallvi *et al.*, 2019).

## 5. Conclusions

In recent years, climate change is broadly focused, while environmental disasters have occurred frequently. Significant land degradation and climate events have been reported for more than decades. Plants are considered as a serious part of environmental change. In response to such ecological crises, this research demonstrates such changes in Yuanjiang valley located in the first half part of a river shared by China and Vietnam by focusing on fern and fern allies affected by other environmental factors. From bottom to top of the mountain, precipitation increased while the temperature decreased. Fern and fern allies are positively correlated with height as a result of atmospheric conditions. Species richness increases further accompanied habitat heterogeneity in the valley. Even with no influence of human disturbance, these current appearances are not the original pattern of the areas, indicating another consequence of climate change.

Acknowledgement: This research is supported by the Research Foundation for Advanced Talents of Yibin Vocational and Technical College (No. ybzysc20bk03), the Innovation Project of the Department of Education of Guangdong Province, China (No. 2019KTSCX71), Scientific Research Platform of Yibin Vocational and Technical College (No. ybzy21kypt05), Project of Science and Technology Innovation Team (No. ybzy21cxtd-04) and Yi Minority Culture Research Center of the Key Research Base of Philosophy and Social Sciences of Sichuan Province (No. YZWH2101). The authors would like to thank Ms. Hong Anh Thi Nguyen for general assistance and also acknowledge Faculty of Environment and Resource studies, Mahidol University's editing service for support in the improvement of this manuscript.

Data Accessibility: All sampling data are online available with https://doi.org/10.5061/dryad.k0p2ngf8h.

## References

Abotsi, K. E., Bose, R., Adjossou, K., Deblauwe ,V., Rouhan, G., Segla, K. N., Atsri, K. H., & Kokou, K. (2020). Ecological drivers of pteridophyte diversity and distribution in Togo (West Africa). *Ecological Indicators*, 108, 105741. DOI: 10.1016/j.ecolind.2019.105741

Andersen, A. N., Del Toro, I., & Parr, C. L. (2015). Savanna ant species richness is maintained along a bioclimatic gradient of increasing latitude and decreasing rainfall in northern Australia. *Journal of Biogeography*, 42(12), 2313-2322. DOI: 10.1111/jbi.12599

Baselga, A. (2017). Partitioning abundance-based multiple-site dissimilarity into components: Balanced variation in abundance and abundance gradients. *Methods in Ecology and Evolution*, 8(7), 799-808.

Brock, J. M., Perry, G. L., Lee, W. G., & Burns, B. R. (2016). Tree fern ecology in New Zealand: A model for southern temperate rainforests. *Forest Ecology and Management*, 375, 112-126. DOI: 10.1016/j.foreco.2016.05.030

Caceres, M., Legendre, P. (2009). Associations between species and groups of sites: Indices and statistical inference. *Ecology*, 90,3566–3574. DOI: 10.1890/08-1823.1

Chao, A. (1984). Nonparametric estimation of the number of classes in a population. *Scandinavian Journal* of *Statistics*, 11, 265-270.

Chao, A. (1987). Estimating the population size for capture-recapture data with unequal catchability. *Biometrics*, 43, 783-791. DOI: 10.2307/2531532

Chao, A. & Chiu, C.H. (2016) Bridging two major approaches (the variance framework and diversity decomposition) to beta diversity and related similarity and differentiation measures. *Methods in Ecology and Evolution*, 8, 919-928.

Chao, A., Colwell, R. K., Chiu, C. H., & Townsend, D. (2017). Seen once or more than once: Applying Good–Turing theory to estimate species richness using only unique observations and a species list. *Methods in Ecology and Evolution*, 8(10), 1221-1232. DOI: 10.1111/2041-210X.12768

Colwell, R.K. (2019, April 4). EstimateS: Statistical Estimation of Species Richness and Shared Species from Samples. Version 9.1. [Software]. Retrieved from http://viceroy.eeb.uconn.edu/estimates/EstimateSPages/AboutEstimateS.htm.

Diserud, O.H. & Odegaard, F. (2007) A multiple-site similarity measure. Biology Letters, 3, 20-22.

Dong, Y., Xiong, D., Su, Z. A., Li, J., Yang, D., Shi, L., & Liu, G. (2014). The distribution of and factors influencing the vegetation in a gully in the Dry-hot Valley of southwest China. *Catena*, 116, 60-67. DOI: 10.1016/j.catena.2013.12.009

Girardello, M., Chapman, A., Dennis, R., Kaila, L., Borges, P. A., & Santangeli, A. (2019). Gaps in butterfly inventory data: A global analysis. *Biological Conservation*, 236, 289-295. DOI: 10.1016/j.biocon.2019.05.053

Giovannini, L., Laiti, L., Serafin, S., & Zardi, D. (2017). The thermally driven diurnal wind system of the Adige Valley in the Italian Alps. *Quarterly Journal of the Royal Meteorological Society*, 143(707), 2389-2402. DOI: 10.1002/qj.3092

GMS Operations Center. (2017). Greater Mekong Subregion Atlas of the Environment (2nd Edition). derived online: http://portal.gms-eoc.org

Hamm, M., & Drossel, B. (2017). Habitat heterogeneity hypothesis and edge effects in model metacommunities. *Journal of theoretical biology*, 426, 40-48. DOI: 10.1016/j.jtbi.2017.05.022

Hea, Z, Dua, J., Chena, L., Zhua, X., Lina, P., Zhao, M., Fan, S. (2018). Impacts of recent climate extremes on spring phenology in arid-mountain ecosystems in China. *Agricultural and Forest Meteorology*, 260-261, 31-40. DOI: 10.1016/j.agrformet.2018.05.022

Hernandez-Rojas, A., Kessler, M., Kromer, T., Carvajal-Hernandez, C., Weigand, A., & Kluge, J. (2018). Richness Patterns of Ferns Along an Elevational Gradient in the Sierra de Juarez, Oaxaca, Mexico: a Comparison with Central and South America. *American Fern Journal*, 108 (3), 76-95. DOI: 10.1640/0002-8444-108.3.76

Jina, J., Wang, Q., Wanga, J., & Otienoc D. (2019). Tracing water and energy fluxes and reflectance in an arid ecosystem using the integrated model SCOPE. *Journal of Environmental Management*, 231, 1082-1090. DOI: 10.1016/j.jenvman.2018.10.090

Karst, J., Gilbert, B., & Lechowicz, M. J. (2005). Fern community assembly: the roles of chance and the environment at local and intermediate scales. *Ecology*, 86(9), 2473-2486. DOI: 10.1890/04-1420

Kessler, M., Parris, B. S., & Kessler, E. (2001). A comparison of the tropical montane pteridophyte floras of Mount Kinabalu, Borneo, and Parque Nacional Carrasco, Bolivia. *Journal of Biogeography*, 28(5), 611-622. DOI: 10.1046/j.1365-2699.2001.00577.x

Kessler, M., Kluge, J., Hemp, A., & Ohlemuller, R. (2011). A global comparative analysis of elevational species richness patterns of ferns. *Global Ecology and Biogeography*, 20(6), 868-880. DOI: 10.1111/j.1466-8238.2011.00653.x

Kluge, J., Kessler, M., & Dunn, R. R. (2006). What drives elevational patterns of diversity? A test of geometric constraints, climate and species pool effects for pteridophytes on an elevational gradient in Costa Rica. *Global Ecology and Biogeography*, 15(4), 358-371. DOI: 10.1111/j.1466-822X.2006.00223.x

Kreft, H., Jetz, W., Mutke, J., & Barthlott, W. (2010). Contrasting environmental and regional effects on global pteridophyte and seed plant diversity. *Ecography*, 33(2), 408-419. DOI: 10.1111/j.1600-0587.2010.06434.x

Magurran, A. E. (2004). *Measuring biological diversity*. Blackwell, New York.

Negrao, R., Sampaio-e-Silva, T., Kortz, A. R., Magurran, A., & Matos, D. M. S. (2017). An endangered tree fern increases beta-diversity at a fine scale in the Atlantic Forest Ecosystem. *Flora*, 234, 1-6. DOI: 10.1016/j.flora.2017.05.020

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., & Wagner, H. (2011). vegan: Community Ecology Package. R Package Version 1.17-8. [Software]. Retrieved from http://CRAN.R-project.org/package=vegan.

Pallvi, S., Chander, D. H., & Bikarma, S. (2019). Four pyrophytic pteridophytes in the Chir pine forest of Kalidhar forest range in Shiwaliks of North West Himalaya in Indian Himalayan region. *Environment Conservation Journal*, 20 (3), 73-77. DOI: 10.36953/ECJ.2019.20310.

Park, B. J., Byeon, J. G., & Cheon, K. I. (2019). Study of Ecological Niche and Indicator Species by Landforms and Altitude of Forest Vegetation in Mt. Myeonbong. Korean Journal of Plant Resources, 32(4), 325-337. DOI: 10.7732/kjpr.2019.32.4.325.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & the R Development Core Team. (2011) nlme: Linear and Nonlinear Mixed Effects Models. R Package Version 3.1-98. [Software]. Retrieved from http://web.mit.edu/~r/current/arch/i386\_linux26/lib/R/library/nlme/html/lme.html

R Development Core Team. (2012). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Rajakaruna, H., Drake, D. A. R., T. Chan, F., & Bailey, S. A. (2016). Optimizing performance of nonparametric species richness estimators under constrained sampling. *Ecology and evolution*, 6(20),7311-7322. DOI: 10.1002/ece3.2463

Ricotta, C., & Podani, J. (2017). On some properties of the Bray-Curtis dissimilarity and their ecological meaning. *Ecological complexity*, 31, 201-205.

Roberts, D.W. (2018, November 9). Ordination and Multivariate Analysis for Ecology [Software]. Retrieved from http://ecology.msu.montana.edu/labdsv/R

Salazar, L., Homeier, J., Kessler, M., Abrahamczyk, S., Lehnert, M., Kromer, T., & Kluge, J. (2015). Diversity patterns of ferns along elevational gradients in Andean tropical forests. *Plant Ecology & Diversity*, 8(1), 13-24. DOI: 10.1080/17550874.2013.843036

Shapiro, S. S., & Wilk, M.B. (1965). An analysis of variance test for normality (complete samples), *Biometrika*, 52, 591–611. DOI: 10.2307/2333709

Silva, V. L., Mehltreter, K., & Schmitt, J. L. (2018). Ferns as potential ecological indicators of edge effects in two types of Mexican forests. Ecological Indicators, 93, 669-676.

Srivastava, P., Singh, R., Tripathi, S., Singh, H., & Raghubanshi, A. (2018). Understanding the complex interaction between soil N availability and soil C dynamics under changing climate conditions. *Soil Management and Climate Change*, 20, 337–348. DOI: 10.1016/B978-0-12-812128-3.00022-7

Su, Z., Xiong, D., Dong, Y., Zhang, B., Zhang, S., Zheng, X., & Fang, H. (2015). Hydraulic properties of concentrated flow of a bank gully in the dry-hot valley region of southwest China. *Earth Surface Processes* 

and Landforms, 40(10), 1351-1363. DOI: 10.1002/esp.3724

Szewczyk, T. M., & McCain, C. M. (2019). Disentangling elevational richness: a multi-scale hierarchical Bayesian occupancy model of Colorado ant communities. *Ecography*, 42(5), 977-988. DOI: 10.1111/ecog.04115

Tan, B. C. (2013). The *Selaginella* Flora, A Good Indicator of the Philippine Pleistocene Island Groups. *Philippine Journal of Science*, 142: 211-213.

Torimaru, T., Akada, S., Ishida, K., Narita, M., & Higaki, D. (2018). Species habitat associations in an old-growth beech forest community organised by landslide disturbances. *Journal of Forest Research*, 23(2), 98-104. DOI: 10.1080/13416979.2017.1396418

Tuanmu, M. N., & Jetz, W. (2015). A global, remote sensing-based characterization of terrestrial habitat heterogeneity for biodiversity and ecosystem modelling. *Global Ecology and Biogeography*, 24(11),1329-1339. DOI: 10.1111/geb.12365

Wang, H. (2008). The Floristic composition and vertical distribution patterns of Pteridophytes in Mt. Fanjingshan Nature Reserve. Matser Dissertation, Guizhou University. (in Chinese with English abstract)

Wang, X., Zhao, L., Yan, B., Shi, L., Liu, G., & He, Y. (2016). Morphological and physiological responses of *Heteropogon contortus* to drought stress in a dry-hot valley. *Botanical Studies*, 57, 17.

Wu S. K. (1981). A study of the genus Aleuritopteris Fee in China. Journal of Systematics and Evolution, 19 (1), 57-74.

Wu, Z. Y. (2006). Flora Yunnanica (Volumes 20, 21). Science Press, Beijing.

Xu, Z. F., Tao, G. H., Yu, P. H., & Wang, Y. L. (1985). An approach to the vegetational changes from yuanjiang dry-hot valley of yunnan in the last 500 years. *Acta Botanica Yunnanica*, 7(4), 403-412. (in Chinese with English abstract)

Yang, F. C., Li, D. F., & Huang, H. P. (2018). *Pteridophytes of Thailand*. Nanhai Publishing House, Haikou.

Yang F. C. & Grote P. J. (2018). Riverine vegetation and environments of a Late Pleistocene river terrace, Khorat Plateau, Southeast Asia. *Palynology*, 42 (2), 158–167. DOI: 10.1080/01916122.2017.1296044

Yang F. C., Grote P. J. & Zhang S. T. (2019). The evolution of Mun River in Southeast Asia, and its relationship with the environmental changes in the late Middle Pleistocene, based on sedimentologic and palynological evidences. *Quaternary International*, 519, 50–57. DOI: 10.1016/j.quaint.2019.02.010

Yang, F. C., Zhang, C. L., Wu, G., Liang, S. Y., & Zhang, X. C. (2011). Endangered pteridophytes and their distribution in Hainan Island, China. *American Fern Journal*, 101 (2), 105-117. DOI: 10.1640/0002-8444-101.2.105

Zhang, B., Shi, K., Liu, C. Q., Ai, N. S., Liu, G. C., & Qin, F. C. (2009). Changes of Seasonal Precipitation of Yuanmou Dry-hot Valley in Recent 50 Years Based on method of DFA. *Scientia Geographica Sinica*, 4.

Zhang, L. L, Xu, C. D., & Feng, J. M. (2017). Biogeographical explanation of distribution pattern of fern diversity in Yunnan province. *Journal of Chuxiong Normal University*, 32(3), 45-50. (in Chinese with English abstract)