

GENERALIZATION OF HERMITE-HADAMARD-MERCER AND TRAPEZOID FORMULA TYPE INEQUALITIES INVOLVING BETA FUNCTION

MUHAMMAD AAMIR ALI AND ZHIYUE ZHANG

ABSTRACT. In this work, we prove a generalized version of Hermite-Hadamard-Mercer type inequalities using the Beta function. Moreover, we prove some new trapezoidal type inequalities involving Beta functions for differentiable convex functions. The main advantage of these inequalities is that these can be converted into similar classical integral inequalities, Riemann-Liouville fractional integral inequalities and k -Riemann-Liouville fractional integral inequalities. Finally, we give applications to special means of real numbers for newly established inequalities.

1. INTRODUCTION

In both pure and applied mathematics, the concept of convex functions is extremely important. Convex functions have a wide range of applications in fields such as finance, economics, and engineering.

Definition 1. [19] A function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$ is called convex for all $\varkappa, \gamma \in [\pi_1, \pi_2]$ and $\tau \in [0, 1]$ if

$$(1.1) \quad F(\gamma + \tau(\varkappa - \gamma)) \leq \tau F(\varkappa) + (1 - \tau) F(\gamma).$$

Function F is said to be a strictly convex if the inequality in (1.1) is strict for F , while F is said to be a concave function if $-F$ is convex.

Many important inequalities hold for convex functions, including Jensen, Jensen-Mercer, Hermite-Hadamard, and support line inequalities. The traditional Jensen's inequality is one of the most well-known inequalities, as stated below:

Theorem 1. [10, 11] For a convex function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:

$$(1.2) \quad F\left(\sum_{i=1}^n w_i \varkappa_i\right) \leq \sum_{i=1}^n w_i F(\varkappa_i),$$

where $\varkappa_i \in [\pi_1, \pi_2]$, $w_i \in [0, 1]$ and $\sum_{i=1}^{\infty} w_i = 1$.

In 2003, Mercer presented a variant of Jensen's inequality known as Jensen-Mercer inequality in [14].

Theorem 2. For a convex function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:

$$(1.3) \quad F\left(\pi_1 + \pi_2 - \sum_{i=1}^n w_i \varkappa_i\right) \leq F(\pi_1) + F(\pi_2) - \sum_{i=1}^n w_i F(\varkappa_i),$$

where $\varkappa_i \in [\pi_1, \pi_2]$, $w_i \in [0, 1]$ and $\sum_{i=1}^{\infty} w_i = 1$.

Due to its geometrical significance and applications, the Hermite-Hadamard inequality is one of the most studied inequality in the theory of convex functions. Because of the relevance of the Hermite-Hadamard inequality, there has been a significant amount of research committed to its expansions, generalizations, modifications, and applications. The Hermite-Hadamard inequality is shown in [7].

1991 *Mathematics Subject Classification.* 26D10, 26D15, 26A51.

Key words and phrases. Jensen-Mercer Inequality, Hermite-Hadamard-Mercer Inequality, Beta Function, Convex Function.

For a convex function $F : I \subset \mathbb{R} \rightarrow \mathbb{R}$ and $\pi_1, \pi_2 \in I$ with $\pi_1 < \pi_2$, we have

$$(1.4) \quad F\left(\frac{\pi_1 + \pi_2}{2}\right) \leq \frac{1}{\pi_2 - \pi_1} \int_{\pi_1}^{\pi_2} F(w) dw \leq \frac{F(\pi_1) + F(\pi_2)}{2}.$$

The inequality (1.4) holds in the reversed way if F is concave.

Recently, Kian and Moslehian [12] presented the following variant of Hermite-Hadamard inequality (1.4) using the inequality (1.3) and it is known as Hermite-Hadamard-Mercer inequality.

Theorem 3. *For a convex function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:*

$$(1.5) \quad \begin{aligned} F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) &\leq \frac{1}{\gamma - \varkappa} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} F(w) dw \\ &\leq \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \\ &\leq F(\pi_1) + F(\pi_2) - \frac{F(\varkappa) + F(\gamma)}{2} \end{aligned}$$

and

$$(1.6) \quad F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \leq F(\pi_1) + F(\pi_2) - \int_{\varkappa}^{\gamma} F(w) dw \leq F(\pi_1) + F(\pi_2) - F\left(\frac{\varkappa + \gamma}{2}\right),$$

where $\varkappa, \gamma \in [\pi_1, \pi_2]$.

Fractional integral operators have been used to expand the Hermite-Hadamard inequality. The Riemann-Liouville fractional operators are defined as:

Definition 2. [15, 18] *For an integrable function F on $[\pi_1, \pi_2]$, the left and right Riemann-Liouville fractional integrals are defined as:*

$$\begin{aligned} J_{\pi_1+}^{\alpha} F(\varkappa) &= \frac{1}{\Gamma(\alpha)} \int_{\pi_1}^{\varkappa} (\varkappa - w)^{\alpha-1} F(w) dw, \\ J_{\pi_2-}^{\alpha} F(\varkappa) &= \frac{1}{\Gamma(\alpha)} \int_{\varkappa}^{\pi_2} (w - \varkappa)^{\alpha-1} F(w) dw, \end{aligned}$$

where Γ represents the Gamma function and defined as $\Gamma(\alpha) = \int_0^{\infty} e^{-\tau} \tau^{\alpha-1} d\tau$.

Definition 3. [16] *For an integrable function F over $[\pi_1, \pi_2]$, the left and right k -Riemann-Liouville fractional integrals are defined as:*

$$\begin{aligned} J_{\pi_1+}^{\alpha, k} F(\varkappa) &= \frac{1}{k\Gamma_k(\alpha)} \int_{\pi_1}^{\varkappa} (\varkappa - w)^{\frac{\alpha}{k}-1} F(w) dw, \\ J_{\pi_2-}^{\alpha, k} F(\varkappa) &= \frac{1}{k\Gamma_k(\alpha)} \int_{\varkappa}^{\pi_2} (w - \varkappa)^{\frac{\alpha}{k}-1} F(w) dw, \end{aligned}$$

where $\Gamma_k(\alpha)$ is k -Gamma functions and defined as $\Gamma_k(\alpha) = \int_0^{\infty} e^{-\frac{\tau}{k}} \tau^{\alpha-1} d\tau$.

Recently, Sarikaya and Ata presented the generalized variant of Hermite-Hadamard inequality (1.4) using the notions of Beta functions.

Theorem 4. [22] *For a convex functions $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:*

$$(1.7) \quad \begin{aligned} F\left(\frac{\pi_1 + \pi_2}{2}\right) \beta(m, n) &\leq \frac{1}{2(\pi_2 - \pi_1)^{m+n-1}} \int_{\pi_1}^{\pi_2} \Xi(w) F(w) dw \\ &\leq \frac{F(\pi_1) + F(\pi_2)}{2} \beta(m, n), \end{aligned}$$

where $\beta(m, n) = \int_0^1 \tau^{m-1} (1 - \tau)^{n-1} d\tau$ is a Beta function and

$$\Xi(w) = (\pi_2 - w)^{m-1} (w - \pi_1)^{n-1} + (\pi_2 - w)^{n-1} (w - \pi_1)^{m-1}.$$

The fundamental benefit of inequality (1.7) is that it can be turned into classical Hermite-Hadamard inequality (1.4), fractional Hermite-Hadamard inequality [21] and k -fractional Hermite-Hadamard inequality [8] without having to prove each one separately.

Many studies have been conducted in the topic of integral inequality, particularly in Hermite-Hadamard inequality, using various integral operators. For example, the authors of [6] demonstrated various Hermite-Hadamard type inequalities for convex functions and applied them to numerical integration formulas. In [13] Kirmaci established Hermite-Hadamard type inequalities for differentiable convex functions. The authors of [21] used Riemann-Liouville fractional integrals to prove Hermite-Hadamard type inequalities for convex functions. The authors of [5] established Hermite-Hadamard type inequalities for (α, m) -logarithmically convex functions. Riemann-Liouville fractional integrals were used by Awan et al. [3] to refine fractional Hermite-Hadamard type inequalities for strongly convex functions. Using the generalized fractional integrals, Hermite-Hadamard type inequalities for convex functions were found in [20]. Using generalized fractional integrals, the authors showed some new Hermite-Hadamard type inequalities for harmonically convex functions in [25].

On the other hand, different authors have offered distinct versions of Hermite-Hadamard-Mercer type inequalities. For example, in [1], fractional Hermite-Hadamard-Mercer type inequalities are demonstrated using the Jensen-Mercer inequalities. Chu et al. [4] showed some new fractional estimates of the Hermite-Hadamard inequality for differentiable convex functions. Using the k -Caputo fractional derivative, the authors demonstrated Hermite-Hadamard type inequalities for convex functions in [26]. In [9], İşcan demonstrated a weighted variant of the Hermite-Hadamard-Mercer inequality. In [24], the authors showed Hermite-Hadamard-Mercer inequality for Harmonically convex functions.

Inspired by recent research, we establish a generalized version of Hermite-Hadamard-Mercer inequalities using the Beta function. The fundamental benefit of these inequalities is that these can be turned into classical Hermite-Hadamard-Mercer inequalities (1.6) and (1.5), fractional Hermite-Hadamard-Mercer inequalities [17] and k -fractional Hermite-Hadamard-Mercer inequalities [2, 23] without having to prove each one separately.

This article's structure is as follows: Section 2 proves the Hermite-Hadamard type inequality for convex functions using the Beta function. We prove some trapezoid formula type inequalities in Section 3. In Section 4, we present various applications to special means based on the recently demonstrated inequalities. Section 5 finishes with a few suggestions for future research.

2. HERMITE-HADAMARD-MERCER INEQUALITIES

In this section, we prove some new Hermite-Hadamard-Mercer type inequalities involving the Beta function.

Theorem 5. *For a convex function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:*

$$\begin{aligned}
 (2.1) \quad & F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \beta(m, n) \\
 & \leq [F(\pi_1) + F(\pi_2)] \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \Lambda(w) F(w) dw \\
 & \leq [F(\pi_1) + F(\pi_2)] \beta(m, n) - \beta(m, n) F\left(\frac{\varkappa + \gamma}{2}\right),
 \end{aligned}$$

where $\beta(m, n)$ is a Beta function,

$$\Lambda(w) = (\gamma - w)^{m-1} (w - \varkappa)^{n-1} + (\gamma - w)^{n-1} (w - \varkappa)^{m-1}$$

and $\varkappa, \gamma \in [\pi_1, \pi_2]$.

Proof. From Jensen-Mercer inequality and $u, v \in [\pi_1, \pi_2]$, we have

$$(2.2) \quad F\left(\pi_1 + \pi_2 - \frac{u+v}{2}\right) \leq F(\pi_1) + F(\pi_2) - \frac{F(u) + F(v)}{2}.$$

Now by setting $u = \tau\varkappa + (1 - \tau)\gamma$ and $v = \tau\gamma + (1 - \tau)\varkappa$ in (2.2), we have

$$(2.3) \quad F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \leq F(\pi_1) + F(\pi_2) - \frac{F(\tau\varkappa + (1 - \tau)\gamma) + F(\tau\gamma + (1 - \tau)\varkappa)}{2}.$$

Integrating to inequality (2.3) with respect to τ over $[0, 1]$ after multiplying $\tau^{m-1} (1 - \tau)^{n-1}$, we have

$$(2.4) \quad F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \beta(m, n) \leq \beta(m, n) [F(\pi_1) + F(\pi_2)] \\ - \frac{1}{2} \left[\int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau \varkappa + (1 - \tau) \gamma) d\tau \right. \\ \left. + \int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau \gamma + (1 - \tau) \varkappa) d\tau \right].$$

Thus, we obtain the first inequality in (2.1) by using the change of variables of integration.

We use the convexity of F to prove the second inequality in (2.1) and we obtain

$$(2.5) \quad F\left(\frac{\varkappa + \gamma}{2}\right) \leq \frac{F(\tau \varkappa + (1 - \tau) \gamma) + F(\tau \gamma + (1 - \tau) \varkappa)}{2}.$$

Integrating to inequality (2.5) with respect to τ over $[0, 1]$ after multiplying $\tau^{m-1} (1 - \tau)^{n-1}$, we have

$$(2.6) \quad F\left(\frac{\varkappa + \gamma}{2}\right) \beta(m, n) \leq \frac{1}{2} \left[\int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau \varkappa + (1 - \tau) \gamma) d\tau \right. \\ \left. + \int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau \gamma + (1 - \tau) \varkappa) d\tau \right].$$

By using change of variables of integration, we have the following inequality

$$F\left(\frac{\varkappa + \gamma}{2}\right) \beta(m, n) \leq \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \Lambda(w) F(w) dw$$

which implies that

$$(2.7) \quad -F\left(\frac{\varkappa + \gamma}{2}\right) \beta(m, n) \leq -\frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \Lambda(w) F(w) dw.$$

As a result, we may get the last inequality of (2.1) by adding $F(\pi_1) + F(\pi_2)$ on both sides of the inequality (2.7). The proof is finished. \square

Remark 1. In Theorem 5, if we set $m = n = 1$, then we recapture inequality (1.6).

Remark 2. In Theorem 5, if we set $m = 1$ and $n = \alpha$ or $m = \alpha$ and $n = 1$, then we recapture inequality (2.1) in [17, Theorem 2].

Theorem 6. For a convex function $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$, the following inequality holds:

$$(2.8) \quad F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \beta(m, n) \leq \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \\ \leq \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) \\ \leq \beta(m, n) \left[F(\pi_1) + F(\pi_2) - \frac{F(\varkappa) + F(\gamma)}{2} \right],$$

where $\beta(m, n)$ is a Beta function and

$$\Delta(w) = (\pi_1 + \pi_2 - \varkappa - w)^{m-1} (w - (\pi_1 + \pi_2 - \gamma))^{n-1} \\ + (\pi_1 + \pi_2 - \varkappa - w)^{n-1} (w - (\pi_1 + \pi_2 - \gamma))^{m-1}$$

and $\varkappa, \gamma \in [\pi_1, \pi_2]$.

Proof. From convexity of F and $u, v \in [\pi_1, \pi_2]$, we have

$$(2.9) \quad F\left(\pi_1 + \pi_2 - \frac{u + v}{2}\right) \leq \frac{F(\pi_1 + \pi_2 - u) + F(\pi_1 + \pi_2 - v)}{2}.$$

Now by setting $\pi_1 + \pi_2 - u = \tau(\pi_1 + \pi_2 - \varkappa) + (1 - \tau)(\pi_1 + \pi_2 - \gamma)$ and $\pi_1 + \pi_2 - v = \tau(\pi_1 + \pi_2 - \gamma) + (1 - \tau)(\pi_1 + \pi_2 - \varkappa)$ in inequality (2.9), we have

$$(2.10) \quad F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \leq \frac{1}{2} [F(\tau(\pi_1 + \pi_2 - \varkappa) + (1 - \tau)(\pi_1 + \pi_2 - \gamma)) + F(\tau(\pi_1 + \pi_2 - \gamma) + (1 - \tau)(\pi_1 + \pi_2 - \varkappa))].$$

Integrating to inequality (2.10) with respect to τ over $[0, 1]$ after multiplying $\tau^{m-1}(1 - \tau)^{n-1}$, we have

$$\begin{aligned} & F\left(\pi_1 + \pi_2 - \frac{\varkappa + \gamma}{2}\right) \beta(m, n) \\ & \leq \frac{1}{2} \left[\int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau(\pi_1 + \pi_2 - \varkappa) + (1 - \tau)(\pi_1 + \pi_2 - \gamma)) d\tau \right. \\ & \quad \left. + \int_0^1 \tau^{m-1} (1 - \tau)^{n-1} F(\tau(\pi_1 + \pi_2 - \gamma) + (1 - \tau)(\pi_1 + \pi_2 - \varkappa)) d\tau \right]. \end{aligned}$$

Thus, we obtain the first inequality in (2.8) by using the change of variables of integration.

We use the convexity of F to prove the second inequality in (2.8) and we obtain

$$(2.11) \quad \begin{aligned} & F(\tau(\pi_1 + \pi_2 - \varkappa) + (1 - \tau)(\pi_1 + \pi_2 - \gamma)) \\ & \leq \tau F(\pi_1 + \pi_2 - \varkappa) + (1 - \tau) F(\pi_1 + \pi_2 - \gamma) \end{aligned}$$

and

$$(2.12) \quad \begin{aligned} & F(\tau(\pi_1 + \pi_2 - \gamma) + (1 - \tau)(\pi_1 + \pi_2 - \varkappa)) \\ & \leq \tau F(\pi_1 + \pi_2 - \gamma) + (1 - \tau) F(\pi_1 + \pi_2 - \varkappa). \end{aligned}$$

By applying Jensen-Mercer inequality after addition of (2.11) and (2.12), we have

$$(2.13) \quad \begin{aligned} & F(\tau(\pi_1 + \pi_2 - \varkappa) + (1 - \tau)(\pi_1 + \pi_2 - \gamma)) + F(\tau(\pi_1 + \pi_2 - \gamma) + (1 - \tau)(\pi_1 + \pi_2 - \varkappa)) \\ & \leq F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa) \\ & \leq 2[F(\pi_1) + F(\pi_2)] - [F(\varkappa) + F(\gamma)]. \end{aligned}$$

Thus, we obtain second and third inequality of (2.8) by integrating to inequality (2.13) with respect to τ over $[0, 1]$ after multiplying $\tau^{m-1}(1 - \tau)^{n-1}$ and using the change of variables of integration. The proof is completed. \square

Remark 3. In Theorem 6, if we set $m = n = 1$, then we recapture inequality (1.5).

Remark 4. In Theorem 6, if we set $m = 1$ and $n = \alpha$ or $m = \alpha$ and $n = 1$, then we recapture inequality (2.2) in [17, Theorem 2].

Remark 5. In Theorem 6, if we set $m = 1$ and $n = \frac{\alpha}{k}$ or $m = \frac{\alpha}{k}$ and $n = 1$, then we obtain [2, Theorem 2.1].

Remark 6. In Theorem 5, if we set $\varkappa = \pi_1$ and $\gamma = \pi_2$, then we recapture inequality (1.7).

3. TRAPEZOIDAL TYPE INEQUALITIES

In this section, we prove some trapezoid formula type inequalities involving Beta functions.

Let's start with the following new Lemma.

Lemma 1. For a differentiable mapping $F : [\pi_1, \pi_2] \rightarrow \mathbb{R}$ over (π_1, π_2) and $F \in L[\pi_1, \pi_2]$, then the following equality holds:

$$(3.1) \quad \begin{aligned} & \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \\ & = \frac{\gamma - \varkappa}{2} \int_0^1 \beta_\tau(m, n) [F'(\pi_1 + \pi_2 - (\tau\varkappa + (1 - \tau)\gamma)) + F'(\pi_1 + \pi_2 - (\tau\gamma + (1 - \tau)\varkappa))] d\tau \\ & = \frac{\gamma - \varkappa}{2} \int_0^1 [\beta_\tau(m, n) - \beta_{1-\tau}(m, n)] F'(\pi_1 + \pi_2 - (\tau\varkappa + (1 - \tau)\gamma)) d\tau, \end{aligned}$$

where

$$\beta_{\tau}(m, n) = \int_0^{\tau} z^{m-1} (1-z)^{n-1} dz$$

and $\varkappa, \gamma \in [\pi_1, \pi_2]$.

Proof. From integration by parts, we have

$$\begin{aligned} (3.2) \quad & \int_0^1 \beta_{\tau}(m, n) F'(\pi_1 + \pi_2 - (\tau\varkappa + (1-\tau)\gamma)) d\tau \\ &= \beta_{\tau}(m, n) \frac{F(\pi_1 + \pi_2 - \varkappa)}{\gamma - \varkappa} \\ &\quad - \frac{1}{\gamma - \varkappa} \int_0^1 \tau^{m-1} (1-\tau)^{n-1} F(\tau(\pi_1 + \pi_2 - \varkappa) + (1-\tau)(\pi_1 + \pi_2 - \gamma)) d\tau \\ &= \beta_{\tau}(m, n) \frac{F(\pi_1 + \pi_2 - \varkappa)}{\gamma - \varkappa} \\ &\quad - \frac{1}{(\gamma - \varkappa)^{m+n}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} (\pi_1 + \pi_2 - \varkappa - w)^{n-1} (w - (\pi_1 + \pi_2 - \gamma))^{m-1} F(w) dw \end{aligned}$$

and

$$\begin{aligned} (3.3) \quad & \int_0^1 \beta_{\tau}(m, n) F'(\pi_1 + \pi_2 - (\tau\gamma + (1-\tau)\varkappa)) d\tau \\ &= \beta_{\tau}(m, n) \frac{F(\pi_1 + \pi_2 - \gamma)}{\gamma - \varkappa} \\ &\quad - \frac{1}{(\gamma - \varkappa)^{m+n}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} (\pi_1 + \pi_2 - \varkappa - w)^{m-1} (w - (\pi_1 + \pi_2 - \gamma))^{n-1} F(w) dw. \end{aligned}$$

Thus, we obtain the required equality by subtracting (3.2) from (3.3) and later multiplying $\frac{2}{(\gamma - \varkappa)}$. The proof is completed. \square

Remark 7. In Lemma 1, if we set $m = n = 1$, then Lemma 1 becomes [17, Corollary 1].

Remark 8. In Lemma 1, if we set $m = 1$ and $n = \alpha$ or $m = \alpha$ and $n = 1$, then we obtain [17, Lemma 1].

Remark 9. In Lemma 1, if we set $m = 1$ and $n = \frac{\alpha}{k}$ or $m = \frac{\alpha}{k}$ and $n = 1$, then we obtain equality (3.3) in [23, Remark 3.1].

Remark 10. In Lemma 1, if we set $\varkappa = \pi_1$ and $\gamma = \pi_2$, then we obtain [22, Lemma 5].

Theorem 7. We assume that the conditions of Lemma 1 hold. If $|F'|$ is a convex function, then we have the following trapezoid formula type inequality:

$$\begin{aligned} (3.4) \quad & \left| \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \right| \\ & \leq (\gamma - \varkappa) \left(|F'(\pi_1)| + |F'(\pi_2)| - \frac{|F'(\varkappa)| + |F'(\gamma)|}{2} \right) \int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_{\tau}(m, n)] d\tau. \end{aligned}$$

Proof. Taking modulus in (3.1) and using Jensen-Mercer inequality, we have

$$\begin{aligned} & \left| \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \right| \\ & \leq \frac{\gamma - \varkappa}{2} \int_0^1 |[\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)]| |F'(\pi_1 + \pi_2 - (\tau\varkappa + (1-\tau)\gamma))| d\tau \\ & \leq \frac{\gamma - \varkappa}{2} \left[\int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_{\tau}(m, n)] [|F'(\pi_1)| + |F'(\pi_2)| - \tau|F'(\varkappa)| - (1-\tau)|F'(\gamma)|] d\tau \right. \end{aligned}$$

$$\begin{aligned}
& + \int_{\frac{1}{2}}^1 [\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)] [|F'(\pi_1)| + |F'(\pi_2)| - \tau |F'(\varkappa)| - (1-\tau) |F'(\gamma)|] d\tau \\
& = (\gamma - \varkappa) \left(|F'(\pi_1)| + |F'(\pi_2)| - \frac{|F'(\varkappa)| + |F'(\gamma)|}{2} \right) \int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_{\tau}(m, n)] d\tau
\end{aligned}$$

and the proof is completed. \square

Remark 11. In Theorem 7, if we set $m = n = 1$, then Theorem 7 becomes [23, Corollary 3.5].

Remark 12. In Theorem 7, if we set $m = 1$ and $n = \alpha$ or $m = \alpha$ and $n = 1$, then we obtain [17, Theorem 4].

Remark 13. In Theorem 7, if we set $m = 1$ and $n = \frac{\alpha}{k}$ or $m = \frac{\alpha}{k}$ and $n = 1$, then we recapture inequality (3.11) in [23, Remark 3.7].

Remark 14. In Theorem 7, if we set $\varkappa = \pi_1$ and $\gamma = \pi_2$, then we obtain [22, Theorem 6].

Theorem 8. We assume that the conditions of Lemma 1 hold. If $|F'|^q$, $q > 1$ is convex functions, then we have the following trapezoidal type inequality:

$$\begin{aligned}
& \left| \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \right| \\
& \leq \frac{\gamma - \varkappa}{2} \left(\int_0^1 |[\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)]|^p d\tau \right)^{\frac{1}{p}} \left(|F'(\pi_1)|^q + |F'(\pi_2)|^q - \frac{|F'(\varkappa)|^q + |F'(\gamma)|^q}{2} \right)^{\frac{1}{q}},
\end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. Taking modulus in equality (3.1) and applying Hölder's inequality, we have

$$\begin{aligned}
& \left| \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \right| \\
& \leq \frac{\gamma - \varkappa}{2} \left(\int_0^1 |[\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)]|^p d\tau \right)^{\frac{1}{p}} \\
& \quad \times \left(\int_0^1 |F'(\pi_1 + \pi_2 - (\tau\varkappa + (1-\tau)\gamma))|^q d\tau \right)^{\frac{1}{q}}.
\end{aligned}$$

From Jensen-Mercer inequality, we have

$$\begin{aligned}
& \left| \frac{F(\pi_1 + \pi_2 - \gamma) + F(\pi_1 + \pi_2 - \varkappa)}{2} \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) F(w) dw \right| \\
& \leq \frac{\gamma - \varkappa}{2} \left(\int_0^1 |[\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)]|^p d\tau \right)^{\frac{1}{p}} \\
& \quad \times \left(\int_0^1 [|F'(\pi_1)|^q + |F'(\pi_2)|^q - \tau |F'(\varkappa)|^q - (1-\tau) |F'(\gamma)|^q] d\tau \right)^{\frac{1}{q}} \\
& = \frac{\gamma - \varkappa}{2} \left(\int_0^1 |[\beta_{\tau}(m, n) - \beta_{1-\tau}(m, n)]|^p d\tau \right)^{\frac{1}{p}} \left(|F'(\pi_1)|^q + |F'(\pi_2)|^q - \frac{|F'(\varkappa)|^q + |F'(\gamma)|^q}{2} \right)^{\frac{1}{q}}
\end{aligned}$$

and the proof is completed. \square

Remark 15. In Theorem 8, if we set $m = n = 1$, then Theorem 8 becomes [23, Corollary 3.8].

Remark 16. In Theorem 8, if we set $m = 1$ and $n = \alpha$ or $m = \alpha$ and $n = 1$, then we obtain [23, Corollary 3.9].

Remark 17. In Theorem 8, if we set $m = 1$ and $n = \frac{\alpha}{k}$ or $m = \frac{\alpha}{k}$ and $n = 1$, then we recapture inequality (3.11) in [23, Corollary 3.10].

Remark 18. In Theorem 8, if we set $\varkappa = \pi_1$ and $\gamma = \pi_2$, then we obtain [22, Theorem 7].

4. APPLICATION TO SPECIAL MEANS

In this section, we give applications of newly established inequalities in the context of special means of real numbers. For arbitrary positive real numbers π_1 and π_2 ($\pi_1 \neq \pi_2$), we consider the means as follows:

(1) The arithmetic mean

$$\mathcal{A} = \mathcal{A}(\pi_1, \pi_2) = \frac{\pi_1 + \pi_2}{2}.$$

(2) The harmonic mean

$$\mathcal{H} = \mathcal{H}(\pi_1, \pi_2) = \frac{2\pi_1\pi_2}{\pi_1 + \pi_2}.$$

(3) The geometric mean

$$\mathcal{G} = \mathcal{G}(\pi_1, \pi_2) = \sqrt{\pi_1\pi_2}.$$

Proposition 1. For $0 < \pi_1 < \pi_2$ and $p \in \mathbb{N}$, $p \geq 2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} (2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma))^p \beta(m, n) &\leq 2\mathcal{A}(\pi_1^p, \pi_2^p) \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \Lambda(w) w^p dw \\ &\leq (2\mathcal{A}(\pi_1^p, \pi_2^p) - \mathcal{A}^p(\varkappa, \gamma)) \beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 5 to the convex function $F(w) = w^p$, $w > 0$. \square

Proposition 2. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} (2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma))^{-1} \beta(m, n) &\leq 2\mathcal{H}^{-1}(\pi_1, \pi_2) \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \frac{\Lambda(w)}{w} dw \\ &\leq (2\mathcal{H}^{-1}(\pi_1, \pi_2) - \mathcal{A}^{-1}(\varkappa, \gamma)) \beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 5 to the convex function $F(w) = \frac{1}{w}$, $w \neq 0$. \square

Proposition 3. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} \ln(2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma)) \beta(m, n) &\leq 2\ln(\mathcal{G}(\pi_1, \pi_2)) \beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\varkappa}^{\gamma} \Lambda(w) \ln w dw \\ &\leq (2\ln(\mathcal{G}(\pi_1, \pi_2)) - \ln(\mathcal{A}(\varkappa, \gamma))) \beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 5 to the convex function $F(w) = \ln w$. \square

Proposition 4. For $0 < \pi_1 < \pi_2$ and $p \in \mathbb{N}$, $p \geq 2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} (2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma))^p \beta(m, n) &\leq \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \Delta(w) w^p dw \\ &\leq \mathcal{A}((\pi_1 + \pi_2 - \varkappa)^p, (\pi_1 + \pi_2 - \gamma)^p) \beta(m, n) \\ &\leq (2\mathcal{A}(\pi_1^p, \pi_2^p) - \mathcal{A}(\varkappa^p, \gamma^p)) \beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 6 to the convex function $F(w) = w^p$, $w > 0$. \square

Proposition 5. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} (2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma))^{-1} \beta(m, n) &\leq \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1 + \pi_2 - \gamma}^{\pi_1 + \pi_2 - \varkappa} \frac{\Delta(w)}{w} dw \\ &\leq \mathcal{H}^{-1}(\pi_1 + \pi_2 - \varkappa, \pi_1 + \pi_2 - \gamma) \beta(m, n) \\ &\leq (2\mathcal{H}^{-1}(\pi_1, \pi_2) - \mathcal{H}^{-1}(\varkappa, \gamma)) \beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 6 to the convex function $F(w) = \frac{1}{w}$, $w \neq 0$. \square

Proposition 6. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} \ln(2\mathcal{A}(\pi_1, \pi_2) - \mathcal{A}(\varkappa, \gamma))\beta(m, n) &\leq \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1+\pi_2-\gamma}^{\pi_1+\pi_2-\varkappa} \Delta(w) \ln w dw \\ &\leq \ln(\mathcal{G}(\pi_1 + \pi_2 - \varkappa, \pi_1 + \pi_2 - \gamma))\beta(m, n) \\ &\leq (2\ln(\mathcal{G}(\pi_1, \pi_2)) - \ln(\mathcal{G}(\varkappa, \gamma)))\beta(m, n). \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 6 to the convex function $F(w) = \ln w$. \square

Proposition 7. For $0 < \pi_1 < \pi_2$ and $p \in \mathbb{N}$, $p \geq 2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} &\left| \mathcal{A}((\pi_1 + \pi_2 - \varkappa)^p, (\pi_1 + \pi_2 - \gamma)^p)\beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1+\pi_2-\gamma}^{\pi_1+\pi_2-\varkappa} \Delta(w) w^p dw \right| \\ &\leq p(\gamma - \varkappa) \left(2\mathcal{A}(\pi_1^{p-1}, \pi_2^{p-1}) - \mathcal{A}(\varkappa^{p-1}, \gamma^{p-1}) \right) \int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_\tau(m, n)] d\tau. \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 7 to the convex function $F(w) = w^p$, $w > 0$. \square

Proposition 8. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} &\left| \mathcal{H}^{-1}(\pi_1 + \pi_2 - \varkappa, \pi_1 + \pi_2 - \gamma)\beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1+\pi_2-\gamma}^{\pi_1+\pi_2-\varkappa} \frac{\Delta(w)}{w} dw \right| \\ &\leq (\varkappa - \gamma) (2\mathcal{H}^{-1}(\pi_1^2, \pi_2^2) - \mathcal{H}^{-1}(\varkappa^2, \gamma^2)) \int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_\tau(m, n)] d\tau. \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 7 to the convex function $F(w) = \frac{1}{w}$, $w \neq 0$. \square

Proposition 9. For $0 < \pi_1 < \pi_2$, the following inequality holds for all $\varkappa, \gamma \in [\pi_1, \pi_2]$:

$$\begin{aligned} &\left| \ln(\mathcal{G}(\pi_1 + \pi_2 - \varkappa, \pi_1 + \pi_2 - \gamma))\beta(m, n) - \frac{1}{2(\gamma - \varkappa)^{m+n-1}} \int_{\pi_1+\pi_2-\gamma}^{\pi_1+\pi_2-\varkappa} \Delta(w) \ln w dw \right| \\ &\leq (\gamma - \varkappa) (2\mathcal{H}^{-1}(\pi_1, \pi_2) - \mathcal{H}^{-1}(\varkappa, \gamma)) \int_0^{\frac{1}{2}} [\beta_{1-\tau}(m, n) - \beta_\tau(m, n)] d\tau. \end{aligned}$$

Proof. The needed inequality can be obtained by applying Theorem 7 to the convex function $F(w) = \ln w$. \square

5. CONCLUSIONS

We used the Beta function in this paper to prove several new generalizations of Hermite-Hadamard-Mercer type inequalities. We also proved some new trapezoidal type inequalities involving the Beta function for differentiable convex functions. Finally, we demonstrated the application of special means of real numbers for newly established inequalities. Researchers can obtain comparable inequalities for different types of convexity and special functions in their future work, which is a new and interesting problem.

AUTHOR CONTRIBUTIONS

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

FUNDING

The work was supported by the Natural Science Foundation of China (Grant No. 11971241).

AVAILABILITY OF DATA MATERIALS

Data sharing not applicable to this paper as no data sets were generated or analyzed during the current study.

ACKNOWLEDGMENTS

The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

CONFLICTS OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

- [1] T. Abdeljawad, M. A. Ali, P. O. Mohammed and Arton Kashuri, On inequalities of Hermite-Hadamard-Mercer type involving Riemann-Liouville fractional integrals. *AIMS Mathematics*, **5** (2020), 7316–7331.
- [2] M. A. Ali, T. Hussain, M. Z. Iqbal and F. Ejaz, Inequalities of Hermite-Hadamard-Mercer type for convex functions via K -fractional integrals. *International Journal of Mathematical Modelling & Computations*, **10** (2020), 227-238.
- [3] M. U. Awan, M. A. Noor, T.-S. Du and K. I. Noor, New refinements of fractional Hermite-Hadamard inequality. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas*. **113** (2019), 21-29.
- [4] H.-H. Chu, S. Rashid, Z. Hammouch and Y.-M. Chu, New fractional estimates for Hermite-Hadamard-Mercer's type inequalities. *Alexandria Engineering Journal*, **59** (2020), 3079–3089.
- [5] J. Deng and W. R. Wang, Fractional Hermite-Hadamard inequalities for (α, m) -logarithmically convex functions. *Journal of Inequalities and Applications*, **2013** (2013), 1-11.
- [6] S. S. Dragomir, R. P. Agrwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula. *Applied mathematics letters*, **11** (1998), 91-95.
- [7] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann. *Journal de Mathématiques Pures et Appliquées*, **58** (1893), 171-215.
- [8] R. Hussain, A. Ali, G. Gulshan, A. Latif and M. Muddassar, Generalized co-ordinated integral inequalities for convex functions by way of k -fractional derivatives, Miskolc Mathematical Notes Publications of the university of Miskolc. (Submitted).
- [9] İ. İşcan, Weighted Hermite-Hadamard-Mercer type inequalities for convex functions. *Numerical Methods for Partial Differential Equations*, **37** (2021), 118–130.
- [10] M. A. Khan, J. Khan and J. Pečarić, Generalization of Jensen's and Jensen-Steffensen's inequalities by generalized majorization theorem. *Journal of Mathematical Inequalities*, **11** (2017), 1049-1074.
- [11] J. Khan, M. A. Khan and J. Pecaric, On Jensen's type inequalities via generalized majorization inequalities, *Filomat*, **32** (2018), 5719-5733.
- [12] M. Kian and M.S. Moslehian, Refinements of the operator Jensen-Mercer inequality. *Electron. J. Linear Algebra*, **26** (2013), 742-753.
- [13] U. S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula. *Applied mathematics and computation*, **147** (2004), 137-146.
- [14] A. Mercer, A variant of Jensen's inequality. *Journal of Inequalities in Pure and Applied Mathematics*, **4** (2003), 73.
- [15] K. S. Miller and B. Ross, *An introduction to the fractional calculus and fractional differential equations*, Wiley, New York, 1993.
- [16] S. Mubeen and G. M. Habibullah, k -Fractional integrals and applications. *Int. J. Contemp. Math. Sci.*, **7** (2012), 89–94.
- [17] H. ÖĞÜLMÜŞ and M. Z. Sarikaya, Hermite-Hadamard-Mercer type inequalities for fractional integrals. ResearchGate Article, <https://www.researchgate.net/publication/337682540>.
- [18] K. B. Oldham and J. Spanier, *The fractional calculus*, Academic Press, San Diego, 1974.
- [19] J. Pečarić, F. Prochan, and Y. L. Tong, *Convex Functions, Partial Ordering and Statistical Applications*, Academic Press, New York, NY, USA, 1991.
- [20] M. Z. Sarikaya and F. Ertugral, On the generalized Hermite-Hadamard inequalities. *Annals of the University of Craiova-Mathematics and Computer Science Series*, **47** (2020), 193-213.
- [21] M. Z. Sarikaya, E. Set, H. Yaldiz and N. Basak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities. *Mathematical and Computer Modelling*, **57** (2013) 2403-2407.
- [22] M. Z. Sarikaya and F. Ata, On the generalized Hermite-Hadamard inequalities involving Beta function. ResearchGate Article, <https://www.researchgate.net/publication/337103933>.
- [23] M. Vivas-Cortez, M. A. Ali, A. Kashuri and H. Budak, Generalizations of fractional Hermite-Hadamard-Mercer like inequalities for convex functions. *AIMS Mathematics*, **6** (2021), 9397–9421.
- [24] X. You, M. A. Ali, H. Budak, J. Reunsumrit and T. Sitthiwirattam, Hermite-Hadamard-Mercer-Type Inequalities for Harmonically Convex Mappings. *Mathematics*, **9** (2021), 2556.
- [25] D. Zhao, M. A. Ali, A. Kashuri, and H. Budak, Generalized fractional integral inequalities of Hermite-Hadamard type for harmonically convex functions. *Adv. Differ. Equ.*, **2020** (2020), 1-37.

- [26] S. Zhao, S.I. Butt, W. Nazeer, J. Nasir, M. Umar and Y. Liu, Some Hermite-Jensen-Mercer type inequalities for k -Caputo-fractional derivatives and related results. *Advances in Difference Equations*, **2020** (2020), 1-17.

JIANGSU KEY LABORATORY FOR NSLSCS, SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING, CHINA.

E-mail address: mahr.muhammad.aamir@gmail.com

JIANGSU KEY LABORATORY FOR NSLSCS, SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING, CHINA.

E-mail address: 05298@njnu.edu.cn