

**Table 1:** Baseline characteristics of all patients, survivors, and non-survivors prior to the initiation of ECPR

	Overall	Survivor	Non-Survivor	p
N	47	12	35	-
Age in years (median, [IQR])	53.00 [45.00, 64.00]	44.00 [32.25, 55.75]	60.00 [46.00, 65.65]	0.034*
Male (%)	33 (70.2)	10 (83.3)	23 (65.7)	0.302
<b>Race and Ethnicity</b>				
White (%)	35 (74.5)	9 (75.0)	26 (74.3)	1.000
Black (%)	7 (14.9)	2 (16.7)	5 (14.3)	1.000
Unknown (%)	4 (8.5)	1 (8.3)	3 (8.6)	1.000
Asian (%)	1 (2.1)	0 (0.0)	1 (2.9)	1.000
Hispanic or Latino (%)	5 (10.6)	0 (0.0)	5 (14.3)	0.309
<b>Past Medical History</b>				
Coronary Artery Disease (%)	27 (57.4)	6 (50.0)	21 (60.0)	0.737
Cardiomyopathy (%)	20 (42.6)	3 (25.0)	17 (48.6)	0.191
Prior Revascularization (%)	18 (38.3)	5 (41.7)	13 (37.1)	1.000
Ventricular Arrhythmia (%)	4 (8.5)	1 (8.3)	3 (8.6)	1.000
Peripheral Arterial Disease (%)	4 (8.5)	1 (8.3)	3 (8.6)	1.000
Stroke (%)	3 (6.4)	0 (0.0)	3 (8.6)	0.560
Dialysis (%)	3 (6.4)	1 (8.3)	2 (5.7)	1.000