

Table.3. Comparisons of complications and outcomes between LDA and control groups

	LDA (N=235)	Control (N=235)	OR (95% CI)	P value
Hypertensive disorders	26(11.1)	40(17.0)	0.65(0.41-1.03)	0.084
Gestational hypertension	15(6.38)	17(7.23)	0.88(0.45-1.73)	0.855
PE	11(4.68)	23(9.79)	0.48(0.24-0.95)	0.048
DC	9	17	--	--
MC	2	6	--	--
Neonatal weight(mean)	2550(2280-2780)	2415(2120-2650)	--	< 0.001
SGA	55(23.40)	74(31.49)	0.74(0.55-1.00)	0.063
DC	31	37	--	--
MC	24	37	--	--
sSGA ^{#1}	45/225(20.0)	62/223(27.80)	0.72(0.51-1.01)	0.060
Gestational week(d)	259.0(252.0-263.0)	254.0(243.0-260.0)	--	< 0.001
Preterm birth^{#2}	18(7.66)	36(15.32)	0.50(0.29-0.86)	0.013
Spontaneous	15(6.38)	32(13.62)	0.47(0.26-0.84)	0.013
DC	6	13	--	--
MC	9	19	--	--
Iatrogenic	3(1.28)	4(1.70)	0.75(0.17-3.31)	1.000
Bleeding amount	505.66±300.96	524.57±445.61	--	0.590
Hemorrhage	8(3.40)	9(3.83)	0.89(0.35-2.26)	1.000

#1: Excluded both of twins were growth restricted (10 cases in the LDA group, 12 cases in the Control group).

#2: Gestational weeks less than 34.