

Table 4: Warthog and bushpig abundance in the four vegetation types in NGR, inferred from camera traps. The total number of wild suid species and events are shown. Mean number of wild suids/events indicate the number of observations per camera day. The numbers in bold indicate warthogs results while the italicised ones are for bushpigs.  $P$ = statistical significance of a 1-way ANOVA on the log-transformed number of images with wild suids, comparing different habitats. Since the total number of camera days is relatively small, differences among habitats are not significant.

<i>Vegetation types</i>	Number of cameras	Camera days	Number of wild suids		Events		Mean number of wild suids		S.E.M (wild suids)		Mean number of events		S.E.M (events)		P	
<i>Sand (S)</i>	3	97	<b>52</b>	<i>7</i>	<b>26</b>	<i>2</i>	<b>0.536</b>	<i>0.072</i>	<b>0.418</b>	<i>2.330</i>	<b>0.268</b>	<i>0.021</i>	<b>0.241</b>	<i>0.660</i>		
<i>Bushveld (B)</i>	5	170	<b>4</b>	<i>6</i>	<b>3</b>	<i>5</i>	<b>0.024</b>	<i>0.035</i>	<b>0.015</b>	<i>0.063</i>	<b>0.018</b>	<i>0.029</i>	<b>0.011</b>	<i>0.064</i>		
<i>Combined S+B</i>	8	267	<b>56</b>	<i>13</i>	<b>29</b>	<i>7</i>	<b>0.210</b>	<i>0.049</i>	<b>0.179</b>	<i>0.869</i>	<b>0.109</b>	<i>0.026</i>	<b>0.090</b>	<i>0.246</i>	<b>0.903</b>	<i>0.634</i>
<i>Thicket</i>	17	514	<b>77</b>	<i>174</i>	<b>20</b>	<i>27</i>	<b>0.150</b>	<i>0.338</i>	<b>0.249</b>	<i>1.007</i>	<b>0.039</b>	<i>0.052</i>	<b>0.069</b>	<i>0.087</i>		
<i>Water</i>	23	602	<b>168</b>	<i>307</i>	<b>36</b>	<i>51</i>	<b>0.279</b>	<i>0.515</i>	<b>0.249</b>	<i>1.323</i>	<b>0.060</b>	<i>0.086</i>	<b>0.069</b>	<i>0.182</i>		
<i>Overall</i>	48	1383	<b>301</b>	<i>491</i>	<b>85</b>	<i>85</i>	<b>0.218</b>	<i>0.355</i>	<b>n/a</b>	<i>n/a</i>	<b>0.061</b>	<i>0.061</i>	<b>n/a</b>	<i>n/a</i>		