

**Table 1. Baseline characteristics of the study population**

Characteristic	Value
Age (years)	60.6 ± 9.4
Male sex (%)	77 (75)
Paroxysmal AF (n°)	102 (100)
Diagnosis-to-ablation (months)	51.5 ± 46.9
LA diameter (mm)	39.8 ± 5.2
LA area (cmq)	21.0 ± 6.0
LV EF (%)	60.4 ± 6.2
BMI (kg/m <sup>2</sup> )	25.8 ± 3.8
AHT (n°)	69 (68)
DM (n°)	3 (3)
CAD (n°)	8 (8)
CHA <sub>2</sub> DS <sub>2</sub> vasc score (0-9)	1.5 ± 1.0

Table 1: Baseline characteristics of the study population (AF, atrial fibrillation; LA, left atrium; LV, left ventricle; EF, ejection fraction; BMI, body mass index; AHT, arterial hypertension; DM, diabetes mellitus; CAD, coronary artery disease).

Categorical variables are expressed as absolute and percentage (in brackets).

Continuous variables are expressed as mean ± SD.

**Table 2. Procedural characteristics**

Characteristic	Value
Procedural time (minutes)	79.9 ± 20.3
Fluoroscopic time (minutes)	17.1 ± 6.6
LSPV freeze >1 (n°)	29 (28)
LIPV freeze >1 (n°)	15 (15)
RIPV freeze >1 (n°)	18 (18)
RSPV freeze >1 (n°)	8 (8)
Total cryo-applications time (seconds)	966 ± 239.1
Nadir LSPV (°C)	-48.2 ± 5.0
Nadir LIPV (°C)	-44.3 ± 4.5
Nadir RIPV (°C)	-47.3 ± 6.2
Nadir RSPV (°C)	-50.9 ± 5.6

Table 2: Procedural characteristics (LSPV, left superior pulmonary vein; LIPV, left inferior pulmonary vein; RIPV, right inferior pulmonary vein; RSPV, right superior pulmonary vein). Categorical variables are expressed as absolute and percentage (in brackets). Continuous variables are expressed as mean ± SD.