

	Women requiring no respiratory support N=54	Women requiring respiratory support N=15	p-value
SYMPTOMS n (%)			
Asymptomatic	4/54 (7)	0/15(0)	ns
Cough	30/54 (56)	13/15 (87)	<0.05
Fever	33/54 (57)	14/15 (93)	<0.05
Shortness of Breath	10/54 (19)	10/15 (67)	<0.001
Other	30/54 (56)	9/15 (60)	ns
Laboratory Findings mean ± SEM			
White cell count (cellsx10 ⁹ /L)	9.2±0.6	7.6 ± 0.5	ns
Neutrophils (cellsx10 ⁹ /L)	7.0 ± 0.6	6.2 ± 0.4	ns
Lymphocytes (cellsx10 ⁹ /L)	1.4 ± 0.1	0.9 ± 0.1	<0.01
Platelets (cellsx10 ⁹ /L)	2144± 10	232 ± 14	ns
C-reactive protein (mg/L)	46.4 ± 6.9	56.3 ± 12.0	ns
Alanine Transaminase (iU/L)	47.3 ± 13.5	28.5 ± 5.7	ns
Alkaline Phosphatase (iU/L)	148 ± 16	131 ± 16	ns
Creatinine (micromole/L)	50.6 ± 1.7	49.5 ± 2.7	ns
Potassium (mmol/L)	4.0 ± 0.06	3.8 ± 0.08	<0.05
Radiography			
Chest radiography performed n (%)	17/54 (31)	15/15 (100)	<0.001
Consolidation on chest xray n (%)	10/17 (59)	13/15 (87)	ns

Table 3: Comparison of clinical and radiographic characteristics of pregnant women infected with confirmed SARS-CoV2 who did not or did required respiratory support across a maternal medicine network in London, UK, 1st March – 1st July 2020.