

**Table 2.** Demographic and clinical characteristics of the cases according to the groups with and without cranial nerve involvement

	No cranial nerve involvement (n = 167)	Cranial nerve involvement present (n = 135)	p-value
<b>Age</b>	40.2 ± 16.7	39.3 ± 16.4	0.624†
<b>Gender</b>			0.138‡
Male	102 (61.1%)	71 (52.6%)	
Female	65 (38.9%)	64 (47.4%)	
<b>Body mass index (kg/m<sup>2</sup>)</b>	26.3 ± 4.9	26.8 ± 5.5	0.348†
<25.00 kg/m <sup>2</sup>	68 (40.7%)	55 (40.7%)	
25.00-29.99 kg/m <sup>2</sup>	70 (41.9%)	49 (36.3%)	
≥30.00 kg/m <sup>2</sup>	29 (17.4%)	31 (23.0%)	
<b>Comorbidities</b>	34 (20.4%)	28 (20.7%)	0.935‡
Hypertension	16 (9.6%)	13 (9.6%)	>0.999¶¶
Diabetes mellitus	9 (5.4%)	8 (5.9%)	>0.999¶¶
Cardiac diseases	11 (6.6%)	5 (3.7%)	0.393¶¶
Asthma – COPD	10 (6.0%)	10 (7.4%)	0.795¶¶
Behçet's disease	0 (0.0%)	1 (0.7%)	N/A
<b>Diagnostic method</b>			<0.001‡
Screening	92 (55.1%)	19 (14.1%)	
Patient complaints	75 (44.9%)	116 (85.9%)	
<b>Intensive care requirement</b>	27 (16.2%)	28 (20.7%)	0.382¶¶
<b>Length of ICU stay (days)</b>	11 (5-32)	10.5 (3-18)	0.548¥
<b>Length of hospital stay (days)</b>	6 (3-37)	7 (3-23)	0.069¥

COPD, Chronic obstructive pulmonary disease; ICU, intensive care unit; † Student's t test, ‡ Pearson's  $\chi^2$  test, ¶¶ Continuity-corrected  $\chi^2$  test, ¥ Mann-Whitney U test, N/A: Not applicable