

TABLE 3 Determining factors associated with incidence of complications (all complications grouped):

Dependent: Complication		No	yes	OR (univariate)	OR (multivariate)
Sex	F	8 (80.0)	2 (20.0)	-	
	M	56 (73.7)	20 (26.3)	1.43 (0.32-10.00, p=0.668)	
Age at the operation	≤24 months	19 (65.5)	10 (34.5)	-	
	>24 months	45 (78.9)	12 (21.1)	0.51 (0.19-1.39, p=0.181)	
Initial Size	≤5cm	32 (72.7)	12 (27.3)	-	
	> 5cm	32 (76.2)	10 (23.8)	0.83 (0.31-2.20, p=0.713)	
Localization	Bladder	22 (73.3)	8 (26.7)	-	
	Bladder_Prostate	42 (75.0)	14 (25.0)	0.92 (0.34-2.60, p=0.866)	
Bladder neck-trigone dissection	no	44 (80.0)	11 (20.0)	-	
	yes	20 (64.5)	11 (35.5)	2.20 (0.82-5.99, p=0.118)	
Posterior bladder wall dissection	no	60 (78.9)	16 (21.1)	-	5.15 (1.29-22.79, p=0.022)
	yes	4 (40.0)	6 (60.0)	5.62 (1.44-24.37, p=0.014)	
Anterior partial cystectomy	no	36 (72.0)	14 (28.0)	-	
	yes	28 (77.8)	8 (22.2)	0.73 (0.26-1.97, p=0.545)	
Partial prostatectomy	no	41 (78.8)	11 (21.2)	-	
	yes	23 (67.6)	11 (32.4)	1.78 (0.67-4.80, p=0.247)	
Ureteral reimplantation	no	22 (71.0)	9 (29.0)	-	
	yes	42 (76.4)	13 (23.6)	0.76 (0.28-2.09, p=0.582)	
Only Mucosectomy	yes	4 (80.0)	1 (20.0)	-	
	no	60 (74.1)	21 (25.9)	1.40 (0.19-28.21, p=0.769)	
Brachytherapy	LDR	47 (70.1)	20 (29.9)	-	0.31 (0.05-1.26, p=0.147)
	PDR	17 (89.5)	2 (10.5)	0.28 (0.04-1.09, p=0.105)	