

**Table 2: Perioperative and postoperative outcomes of 120 patients stratified according to antithrombotic therapy**

	All Patients Median (IQR) N=120	Group 1 Median (IQR) N=60	Group 2 Median (IQR) N=60	p value
<i>Perioperative</i>				
Specimen weight (g)	42 (10-152)	44 (10-134)	40 (10-152)	0.765
Enucleation time (min)	40 (17-90)	38 (17-90)	40 (20-75)	0.491
Enucleation efficiency (g/min)	1.14 (0.33-2.8)	1.15 (0.33-2.2)	1.11 (0.4-2.8)	0.775
Morcellation time (min)	10 (3-44)	10 (3-37)	10 (3-44)	0.748
Morcellation efficiency (g/min)	4.45 (1.1-10)	4.45 (2.2-8)	4.53 (1.1-10)	0.579
<i>Postoperative</i>				
Catheterization (days)	2 (2-3)	2 (2-3)	2 (2-3)	0.649
Hospital stay (days)	3 (3-4)	3 (3-4)	3 (3-4)	0.649
Hemoglobin decrease (g/dL)	1.3 (0.1-4)	1.3 (0.1-4)	1.4 (0.4-3.1)	0.736

*IQR: Inter Quantile Range.*