

	KT group Mean±SD (n=25)	Sham KT group Mean±SD (n=25)	P-value
Age (years)	25.60±5.90	24.71±5.95	0.557
Parity	1.2±1.60	1.2±1.49	0.568
Gravidas	2.24±1.69	2.42±1.44	0.324
Gestation, week	24.14±3.82	23.22±4.12	0.620
Body mass index (kg/m ²)	27.32±2.10	26.45±2.45	0.812

Table 1. Comparison of clinical characteristics of pregnant women with sacroiliac dysfunction

KT: Kinesio taping; SD: Standart deviation

Table 2. Assessment of VAS, RMDQ and PGQ values before and after treatment between and within groups

		KT group (n=25)		Sham KT group (n=25)		P-value (inter group)
		Mean ± SD	P-value (intra-group)	Mean ± SD	P-value (intra-group)	
VAS rest						
	Basal	6.55±1.50		7.11±1.31		0.432
	5 th week	2.80±1.67	0.013	6.73±1.19	0.08	<0.001
VAS motion						
	Basal	7.41±1.22		7.83±0.90		0.352
	5 th week	1.42±1.05	<0.001	6.32±0.89	0.085	<0.001
RMDQ						
	Basal	17.73±3.68		18.07±3.63		0.451
	5 th week	4.50±3.71	<0.001	16.52±3.59	0.064	<0.001
PGQ Activity Subscale						
	Basal	42.10±8.22		41.71±8.18		0.464
	5 th week	24.42±8.57	<0.001	39.58±8.55	0.058	<0.001
PGQ Symptom Subscale						
	Basal	10.23±2.48		11.08±2.40		0.482
	5 th week	3.44±2.24	<0.001	9.32±2.38	0.056	<0.001

KT: Kinesiotaping; VAS: Visual analog scale; RMDQ: Roland-Morris Disability Questionnaire; PGQ: Pelvic Girdle Questionnaire SD: Standard deviation; Samples t-test; Paired samples t-test