

Using Community Science to Better Understand Lead Exposure Risks

Matthew Dietrich^{1*}, John Shukle¹, Mark P.S. Krekeler,^{2,3} Gabriel Filippelli^{1,4}

¹Department of Earth Sciences, Indiana University – Purdue University Indianapolis, Indianapolis, IN, USA

²Department of Geology & Environmental Earth Science, Miami University, Oxford, OH 45056, USA

³Department of Mathematical and Physical Sciences Miami University-Hamilton, Hamilton, OH 45011, USA

⁴Environmental Resilience Institute, Indiana University, Bloomington, IN USA

*Corresponding Author Email: mjdietri@iu.edu

To be submitted to: *GeoHealth*

Abstract

Lead (Pb) is a neurotoxicant that particularly harms young children. Urban environments are often plagued with elevated Pb in soils and dusts, posing a health exposure risk from inhalation and ingestion of these contaminated media. Thus, a better understanding of where to prioritize risk screening and intervention is paramount from a public health perspective. We have synthesized a large national dataset of Pb concentrations in household dusts from across the United States (U.S), part of a community science initiative called “DustSafe.” Using these results, we have developed a simplistic logistic regression model that correctly predicts whether Pb is elevated (> 80 ppm) or low (< 80 ppm) in household dusts 75% of the time. Additionally, our model estimated 18% false negatives for elevated Pb, displaying that there was a low probability of elevated Pb in homes being misclassified. Our model uses only variables of approximate housing age and whether there is peeling paint in the interior of the home, illustrating how a simple and successful Pb predictive model can be generated if researchers ask the right screening questions. Scanning electron microscopy supports a common presence of Pb paint in several dust samples with elevated bulk Pb concentrations, which explains the predictive power of housing age and peeling paint in the model. This model was also implemented into an interactive mobile app that aims to increase community-wide participation with Pb household screening. The app will hopefully provide greater awareness of Pb risks and a highly efficient way to begin mitigation.

Plain Language Summary

Community science has been gaining traction in many locales throughout the United States, particularly in the field of urban pollution. While this has helped with science education and informing communities of potential hazards and mitigation tools, little has been done to effectively assimilate this information in a useful way to help other people in communities. Thus, we utilized a large dataset of household dust samples provided via community scientists across the United States to build a simple predictive model that lets users know if their dust is likely to be high in a toxic metal, lead. Additionally, we built this model into an interactive mobile app that we plan to use as a recruitment tool for usage of lead screening kits. Ultimately, we plan to assess whether this mobile app improves user knowledge of household lead risks and increases participation from start to finish for free lead screening services.

Key Points

- Community science sampling can provide national-level insight
- Mobile apps can be utilized as a lead intervention tool
- Elevated lead in house dust can be reasonably predicted with a simple model and two variables

1. Introduction

Lead (Pb) is a naturally occurring heavy metal neurotoxicant that causes many deleterious effects in humans, even in small quantities (e.g., Assi et al., 2016; Dórea, 2019). It is a biologically non-essential element that is especially detrimental to young children (e.g., Koller et al., 2004). In the United States (U.S.), it has largely been phased out of products, most notably leaded gasoline and paint, but remains in many urban environments as a form of legacy pollution (e.g., Laidlaw et al., 2012). Thus, modern sources of Pb are primarily lead paint in older homes and soil/dusts that contain remnants of both leaded paint and gasoline. Ingestion and inhalation of paint, soil, and dust containing elevated levels of Pb still poses a health risk, particularly for children due to their increased hand-to-mouth behavior (e.g., Ko et al., 2007; Needleman, 2004; Stewart et al., 2014).

Household dust Pb concentrations and loadings have been shown to be strongly related to children's blood Pb levels (BLLs) (e.g., Lanphear et al., 1996; Gulson and Taylor, 2017; Rhoads et al., 1999). Thus, a better understanding of risk factors associated with Pb in household dusts can help predict what homes may have elevated Pb concentrations in dusts, and thus help mitigate Pb exposure and elevated BLLs in children. Predictive modeling of Pb in soil samples with variables such as race and house age has already been shown to be effective in predicting at-risk areas (Obeng-Gyasi et al., 2021), but this has not been attempted with household indoor dust Pb concentrations across a wide geographic area through community-provided samples.

Citizen/community science sampling of environmental media such as soil has been shown to not only aid as an educational tool to those collecting the samples, but also provides important scientific data of inorganic contaminants such as Pb and how they're distributed throughout the environment (e.g., Filippelli et al., 2018; Ringwald et al., 2021; Taylor et al., 2021). Thus, we have utilized an ongoing community science project, "DustSafe (<https://www.360dustanalysis.com/>)," to analyze approximately 434 household dust samples from across the United States (Fig. 1) to determine whether homes at risk for elevated dust Pb can be accurately predicted. While individual variables such as housing age and automobile traffic near homes have been shown to be correlated with indoor dust Pb concentrations (e.g., Meyer et al., 1999; Rasmussen et al., 2011), variables have not been collectively applied in a predictive model across multiple states and cities in the U.S. Additionally, we sought to utilize this predictive model as part of an interactive mobile app to encourage greater community engagement for household Pb screening, which can not only help individuals gain agency in possible Pb mitigation measures, but can also help policymakers and the community at large better understand where/how to focus household Pb intervention efforts.

2. Methods

2.1 "DustSafe" sampling

Details of the household dust sampling are provided in Isley et al. (in review). Briefly, DustSafe was advertised as a program to thousands of households through social media, e-mail, etc. to gain community science participants. Project protocols were approved following ethical review at Indiana

University, USA (project #1810831960). Participants completed an online survey (Isley et al., in review—their SI Text 1) and collected vacuum cleaner dust in a polyethylene bag. Samples were collected from 2019 to present. Once samples were collected by researchers, they were sieved to 250 µm and analyzed for Pb, As, Cd, Cr, Cu, and Zn using X-ray fluorescence spectrometry (XRF). They were dry by virtue of the vacuum sampling and needed no desiccation. NIST 2702 was run periodically as an external standard on the XRF between dust samples, and the arithmetic mean (average) % error for Pb was 14.7% ± 8.6% (n = 9).

Results were reported back to participants following data collection (Example for Pb in Fig. S1), and then plotted on the “Map My Environment” website (www.mapmyenvironment.com) with locations randomly double jittered within ~2 city blocks to protect privacy.

2.2 Data filtering/building of logistic regression model

The initial dataset (link to data provided in Supplementary Materials) of potentially relevant data for this analysis contained 434 samples with matching Pb data (greater than detection limit) from the United States (and three samples from Canada). After deducing the most important potential predictive variables of housing age, interior peeling, exterior peeling, and recent renovation through log-normalized Pb concentration comparative t-tests, ANOVA tests and our global dust data (Isley et al., in review—their Table 1), the data was further filtered down to 342 samples that contained Pb concentrations and questionnaire responses for the aforementioned independent variables. Because exact housing age is difficult to deduce for many respondents, particularly renters and those who may be surveyed in-person at future community Pb screening events, we classified housing age into categories of Pre-1940, 1940-1959, 1960-1979, 1980-Present, and “Not Sure” so that this predictor variable may be more useful/applicable in future surveys.

A logistic regression model was applied using independent potential predictor variables to predict whether an indoor housing dust sample was either ≥80 ppm Pb or <80 ppm Pb. This was used as a conservative cut-off based on California’s safe screening level for soils, because we didn’t collect indoor dust loading data and most other standards used in the U.S. for soil Pb are outdated and likely too high (e.g., the U.S. EPA’s 400 ppm residential soil standard; Gailey et al., 2020). Our model was run in RStudio using the “glm” function based on the general equation:

$$\log \left[\frac{p}{1-p} \right] = b_0 + b_1 * x_1 + b_2 * x_2 \dots + b_n * x_n \quad (1)$$

Where p is the probability of an event occurring, b₀ is the intercept, b_n is the regression beta coefficient, and x_n is a given predictor variable.

Each potential independent predictor variable (besides housing age) categorical response of “No,” “Yes,” and “Not Sure” were reclassified as numeric variables of 0, 1, and 2, respectively, for the model. Housing age categories were reclassified as numeric variables of 0, 1, 2, 3, and 4 for the responses, “1980-Present,” “1960-1979,” “1940-1959,” “Pre-1940,” and “Not Sure,” respectively.

Our most successful model contained the independent variables of housing age (p = 0.0002) and interior peeling paint (p = 0.008), which generated the following equation:

$$\log \left[\frac{p}{1-p} \right] = 2.1413 - 0.4506 (Housing) - 1.1535 (Interior Paint Peeling) \quad (2)$$

This was based on a random training set of 240 samples from our original 342 samples. We evaluated the model on a testing dataset of 102 samples from our original 342 samples. All input and output files are

freely available on GitHub (link provided in Supplementary Materials), as well as the logistic regression model R code.

2.3 Mobile app development

An interactive online web application was developed to implement our predictive model in a simple and straightforward manner (link provided in Supplementary Materials). The application was built using the shiny, shinydashboard, shinydashboardPlus, and shinyjs packages in R (Attali, 2020; Chang et al., 2018; Chang and Borges Ribeiro, 2018; Granjon, 2021). Along with providing users with a straightforward interface for answering questions about house age and peeling paint and a custom risk assessment based on the embedded logistic predictive model, the application also provides users with direct links to our mapmyenvironment.com web portal, where they can register for free dust and soil Pb screening. Finally, the application offers background information about the current model version used to make the predictions, and offers direct links to model, data, and application code repositories.

2.4 Scanning electron microscopy (SEM)

A subset of DustSafe household dust samples were prepared on aluminum samples stubs using carbon sticky tab substrates for analysis using a scanning electron microscope (SEM) and energy dispersive X-ray spectroscopy (EDS). EDS lines used to identify Pb specifically include the $L_{\alpha} = 10.541$ keV (nominally $M_{\alpha} = 2.342$ keV, $M_{\beta} = 2.444$ keV). All analyses were conducted at Indiana University-Purdue University Indianapolis with a Zeiss EVO-10 SEM and Bruker XFlash6, 60 mm² EDS detector. Backscatter electron (BSE) images were collected at a setting of 15 kV in variable pressure mode. Qualitative elemental composition data (EDS data) were collected at the same conditions.

3. Results & Discussion

3.1 Significant findings between Pb in dust and housing age, vacuum frequency, and peeling paint

Household dust Pb concentrations were significantly higher in homes where there was interior or exterior paint peeling (Fig. 2; Table 1), which is in line with recent global household Pb dust data from the same DustSafe project (Isley et al., in review). This suggests that leaded paint is still a significant contributor of Pb to dust in many homes. However, it does not exclude outside sources such as soil/street dust that may include Pb from leaded gasoline. For example, indoor dusts have been shown to contain significant Pb sources from outdoor sources such as soils, dust, and industrial pollution as well (e.g., Adgate et al., 1998; Kelepertzis et al., 2020).

Greater housing age has long been known to be associated with increased Pb concentrations in household dusts, such as in Canada and the U.S. (e.g., Rasmussen et al., 2011; Rasmussen et al., 2013; Spalinger et al., 2007). Our results support this, as a moderate positive correlation was seen between housing age and Pb concentration in our samples (Fig. 3A), with more recent housing age categories generally lower in dust Pb as well (Fig. 3B; Table 1). This is most likely due to older homes containing Pb-based paints that can contribute to dust samples, as Pb housing paint was outlawed in the U.S. in 1978 and housing built before 1940 is the most likely to contain Pb paint (e.g., Levin et al., 2021). Furthermore, our global DustSafe dataset also observed a strong increase in Pb house dust concentration with home age (Isley et al., in review), suggesting that this is a common trend in many countries.

Regular cleaning of homes and the surrounding environment, including measures such as vacuuming, have been shown to effectively lower BLLs in children (e.g., Laidlaw et al., 2017; Rhoads et al., 1999). We also found that those vacuuming more frequently than once a month contained significantly lower concentrations of Pb in their house dust compared to those vacuuming monthly or less (Fig. S2A). However, we did not see any significant differences in Pb house dust concentrations in subcategories where people performed more than monthly vacuuming (Fig. S2B), which corresponds to our general trends in global dust data where increased vacuuming frequency was not associated with Pb dust concentration at all (Isley et al., in review). Our findings suggest that households that hardly vacuum may be more likely to accumulate Pb-rich larger particles when they do finally vacuum and gather samples, such as Pb-paint chips, which would skew the bulk chemistry Pb concentration to higher values (since we didn't measure loading rates). Households that more frequently vacuum may be less likely to sample larger, Pb-rich particles for their DustSafe sample submission.

3.2 Predictive accuracy of logistic regression model

Application of our logistic regression model on a “test” dataset of 102 samples from our original dataset reveals an overall prediction accuracy of 75% when using a probability threshold of 0.8 to determine “High” or “Low” Pb (Table 2). Importantly, only 4 samples out of 102 test samples (4%) were classified as “Low” Pb when they were actually a “High” Pb sample (Table 2). This implies that from an intervention standpoint our model contains few false negatives, and thus has excellent sensitivity (82%).

3.3 Usefulness and “App”lication of model for household Pb screening

While more sophisticated models can be effective in predicting high risk exposure areas for Pb in soils or dusts (e.g., Obeng-Gyasi et al., 2021), we believe that from a public health intervention standpoint, sometimes a simpler model is better. Because only two independent variables with categorical responses were proven statistically significant in our model and yielded an effective prediction accuracy of 75%, we decided to incorporate our model into a mobile-based app to aid in household Pb screening recruitment efforts (Fig. 4). The goal is to help people understand whether there is an increased chance of elevated Pb in their home based on our model, then give them an opportunity to freely test their home so that they can gain agency in decision-making regarding Pb mitigation. Additionally, we sought to include decision variables of “Not sure” in our app/model, because this helps with realistic in-person usage of the app at community events, and many people taking the survey may be renters and unsure of home age. Furthermore, renters are often one of the more likely subgroups of people to contain elevated household Pb in soil or dust (e.g., Masri et al., 2020). Within our model, approximately 28 individuals or 8% were uncertain of their exact home age (Fig. S3). Moving forward, it would be useful to include home ownership in our DustSafe surveys, to understand whether this is correlated to uncertainty in home age and the predictive power this has for elevated dust Pb.

Because our mobile app screening questions are simple, straightforward, and contain only categorical multiple-choice responses, we envision that its usage will be highly effective as a quick screening tool that many in-person events (i.e., community events, schools) can implement to help people know if Pb exposure is a hazard they should be concerned about. Furthermore, because our dataset is based on national-scale data, the mobile app can be utilized in many different locations, further aiding in its “app”licability and versatility as a Pb screening recruitment tool.

3.4 Evidence of Pb-paint in dust samples

Through SEM work on several household dust samples that contained elevated bulk Pb concentrations, we were able to identify numerous examples of particles consistent in composition and morphology to Pb paint, ranging from ~10 μm in diameter to >100 μm in diameter (Fig. 5). Our Pb paint chips were similar to Pb paint analyzed by SEM in Hunt (2016), including several Pb-carbonate paints and the presence of Zn in the paint (Fig. 5). Additionally, the Mg-Al-Si EDS peaks in several paint samples (i.e., Figs. S6, S7, S8) are consistent with montmorillonite, an additive commonly used in Pb-based paint as organo-clays to aid in the suspension of the pigments. This helps explain why the predictor variables of housing age and interior peeling paint were so significant—many household dust samples with elevated concentrations of Pb likely have the Pb predominantly sourced from house paint. However, this does not mean that Pb in house dusts is exclusively from house paint, or that other metals are from exclusively indoor sources. As mentioned earlier, outdoor sources of pollutants can enter homes, such as through dust brought indoors (e.g., Adgate et al., 1998; Kelepertzis et al., 2020). For example, we found clear examples of technogenic Fe-oxide spherules, likely a byproduct of anthropogenic combustion, in house dust samples (Fig. S4). These particles likely came from an outdoor source, such as vehicle exhaust or industrial combustion, as they are similar to Fe-rich spherical particles commonly found in industrial areas from high temperature formation processes (e.g., Dietrich et al., 2019; Gaberšek and Gosar, 2021; Miler and Gosar, 2013; Teran et al., 2020). Furthermore, we found one sample that contains EDS spectra consistent with PbCrO_4 , or Pb-chromate paint (Fig. 5A), which could have come from yellow-paint inside the home, but may have also been brought in from outdoors where Pb-chromate is often used in traffic paint (e.g., O'Shea et al., 2021).

3.5 Future goals and directions

We based our initial model on predominantly U.S. house dust samples, because of statistically significant differences in bulk metal composition of dusts between other countries (Isley et al., in review) and there are likely other confounding factors between countries that affect Pb in dusts (i.e., different regulation of Pb paints and Pb gasoline). However, as more data is collected and as we gain a better understanding of what variables predominantly influence Pb in house dust, our model can be applied to additional countries and refined within the U.S. to more accurately differentiate what homes likely contain elevated Pb. A specific area for refinement of the model may lie in spatial data, such as relating zip codes of samples with socioeconomic (i.e., % poverty, racial distribution) and public health data (i.e., blood lead levels) within those zip codes, which may add to the predictive power of our model.

Additionally, this type of simplistic predictive model usage in a mobile app as an intervention tool can be applied beyond Pb in household dusts, such as other contaminants of concern in homes like arsenic (As) or radon (Rn). Lastly, community science sampling endeavors should continue to grow, as they are not only a great opportunity for direct household contamination intervention, but also contribute to a greater general understanding of important issues such as Pb pollution and what areas community remediation should be focused in. Scientific information **from the public** is one of the easiest ways to **help the public** in pollution remediation, as we have shown with our basic Pb dust logistic regression model and mobile app.

We plan to conduct a follow-up study on the effectiveness of this type of simple intervention in engaging participants to have full-cycle involvement, going from initial completion of app input to

submittal of samples to opening results once generated. Ample examples of citizen science exist with various ways that the engagement does, or does not, provide real, tangible benefits to participants (e.g., Hayhow et al., 2021), but they are typically poorly assessed. We hypothesize that this simple app engagement will generate greater “engage to completion” metrics because of simplicity of message. We will also develop a back-end follow-up survey once the sample results are generated and returned to users to determine what, if any, impacts the data had on participant’s behavior, including any mitigation steps that they took in response to results.

4. Conclusions

A simple logistic regression model based on real-world samples proved to be effective at identifying homes at risk for higher Pb in household dusts across the United States. Application of the model on a test dataset of 102 samples revealed a 75% classification accuracy of either “high” or “low” Pb in household dust, with the cutoff based on 80 ppm Pb. This illustrates how community science gathered data can provide valuable insight into primary predictor variables for elevated Pb. Additionally, we showed how simplistic, yet effective Pb predictive models can be incorporated into interactive mobile apps such as a Pb screening recruitment tool. Collectively, we hope that modeling efforts such as these and engagement with local communities will aid in Pb exposure prevention and remediation, so that no child grows up with an unnecessarily high risk of Pb exposure.

Acknowledgements

The authors are deeply thankful to the households that supplied dust samples for this work and the lab techs who helped process samples. The MapMyEnvironment program and the related DustSafe sampling effort are partially supported by National Science Foundation Grant ICER-1701132 to G.M.F. and the Environmental Resilience Institute, funded by Indiana University’s Prepared for Environmental Change Grand Challenge Initiative. Partial support for this work was provided by NSF-EAR-PF Award #2052589 to M.D. Special thanks to Miguel Cruz for help with SEM instrumentation, Leah Wood and Angela Herrmann for input on mobile app development, and Michael O’Shea with input on SEM image interpretation.

Open Research

All data and source code used in this manuscript are freely available on GitHub (<https://github.com/dietrimj/Community-Science-Pb-Prediction>).

Tables:

Table 1: Summary statistics of household dust Pb concentrations (mg/kg)

		<i>Mean</i>	<i>Std Dev</i>	<i>Median</i>	<i>Max</i>	<i>Min</i>	<i>(n)</i>
Total Pb		99	239	32	2328	3	434
Exterior Paint Peeling	Yes	131	179	41	815	4	48
	No	94	248	30	2328	3	272
	Not Sure	40	46	28	205	5	23
Interior Paint Peeling	Yes	142	175	81	729	7	40
	No	91	240	30	2328	4	302
	Not Sure	35	N/A	35	35	35	1
Housing Age	Pre-1940	228	306	134	1665	7	54
	1940-1959	121	221	53	1304	10	33
	1960-1979	78	193	32	1377	6	52
	1980-Present	44	113	24	1205	3	178
	Not Sure	114	319	32	2328	5	117

Table 2: Confusion matrix output of logistic regression model from test dataset (n = 102)

	Actual High Pb	Actual Low Pb
Predicted High Pb	18	21
Predicted Low Pb	4	59

303 **Figures:**

304




Figure 1 Samples with Pb (and other heavy metal) results reported back to households from the "DustSafe" project in the U.S. and Canada (samples also taken from Hawaii).

305




Figure 2 Embedded boxplots within violin plots for both interior (A) and exterior peeling paint (B) questionnaire responses. The boxes represent the interquartile range (IQR) of 25th-75th percentiles of data, the horizontal line is the median, and the whiskers represent 1.5 times the IQR. Two-sample paired t-test results between yes/no responses are also provided. The y-axes are transformed on a log₁₀ scale, and the dashed red lines represent California's safe screening soil Pb level of 80 ppm.




Figure 3 (A) Scatterplot between approximate housing ages and \log_{10} Pb concentrations with the Pearson correlation coefficient and associated p-value provided, as well as a linear regression line in blue with the shaded 95% confidence interval. (B) Embedded boxplots within violin plots for housing age categories used in the predictive model. The boxes represent the interquartile range (IQR) of 25th-75th percentiles of data, the horizontal line is the median (which is connected between housing age categories with a black line), and the whiskers represent 1.5 times the IQR. An analysis of variance (ANOVA) test associated p-value between all housing age categories is provided. The y-axis is transformed on a \log_{10} scale, and the dashed red line represents California's safe screening soil Pb level of 80 ppm.

360

361

362

363




Figure 4 Screenshots from the beginning of the interactive Pb household dust screening app (<https://iupui-earth-science.shinyapps.io/IUPUI-LeadRiskApp/>).




Figure 5 SEM images of particles resembling Pb paint, surrounded by other particulates in various high Pb DustSafe household dust samples (corresponding EDS spectra provided in Supplementary Materials; Figs. S5-S10). Pb paint particles are evident by very high contrast of electron backscatter detection—more so than surrounding particles because of the high atomic number of Pb. Most Pb-bearing particles are angular or jagged, with clear flaky particles on their surface.

References

- Adgate, J. L., Rhoads, G. G., & Liroy, P. J. (1998). The use of isotope ratios to apportion sources of lead in Jersey City, NJ, house dust wipe samples. *Science of the Total Environment*, 221(2-3), 171-180.
- Assi, M. A., Hezmee, M. N. M., Abd Wahid Haron, M. Y. M., & Sabri, M. A. R. (2016). The detrimental effects of lead on human and animal health. *Veterinary World*, 9(6), 660.
- Attali, D. (2020). shinyjs: Easily Improve the User Experience of Your Shiny Apps in Seconds. R package version 2.0.0. <https://CRAN.R-project.org/package=shinyjs>
- Chang, W., Cheng, J., Allaire, J.J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., Borges, B. (2021). shiny: Web Application Framework for R. R package version 1.6.0. <https://CRAN.R-project.org/package=shiny>
- Chang, W. & Borges Ribeiro, B. (2018). shinydashboard: Create Dashboards with 'Shiny'. R package version 0.7.1. <https://CRAN.R-project.org/package=shinydashboard>
- Dietrich, M., Wolfe, A., Burke, M., & Krekeler, M. P. (2019). The first pollution investigation of road sediment in Gary, Indiana: Anthropogenic metals and possible health implications for a socioeconomically disadvantaged area. *Environment International*, 128, 175-192.
- Dórea, J. G. (2019). Environmental exposure to low-level lead (Pb) co-occurring with other neurotoxicants in early life and neurodevelopment of children. *Environmental Research*, 177, 108641.
- Filippelli, G. M., Adamic, J., Nichols, D., Shukle, J., & Frix, E. (2018). Mapping the urban lead exposome: a detailed analysis of soil metal concentrations at the household scale using citizen science. *International Journal of Environmental Research and Public Health*, 15(7), 1531.
- Gaberšek, M., & Gosar, M. (2021). Towards a holistic approach to the geochemistry of solid inorganic particles in the urban environment. *Science of the Total Environment*, 763, 144214.
- Gailey, A. D., Schachter, A. E., Egendorf, S. P., & Mielke, H. W. (2020). Quantifying soil contamination and identifying interventions to limit health risks. *Current Problems in Pediatric and Adolescent Health Care*, 50(1), 100740.
- Granjon, D. (2021). shinydashboardPlus: Add More 'AdminLTE2' Components to 'shinydashboard'. R package version 2.0.2. <https://CRAN.R-project.org/package=shinydashboardPlus>
- Gulson, B., & Taylor, A. (2017). A simple lead dust fall method predicts children's blood lead level: New evidence from Australia. *Environmental Research*, 159, 76-81.
- Hayhow, C., Brabander, D.J., Jim, R., Lively, M., Filippelli, G. (2021). "We have been the ladder and held the ladder": Evolving GeoHealth models for actionable, community-engaged research. *Earth and Space Science Open Archive*. <https://doi.org/10.1002/essoar.10507853.1>
- Hunt, A. (2016). Relative bioaccessibility of Pb-based paint in soil. *Environmental Geochemistry and Health*, 38(4), 1037-1050.
- Isley et al. (in review). Global analysis of sources and human health risk associated with trace metal contaminants in residential indoor dust. *Environmental Science & Technology*.

401 Kelepertzis, E., Argyraki, A., Chrastný, V., Botsou, F., Skordas, K., Komárek, M., & Fouskas, A. (2020).
 402 Metal (loid) and isotopic tracing of Pb in soils, road and house dusts from the industrial area of Volos
 403 (central Greece). *Science of The Total Environment*, 725, 138300.

404 Ko, S., Schaefer, P. D., Vicario, C. M., & Binns, H. J. (2007). Relationships of video assessments of
 405 touching and mouthing behaviors during outdoor play in urban residential yards to parental perceptions of
 406 child behaviors and blood lead levels. *Journal of Exposure Science & Environmental*
 407 *Epidemiology*, 17(1), 47-57.

408 Koller, K., Brown, T., Spurgeon, A., & Levy, L. (2004). Recent developments in low-level lead exposure
 409 and intellectual impairment in children. *Environmental Health Perspectives*, 112(9), 987-994.

410 Laidlaw, M. A., Zahran, S., Mielke, H. W., Taylor, M. P., & Filippelli, G. M. (2012). Re-suspension of
 411 lead contaminated urban soil as a dominant source of atmospheric lead in Birmingham, Chicago, Detroit
 412 and Pittsburgh, USA. *Atmospheric Environment*, 49, 302-310.

413 Laidlaw, M.A., Filippelli, G.M., Brown, S., Paz-Ferreiro, J., Reichman, S.M., Netherway, P.,
 414 Truskewycz, A., Ball, A.S. and Mielke, H.W. (2017). Case studies and evidence-based approaches to
 415 addressing urban soil lead contamination. *Applied Geochemistry*, 83, 14-30.

416 Lanphear, B.P., Weitzman, M., Winter, N.L., Eberly, S., Yakir, B., Tanner, M., Emond, M. and Matte,
 417 T.D. (1996). Lead-contaminated house dust and urban children's blood lead levels. *American Journal of*
 418 *Public Health*, 86(10), 1416-1421.

419 Levin, R., Vieira, C. L. Z., Rosenbaum, M. H., Bischoff, K., Mordarski, D. C., & Brown, M. J. (2021).
 420 The urban lead (Pb) burden in humans, animals and the natural environment. *Environmental Research*,
 421 110377.

422 Masri, S., LeBrón, A., Logue, M., Valencia, E., Ruiz, A., Reyes, A., Lawrence, J.M. and Wu, J. (2020).
 423 Social and spatial distribution of soil lead concentrations in the City of Santa Ana, California:
 424 Implications for health inequities. *Science of The Total Environment*, 743, 140764.

425 Meyer, I., Heinrich, J., & Lippold, U. (1999). Factors affecting lead, cadmium, and arsenic levels in house
 426 dust in a smelter town in eastern Germany. *Environmental Research*, 81(1), 32-44.

427 Miler, M., & Gosar, M. (2013). Assessment of metal pollution sources by SEM/EDS analysis of solid
 428 particles in snow: a case study of Žerjav, Slovenia. *Microscopy and Microanalysis*, 19(6), 1606-1619.

429 Needleman, H. (2004). Lead poisoning. *Annual Review of Medicine*, 55, 209-222.

430 Obeng-Gyasi, E., Roostaei, J., & Gibson, J. M. (2021). Lead Distribution in Urban Soil in a Medium-
 431 Sized City: Household-Scale Analysis. *Environmental Science & Technology*, 55(6), 3696-3705.

432 O'Shea, M. J., Vigliaturo, R., Choi, J. K., McKeon, T. P., Krekeler, M. P., & Gieré, R. (2021). Alteration
 433 of yellow traffic paint in simulated environmental and biological fluids. *Science of the Total Environment*,
 434 750, 141202.

435 Rasmussen, P.E., Beauchemin, S., Chénier, M., Levesque, C., MacLean, L.C., Marro, L., Jones-Otazo,
 436 H., Petrovic, S., McDonald, L.T. and Gardner, H.D. (2011). Canadian house dust study: lead
 437 bioaccessibility and speciation. *Environmental Science & Technology*, 45(11), 4959-4965.

438 Rasmussen, P. E., Levesque, C., Chénier, M., Gardner, H. D., Jones-Otazo, H., & Petrovic, S. (2013).
 439 Canadian House Dust Study: Population-based concentrations, loads and loading rates of arsenic,

440 cadmium, chromium, copper, nickel, lead, and zinc inside urban homes. *Science of the Total*
441 *Environment*, 443, 520-529.

442 Rhoads, G. G., Ettinger, A. S., Weisel, C. P., Buckley, T. J., Goldman, K. D., Adgate, J., & Liroy, P. J.
443 (1999). The effect of dust lead control on blood lead in toddlers: a randomized trial. *Pediatrics*, 103(3),
444 551-555.

445 Ringwald, P., Chapin, C., Iceman, C., Tighe, M.E., Sisk, M., Peaslee, G.F., Peller, J. and Wells, E.M.
446 (2021). Characterization and within-site variation of environmental metal concentrations around a
447 contaminated site using a community-engaged approach. *Chemosphere*, 272, 129915.

448 Spalinger, S. M., von Braun, M. C., Petrosyan, V., & von Lindern, I. H. (2007). Northern Idaho house
449 dust and soil lead levels compared to the Bunker Hill Superfund site. *Environmental Monitoring and*
450 *Assessment*, 130(1), 57-72.

451 Stewart, L. R., Farver, J. R., Gorsevski, P. V., & Miner, J. G. (2014). Spatial prediction of blood lead
452 levels in children in Toledo, OH using fuzzy sets and the site-specific IEUBK model. *Applied*
453 *Geochemistry*, 45, 120-129.

454 Taylor, M.P., Isley, C.F., Fry, K.L., Liu, X., Gillings, M.M., Rouillon, M., Soltani, N.S., Gore, D.B. and
455 Filippelli, G.M. (2021). A citizen science approach to identifying trace metal contamination risks in urban
456 gardens. *Environment International*, 155, 106582.

457 Teran, K., Žibret, G., & Fanetti, M. (2020). Impact of urbanization and steel mill emissions on elemental
458 composition of street dust and corresponding particle characterization. *Journal of Hazardous*
459 *Materials*, 384, 120963.