


Numerically Modeling Infiltration and Root Water Uptake in a Montane Forest using High Frequency Stable Isotope Field Measurements

Raymond Hess¹, Matthias Sprenger², Ali Zahori³, Nathaniel Bogie¹¹San José State University, Department of Geology, San José, CA²Lawrence Berkeley National Laboratory, Earth & Environmental Sciences Area, Berkeley, CA³San José State University, Department of Environmental Studies, San José, CA

Background




- In the coming years, climate models forecast mountainous watersheds to undergo a reduction in snowpack, early season melt, and increases in evapotranspiration.
- As a result, dry soil conditions will stress vegetation at elevations of 1,850 to 2,900 meters above sea level.
- We investigate infiltration patterns and root water uptake in response to drying within the East River catchment in Colorado.
- This study highlights the benefits of coupling innovative high frequency tracer data and commonly used hydrometric data to better constrain parameters used in numerical modeling.



Research Objective & Methods

Observe and model water dynamics in soil beneath Engelmann spruce and Aspen profiles.

Instrumentation: Pressure head and sap flow sensors, vapor probes (24), daily measurements of concentration recorded using an in situ vapor analyzer.




Vapor probe schematic


Seeger & Weiler, 2021

- We collected soil cores, measured matric potential, sap flow, and monitored tree xylem and soil for stable isotopes of water (²H, ¹⁸O) along two profiles to 90 cm depth (three Engelmann spruce and three aspen trees).
- Field isotope dynamics were analyzed on a daily basis between mid-July and late October using an in situ cavity ring-down spectrometer.
- The numerical model HYDRUS-1D was trained and calibrated with pressure head and isotope data, simulating the response to late summer dry spells and monsoonal rainfall across the 128 day period.

Field Setup




Conceptual Approach




Model Optimization & Goodness of Fit

Parameter Optimization for Aspen Model



Sensitivity Analysis for Aspen Interception




- Plots showing Aspen model runs for parameter optimization and sensitivity analysis for interception parameter (30,000 runs).
- A multi-objective approach and Monte Carlo analysis was used to identify a best parameter set for the model.
- The Kling-Gupta Efficiency (KGE) was utilized as a goodness of fit metric, with a range of $-inf$ (poorest fit) to 1 (best possible fit).
- KGE_{iso} : Goodness of fit for the of stable isotope time series.
- KGE_h : Goodness of fit for the of pressure head time series.


Parameters Calibrated

θ_s : Saturated Water Content
 α : Aspect $\approx [1 / (\text{Air Entry})]$
 n : Pore-size distribution
 K_s : Saturated hydraulic conductivity
 D_L : Dispersivity
 I : Interception

Simulating Infiltration and Uptake Dynamics



- Simulated $\delta^2\text{H}$ compared with in-situ $\delta^2\text{H}$.**
- Observations plotted as circles.
- Simulations plotted as curves
- Red:** $\delta^2\text{H}$ concentration in the Aspen soil profile
- Green:** $\delta^2\text{H}$ concentration in Aspen trees
- (KGE_{iso} : 0.24)



- Modeled h (pressure heads, matric potential)**
- Observed data plotted as circles
- Modeled data plotted as curves
- Tan:** 15 cm, (KGE : 0.08)
- Orange:** 30 cm, (KGE : -0.04)
- Red:** 60 cm (KGE : 0.77)
- KGE_h : 0.32.

Results & Conclusions

- Model simulations roughly show the three distinct dry down events from late July to September, each followed by rapid infiltration of rainfall (42, 47, and 68 mm of total precipitation per event).
- Compared to Aspen trees, shallow soil under Engelmann spruce repeatedly dries out beyond the permanent wilting point of -1.5 MPa (higher rates of canopy interception).
- That said, these efforts aim to help predict and better understand quantification of certain plant water responses during ecosystem changes and future climate conditions (i.e. less frequent monsoonal rains and even fewer high magnitude rain events).

Next Steps

- Further optimization and sensitivity analysis for best fit at shallow depths.
- Model and compare isotope dynamics for both WY-22 and WY-23
- Simulate infiltration and uptake under high ET and low P conditions.

Acknowledgements

We thank Ginny Smith and Alphonse Odisho for technical support, and Leslie Blum for administrative support. This work was supported in part by the U.S. Department of Energy, Office of Science, Office of Workforce Development for Teachers and Scientists (WDTS) under the Visiting Faculty Program (VFP). This material is based upon work supported as part of the Watershed Function Scientific Focus Area and the Laboratory Directed Research and Development Program of Lawrence Berkeley National Laboratory "Latest-Iso" funded by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-AC02-05CH11231.